

Figure S1. Related to Figure 1. Clumping of GWAS results from NHGRI-EBI GWAS catalog. For each GWAS traits, we performed LD clumping to only keep the lead SNP in each region.




Figure S2. Histogram of shared SNPs for each trait pair in NHGRI-EBI GWAS catalog from iCPAGdb at false discovery rate of 0.1. Among 76127 trait pairs (including compound phenotypes), the mean number of shared independent SNPs was 3.10, and 38.8% of trait pairs shared more than 1 SNP. "Mathematical ability" and "self reported educational attainment" shared the most SNPs with 739.




Figure S3. Related to Figure 1. Comparison of CPAG1 and iCPAGdb using the NHGRI-EBI GWAS catalog summary statistics downloaded on September 4, 2013. CPAG1 finished this analysis in 73.2 minutes while iCPAGdb used 4.1 minutes on a MacBook with Quad-Core intel i7 CPU and 16 GBs of RAM. CPAG1 cannot run efficiently enough to allow for comparison with the 2020 NHGRI-EBI GWAS catalog (estimated completion time > ~10 days for CPAG vs. ~100 minutes for iCPAGdb). A total of 887 traits were included in the dataset, and p values were corrected using multiple test correction for 392,941 comparisons. A) iCPAGdb detected 2598 cross-phenotype associations, 37% more than CPAG1 at FDR of 0.1. B) The number of cross-phenotype associations from directly shared SNPs were nearly unchanged between iCPAGdb and CPAG1 but indirect associations increased 76%.

- **Table S1.** Related to Figure 1. iCPAGdb output for cross-phenotype associations from NHGRI-EBI GWAS catalog (downloaded on August 5, 2020).
- **Table S2.** Related to Figure 2. iCPAGdb output for cross-phenotype associations between molecular and cellular datasets and NHGRI-EBI GWAS catalog.
- **Table S3.** Related to Figure 2. COLOC analysis of *CXCL10* following *C. trachomatis* infection and levels of *CXCL9* (MIG) in whole blood.
- **Table S4.** Related to Figure 3. Cross-phenotype associations for COVID-19 (p value < 1 x 10^{-5}) against 4414 GWAS traits in iCPAGdb (p value < 5×10^{-8}).
- **Table S5.** Related to Figure 3. COLOC analysis output for *ABO* region between COVID-19 and CD209 antigen levels.
- **Table S6.** Related to Figure 4. COLOC analysis output for *DPP9* region between COVID-19 and idiopathic pulmonary fibrosis.