

Supplemental material belonging to:


Drinking habits and executive functioning: a propensity score-weighted analysis of 78,832 adults

Schweren, Haavik, Li, Solberg, Larsson & Hartman. 2020

Contents:

1. Participant flowchart
2. Heatmap: propensity scores vs. all potential confounders
3. Balance before and after propensity score weighing
 - i. Means by drinking level before and after weighing
 - ii. Means by binge-drinking (yes/no) before and after weighing
 - iii. Adjustable and non-adjustable differences between weighted groups
4. Results among participants of white ethnicity only
5. Results stratified by age and sex
6. Results stratified by drink type


Supplement 1: Participant flowchart



Supplemental Figure S1. Participant flowchart.

RFFT = Ruff Figural Fluency Task; MMSE = Mini Mental State Examination; ¹ For logistical reasons, the RFFT was administered in all Lifelines participants until April 2012, and in a random half of all participants thereafter, resulting in a relatively large proportion of missing data; ² By protocol, the RFFT was not administered in participants with MMSE<26. Moreover, MMSE was only administered in participants age 65 and older. Thus, the vast majority of participants with MMSE<26 were already excluded in a previous step.

Supplement 2: Correlations between propensity scores and all potential confounders



Supplemental Figure S2. Heatmap of correlations between all confounders and each propensity score.

Supplement 3: Balance before and after weighing

Supplemental Table S3.1. Means by drinking level before and after weighing

	Before weighing					After weighing				
	abst.	occ.	light	mod.	heavy	abst.	occ.	light	mod.	heavy
Age	42.9	41.7	42.6	45.1	44.4	43.2	42.8	42.9	43.5	44.4*
Income	2309.8	2407.9	2498.1	2543.6	2415.4	2432.6	2442.1	2448.2	2469.6	2505.2
Income equivalised	1483.3	1561.1	1636.1	1694.3	1643.2	1586.7	1589.1	1598.5	1618.0	1665.4*
Neighbourhood status	-0.6	-0.6	-0.6	-0.6	-0.7	-0.6	-0.6	-0.6	-0.6	-0.6
Occupational status	42.4	46.0	47.8	47.8	43.5	46.0	46.1	46.3	46.7	46.4
BMI	26.7	26.0	25.7	25.9	26.8	26.1	26.0	26.0	25.9	26.1
Diet quality	23.7	23.9	23.9	23.0	20.4	23.7	23.7	23.7	23.6	23.1*
Physical activity	249.3	259.1	275.0	270.2	254.9	259.1	260.9	264.3	257.4	259.8
SLE	1.3	1.2	1.1	1.2	1.4	1.2	1.2	1.2	1.2	1.2
LTD	2.7	2.7	2.5	2.4	2.6	2.6	2.6	2.6	2.6	2.5
Impulsivity	22.1	22.3	22.4	22.6	23.2	22.3	22.3	22.4	22.4	22.6
Excitement seeking	20.5	21.8	22.9	23.2	23.9	22.0	22.1	22.3	22.3	22.6*
Self-discipline	29.2	29.1	29.4	29.4	29.0	29.3	29.3	29.3	29.3	29.5
Social contacts	16.7	16.7	18.1	18.2	18.7	17.3	17.4	17.5	17.3	17.9
Male sex(%)	19.4	31.8	48.3	63.5	82.2	41.6	40.0	42.1	45.4	51.3*
Nonwhite ethnicity (%)	2.1	1.8	1.1	0.9	0.5	1.6	1.5	1.4	1.1	1.1
Low education (%)	34.9	26.6	25.6	29.7	37.5	29.3	28.6	28.4	28.8	29.6
High education (%)	22.7	30.7	22.8	32.5	23.2	28.9	30.2	30.6	31.2	30.2
Cancer (%)	3.7	3.8	3.6	3.7	3.5	3.7	3.8	3.8	3.5	4.7
CVD (%)	34.3	28.5	28.6	32.4	37.1	31.2	30.3	30.3	30.6	33.3
Diabetes (%)	2.9	1.8	1.4	1.7	1.8	2.1	1.9	1.8	2.2	1.6
Liver cirrhosis (%)	2.5	1.8	1.7	1.9	2.2	2.0	1.9	1.8	1.9	1.3
Depression (%)	5.0	3.2	2.5	2.5	3.3	3.3	3.1	3.1	3.1	2.9
Anxiety (%)	10.7	8.3	6.9	7.3	8.1	8.2	8.1	7.9	8.7	7.7
Fam addiction (%)	5.6	4.9	5.3	6.0	7.3	5.8	5.3	5.4	5.7	5.6
Fam dep/anx (%)	27.0	26.8	26.1	25.0	24.1	26.2	26.4	26.1	25.7	26.7
Fam CVD (%)	10.7	9.4	8.9	9.2	8.9	9.9	9.5	9.4	8.9	9.7

Past smoking (%)	25.8	28.1	34.3	40.0	34.5	30.9	31.8	32.1	33.8	36.0*
Current smoking (%)	17.7	17.1	22.1	33.3	44.9	20.4	21.3	22.4	23.4	27.2*
Short sleep (%)	11.0	9.7	9.2	10.9	13.7	10.3	10.1	10.1	10.7	10.0
Long sleep (%)	12.0	10.8	8.8	8.8	10.4	10.2	10.1	9.7	9.3	9.6

* adjustable difference between indicated group and abstinent group after weighing. Abbreviations: abst. = abstinent, occ. = occasional drinker, mod. = moderate drinker.


Supplemental Table S3.2. Means by binge-drinking (yes/no) before and after weighing

	Before weighing		After weighing	
	non-binge-drinking	binge-drinking	non-binge-drinking	binge-drinking
Age	43.5	36.8	42.8	42.6
Income	2466.3	2224.1	2442.9	2444.1
Income equivalised	1597.5	1557.0	1593.6	1608.2
Neighbourhood status	-0.6	-0.8	-0.6	-0.6
Occupational status	46.6	42.2	46.2	45.5
BMI	26.0	26.2	26.0	26.1
Diet quality	24.0	21.0	23.7	23.4
Physical activity	264.2	270.6	264.6	267.4
SLE	1.2	1.3	1.2	1.2
LTD	2.6	2.8	2.6	2.6
Impulsivity	22.2	23.6	22.4	22.5
Excitement seeking	21.9	24.9	22.2	22.5
Self-discipline	29.3	28.7	29.3	29.4
Social contacts	17.4	18.3	17.5	17.2
Male sex(%)	39.0	57.4	40.8	43.3
Nonwhite ethnicity (%)	1.5	1.2	1.5	1.4
Low education (%)	28.2	32.4	28.6	29.3
High education (%)	31.1	23.2	30.3	28.1
Cancer (%)	3.8	2.5	3.7	3.8
CVD (%)	31.2	23.9	30.5	30.3
Diabetes (%)	1.9	1.3	1.9	1.7
Liver cirrhosis (%)	2.0	1.4	1.9	1.6
Depression (%)	3.1	3.7	3.2	3.1

Anxiety (%)	8.1	7.9	8.1	8.3
Fam addiction (%)	5.4	5.5	5.4	5.7
Fam dep/anx (%)	26.8	21.8	26.3	26.0
Fam CVD (%)	9.6	8.0	9.5	9.0
Past smoking (%)	32.6	24.4	31.8	33.9
Current smoking (%)	19.8	39.2	21.8	23.9
Short sleep (%)	10.0	10.3	10.0	9.6
Long sleep (%)	10.0	9.5	10.0	9.9

Supplemental Table S3.3. Adjustable and non-adjustable differences between weighted groups

	Contrast	Confounder	Group 1	Group 2	ASMD
Adjustable	Abstinent vs. moderate	Smoking: never	48.7%	42.8%	0.1185
	Abstinent vs. heavy	Age	43.2	44.4	0.1103
	Abstinent vs. heavy	Sex: male	41.6%	51.3%	0.1946
	Abstinent vs. heavy	Income equivalised	1586.7	1665.4	0.1553
	Abstinent vs. heavy	Diet quality	23.7	23.1	0.1148
	Abstinent vs. heavy	Smoking: never	48.7%	36.8%	0.2385
	Abstinent vs. heavy	Smoking: past	30.9%	36.0%	0.1066
	Abstinent vs. heavy	Smoking: current	20.4%	27.2%	0.1614
	Abstinent vs. heavy	Excitement-seeking	22.0	22.6	0.1404
Non-adjustable	<i>None</i>				



Supplemental Figure S3.4. Absolute standardised mean differences (ASMD) in pairwise between-group contrasts (e.g. $ASMD_{AGE}$ abstinent vs. light drinkers, $ASMD_{AGE}$ abstinent vs. moderate drinkers, etc.) before (red circle) and after (blue triangle) propensity score weighing. On the X-axis, contrasts are ordered by their pre-weighting ASMD. Dashed horizontal lines indicate acceptable ($ASMD < 0.1$) and adjustable ($ASMD < 0.25$) residual imbalance after weighing. No residual between-group difference exceeds $ASMD = 0.25$.

Supplement 4: Results among participants of white ethnicity only

Supplemental Table S4.1. Adjustable and non-adjustable differences between weighted groups

	Contrast	Confounder	Group 1	Group 2	ASMD
Adjustable	Abstinent vs. moderate	Smoking: never	48.4%	43.3%	0.1016
	Abstinent vs. heavy	Diet quality	23.7	23.0	0.1320
	Abstinent vs. heavy	Smoking: never	48.4%	40.9%	0.1502
	Abstinent vs. heavy	Smoking: current	20.6%	25.5%	0.1165
	Abstinent vs. heavy	Excitement-seeking	21.9	22.7	0.1549
Non-adjustable	<i>None</i>				

Supplemental Table S4.2. Propensity weighted models comparing abstinent participants to each drinking level, and binge-drinkers to non-binge-drinkers

	Non-standardised		Standardised		p
	Est	95% CI	β	95% CI	
Occasional (0-2.5 g/day) vs. abstinent	0.343	-0.179 – 0.864	0.015	-0.008 – 0.038	0.1974
Light (2.5-14.9 g/day) vs. abstinent	0.849	0.389 – 1.318	0.038	0.017 – 0.058	0.0004
Moderate (15-29.9 g/day) vs. abstinent	1.793	1.122 – 2.465	0.079	0.050 – 0.109	<0.0001
Heavy (30-60 g/day) vs. abstinent	0.789	-0.977 – 2.736	0.039	-0.043 – 0.121	0.3531
Binge-drinking yes vs no	0.233	-0.528 – 0.994	0.010	-0.023 – 0.044	0.5479

Supplement 5: Results stratified by age and sex

Supplemental Table S5.1. Age- and sex-stratified groups

Sex	Age	N	%
Male	Young (18-38)	10,392	13.2
	Middle (39-48)	11,999	15.2
	Old (49-65)	9,876	12.5
Female	Young (18-38)	15,442	19.6
	Middle (39-48)	17,287	21.9
	Old (49-65)	13,836	17.6

Supplemental Table S5.2. Adjustable and non-adjustable differences between weighted groups

	Group	Contrast	Confounder	Group 1	Group 2	ASMD	
Adjustable	Male – young	Abst. vs. light	Excitement-seeking	25.1	25.6	0.1217	
		Abst. vs. moderate	Smoking: never	55.3%	48.4%	0.1377	
		Abst. vs. moderate	Excitement-seeking	25.1	25.7	0.1582	
		Abst. vs. heavy	Ethnicity: white	98.4%	99.7%	0.1267	
	Male – middle	Abst. vs. heavy	Ethnicity: white	98.7%	99.9%	0.1114	
		Abst. vs. heavy	Excitement-seeking	23.2	23.7	0.1172	
	Male – old	Abst. vs. moderate	Excitement-seeking	21.5	21.9	0.1003	
		Abst. vs. heavy	Ethnicity: white	98.6%	99.8%	0.1252	
		Abst. vs. heavy	Family: addiction	5.3%	3.1%	0.1029	
		Abst. vs. heavy	Excitement-seeking	21.5	22.2	0.1688	
		Female – young	Abst. vs. light	Excitement-seeking	22.9	23.3	0.1019
		Abst. vs. moderate	Education: middle	49.1%	43.7%	0.1085	
		Abst. vs. moderate	Education: high	36.1%	43.7%	0.1532	
		Abst. vs. moderate	BMI	25.1	24.5	0.1281	
		Abst. vs. moderate	Family: cardiovascular	7.3%	2.9%	0.1832	
		Abst. vs. moderate	Smoking: never	60.2%	52.3%	0.1575	
		Abst. vs. moderate	Smoking: current	21.1%	26.2%	0.1164	
		Abst. vs. moderate	Excitement-seeking	22.9	23.7	0.1782	

		Abst. vs. heavy	Sleep: short	7.3%	2.4%	0.1904
		Abst. vs. heavy	Sleep: long	12.7%	5.2%	0.2331
		Binge vs. non-binge	Age	29.6	30.2	0.1032
		Binge vs. non-binge	Smoking: never	52.2%	57.2%	0.1017
		Binge vs. non-binge	Excitement-seeking	23.7	23.2	0.1302
	Female – middle	Abst. vs. moderate	Income	2500.8	2578.1	0.1008
		Abst. vs. moderate	Income (equivalised)	1490.7	1552.2	0.1325
		Abst. vs. moderate	Smoking: never	50.7%	44.9%	0.1166
		Abst. vs. moderate	Sleep: long	9.9%	6.6%	0.1156
		Abst. vs. heavy	Ethnicity: white	98.5%	100.0%	0.1276
		Binge vs. non-binge	Impulsivity	22.9	22.5	0.1002
	Female – old	Abst. vs. moderate	Income (equivalised)	1651.2	1705.4	0.1055
		Abst. vs. moderate	Smoking: never	37.2%	31.8%	0.1163
		Abst. vs. heavy	Ethnicity: white	99.0%	100.0%	0.1076
		Abst. vs. heavy	Income	2372.1	2525.2	0.1990
		Abst. vs. heavy	Income (equivalised)	1651.2	1772.0	0.2350
		Abst. vs. heavy	Diabetes	3.8%	0.3%	0.2096
		Abst. vs. heavy	Smoking: past	46.4%	57.8%	0.2281
		Abst. vs. heavy	Smoking: current	16.4%	23.1%	0.1787
		Binge vs. non-binge	Excitement-seeking	20.0	19.5	0.1426
Non-adjustable	Female – young	Abst. vs. heavy	Income	2254.0	1777.0	0.5428
		Abst. vs. heavy	Income (equivalised)	1514.8	1372.2	0.2681
		Abst. vs. heavy	Family: dep/anx	28.5%	10.0%	0.4118
		Abst. vs. heavy	Sleep: normal	80.0%	92.4%	0.3169
	Female – middle	Abst. vs. heavy	Income (equivalised)	1490.7	1609.7	0.2568
		Abst. vs. heavy	Cancer	3.8%	12.1%	0.4224
	Female – old	Abst. vs. heavy	Smoking: never	37.2%	19.1%	0.3875

Abbreviations: abst. = abstinent

Supplemental Table S5.3. Fully adjusted models comparing abstinent participants to each drinking level, and binge-drinkers to non-binge-drinkers, in age- and sex stratified groups

		N	β	95% CI	p
Occasional (0-2.5 g/day) vs. abstinent	Full sample	16,838	0.018	-0.006 – 0.043	0.1432
	Male – young	1,710	-0.018	-0.109 – 0.074	0.7084
	Male – middle	2,181	0.044	-0.031 – 0.119	0.2494
	Male – old	1,456	-0.026	-0.109 – 0.056	0.5329
	Female – young	4,487	0.024	-0.022 – 0.069	0.3053
	Female – middle	4,142	0.045	0.004 – 0.086	0.0312
	Female – old	2,862	0.069	0.020 – 0.117	0.0056
Light (2.5-14.9 g/day) vs. abstinent	Full sample	33,827	0.056	0.033 – 0.078	<0.0001
	Male – young	5,605	-0.002	-0.085 – 0.080	0.9541
	Male – middle	5,916	0.093	0.026 – 0.160	0.0065
	Male – old	4,821	0.005	-0.067 – 0.077	0.8912
	Female – young	5,862	0.059	0.014 – 0.104	0.0104
	Female – middle	6,095	0.108	0.069 – 0.147	<0.0001
	Female – old	5,528	0.106	0.064 – 0.149	<0.0001
Moderate (15-29.9 g/day) vs. abstinent	Full sample	8,955	0.111	0.079 – 0.143	<0.0001
	Male – young	1,695	0.033	-0.061 – 0.127	0.4893
	Male – middle	1,966	0.174	0.098 – 0.250	<0.0001
	Male – old	2,030	0.052	-0.026 – 0.131	0.1917
	Female – young	584	0.131	0.024 – 0.238	0.0165
	Female – middle	1,094	0.145	0.067 – 0.224	0.0003
	Female – old	1,586	0.153	0.089 – 0.218	<0.0001
Heavy (30-60 g/day) vs. abstinent	Full sample	1,839	0.075	-0.009 – 0.158	0.0791
	Male – young	452	0.033	-0.104 – 0.169	0.6389
	Male – middle	529	0.111	-0.028 – 0.249	0.1168
	Male – old	531	0.054	-0.063 – 0.171	0.3655
	Female – young *	40	-0.401	-0.740 – -0.062	0.0206
	Female – middle *	122	0.021	-0.267 – 0.309	0.8869
	Female – old *	165	0.094	-0.065 – 0.252	0.2484
Binge-drinking yes vs no	Full sample	8,380	0.032	-0.002 – 0.066	0.0654
	Male – young	2,426	0.022	-0.029 – 0.073	0.3940

	Male – middle	1,534	0.021	-0.038 – 0.080	0.4816
	Male – old	851	-0.071	-0.162 – 0.020	0.1249
	Female – young	2,111	0.036	-0.028 – 0.101	0.2713
	Female – middle	909	0.134	0.046 – 0.222	0.0030
	Female – old	549	0.052	-0.061 – 0.165	0.3670

Supplement 6: Results stratified by drink type

Supplemental Table S6.1. Adjustable and non-adjustable differences between weighted groups

	Group	Contrast	Confounder	Group 1	Group 2	ASMD
Adjustable	Beer	Abst vs. occasional	Sex: male	41.8%	48.0%	0.1698
		Abst vs. occasional	Smoking: never	52.1%	46.2%	0.1202
		Abst vs. occasional	Smoking: current	21.6%	26.5%	0.1068
		Abst vs. light	Smoking: never	52.1%	44.6%	0.1523
		Abst vs. light	Smoking: current	21.6%	27.1%	0.1196
		Abst vs. light	Excitement-seeking	21.7	22.6	0.1884
		Abst vs. moderate	Diet quality	22.8	22.0	0.1698
		Abst vs. moderate	Excitement-seeking	21.7	22.8	0.2304
		Abst vs. heavy	Ethnicity: white	98.2%	100.0%	0.1755
		Abst vs. heavy	Rheumatoid/liver disease	2.2%	0.7%	0.1258
		Abst vs. heavy	Diet quality	22.8	21.8	0.2017
		Binge vs. non-binge	Age	40.6	41.8	0.1159
		Binge vs. non-binge	Sex: male	55.1%	44.4%	0.2149
		Binge vs. non-binge	Diet quality	22.2%	22.8%	0.1111
		Binge vs. non-binge	Smoking: never	43.1%	51.1%	0.1585
		Binge vs. non-binge	Excitement-seeking	22.7	21.9	0.1758
	Wine	Abst vs. moderate	Smoking: never	49.4%	43.8%	0.1122
		Abst vs. heavy	Age	43.6	45.9	0.2185
		Abst vs. heavy	Sex: male	30.1%	38.6%	0.1819
		Abst vs. heavy	Smoking: past	32.1%	39.8%	0.1607
		Abst vs. heavy	Smoking: current	18.5%	25.7%	0.1795
Non-adjustable	Beer	Abst vs. light	Sex: male	41.8%	52.7%	0.2991
		Abst vs. moderate	Sex: male	41.8%	56.2%	0.3968
		Abst vs. moderate	Smoking: never	52.1%	37.0%	0.3060
		Abst vs. moderate	Smoking: current	21.6%	33.2%	0.2536
		Abst vs. heavy	Sex: male	41.8%	59.8%	0.4936
		Abst vs. heavy	Smoking: never	52.1%	33.6%	0.3749

		Abst vs. heavy	Smoking: current	21.6%	40.8%	0.4197
		Abst vs. heavy	Excitement-seeking	21.7	23.0	0.2792
	Wine	Abst vs. heavy	Smoking: never	49.4%	34.5%	0.3002

Abbreviations: abst. = abstinent

Supplemental Table S6.2. Fully adjusted models comparing abstinent participants to each drinking level, and binge-drinkers to non-binge-drinkers, in age- and sex stratified groups

		N	β	95% CI	p
Occasional (0-2.5 g/day) vs. abstinent	Full sample	16,838	0.018	-0.006 – 0.043	0.1432
	Beer	2,369	0.036	-0.015 – 0.087	0.1632
	Wine	11,104	0.022	-0.002 – 0.047	0.0771
Light (2.5-14.9 g/day) vs. abstinent	Full sample	33,827	0.056	0.033 – 0.078	<0.0001
	Beer *	5,637	0.037	-0.010 – 0.085	0.1248
	Wine	22,399	0.065	0.043 – 0.087	<0.0001
Moderate (15-29.9 g/day) vs. abstinent	Full sample	8,955	0.111	0.079 – 0.143	<0.0001
	Beer *	2,194	0.134	0.061 – 0.207	0.0003
	Wine	5,010	0.119	0.080 – 0.157	<0.0001
Heavy (30-60 g/day) vs. abstinent	Full sample	1,839	0.075	-0.009 – 0.158	0.0791
	Beer *	793	0.051	-0.070 – 0.173	0.4062
	Wine *	626	0.037	-0.074 – 0.147	0.5151
Binge-drinking yes vs no	Full sample	8,380	0.032	-0.002 – 0.066	0.0654
	Beer	3,563	0.027	-0.026 – 0.080	0.3134
	Wine	2,958	0.055	0.001 – 0.110	0.0470

* Residual non-adjustable imbalance (ASMD>0.25) remained after propensity score weighing, rendering this between-group comparison prone to residual confounding.