

Appendix

Rapid disappearance of influenza following the implementation of COVID-19 mitigation measures in Hamilton, Ontario

Kevin Zhang,^{1,*} Avika Misra,² Patrick J. Kim,³ Seyed M. Moghadas,⁴ Joanne M. Langley,⁵ Marek Smieja^{3,6}

¹ Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada, M5S 1A8

² Faculty of Medicine, University of Ottawa, Ottawa, Ontario, Canada, K1H 8M5

³ Department of Medicine, McMaster University, Hamilton, Ontario, Canada, L8S 4L8

⁴ Agent-Based Modelling Laboratory, York University, Toronto, Ontario, Canada, M3J 1P3

⁵ Canadian Center for Vaccinology, Dalhousie University, IWK Health Centre and Nova Scotia Health Authority, Halifax, Nova Scotia, Canada, B3K 6R8

⁶ St. Joseph's Healthcare Hamilton, Hamilton, Ontario, Canada, L8N 4A6

*Corresponding author: kevink.zhang@mail.utoronto.ca

Table A1. Percent positivity of laboratory-confirmed influenza in Hamilton, Ontario in weeks 1-26 of 2010-2019.

Week	Mean	Median	Quartile 1	Quartile 3	Standard Deviation
1	18.5	21.1	11.0	26.2	10.3
2	16.0	15.6	3.6	29.8	11.2
3	17.4	21.0	5.4	26.1	10.0
4	16.5	12.2	5.8	26.6	11.9
5	16.8	17.3	8.8	24.9	10.3
6	17.5	16.7	12.7	23.8	9.2
7	15.2	16.0	7.4	21.5	8.8
8	15.9	15.2	6.8	23.8	9.3
9	14.6	13.9	8.2	20.7	9.1
10	14.8	14.2	5.8	20.0	10.1
11	12.7	11.2	3.9	21.6	8.5

12	10.2	10.9	5.1	16.0	6.2
13	11.4	10.9	5.6	18.4	6.8
14	11.5	9.8	3.6	20.3	8.4
15	8.5	7.6	4.1	11.7	5.8
16	7.4	5.4	2.5	13.1	5.9
17	7.3	5.8	2.2	11.9	5.6
18	5.5	6.3	2.2	7.1	3.4
19	3.2	2.0	0.9	5.8	3.2
20	3.3	1.6	0.7	6.3	3.3
21	1.9	0.7	0.0	2.4	3.0
22	1.4	0.7	0.0	3.0	1.5
23	1.2	0.0	0.0	2.1	1.9
24	0.8	0.0	0.0	1.9	1.1
25	0.3	0.0	0.0	0.3	0.6
26	0.0	0.0	0.0	0.0	0.0

Table A2. Percent positivity of laboratory-confirmed influenza, respiratory syncytial virus, metapneumovirus, parainfluenza, adenovirus, and rhinovirus/enterovirus in Hamilton, Ontario in weeks 1-26 of 2020.

Week	Influenza Percent Positivity	Respiratory Syncytial Virus Percent Positivity	Metapneumovirus Percent Positivity	Parainfluenza Percent Positivity	Adenovirus Percent Positivity	Rhinovirus/Enterovirus Percent Positivity
1	17.7	5.9	0.5	0.0	0.5	2.2
2	9.6	5.6	0.8	0.4	0.8	1.6
3	9.6	1.8	0.9	0.4	0.4	3.9
4	11.5	5.8	0.8	0.0	0.4	1.2
5	6.2	4.5	0.7	0.0	0.3	2.4
6	10.4	2.3	1.2	0.4	0.8	1.5
7	5.8	1.9	1.0	0.0	1.0	1.4
8	8.8	2.4	2.4	0.0	0.0	2.4
9	3.2	0.5	0.5	0.0	0.5	0.0
10	5.8	1.5	1.9	1.0	1.0	4.4

11	5.1	0.6	2.4	0.4	0.4	6.1
12	0.0	0.0	0.5	0.2	0.0	0.2
13	0.1	0.0	0.1	0.1	0.0	0.2
14	0.0	0.0	0.0	0.0	0.0	0.0
15	0.0	0.0	0.0	0.0	0.0	0.0
16	0.0	0.0	0.0	0.0	0.0	0.0
17	0.0	0.0	0.0	0.0	0.0	0.0
18	0.0	0.0	0.0	0.0	0.0	0.0
19	0.0	0.0	0.0	0.0	0.0	0.0
20	0.0	0.0	0.0	0.0	0.0	0.0
21	0.0	0.0	0.0	0.0	0.0	0.0
22	0.0	0.0	0.0	0.0	0.0	0.0
23	0.0	0.0	0.0	0.0	0.0	0.0
24	0.0	0.0	0.0	0.0	0.0	0.0
25	0.0	0.0	0.0	0.0	0.0	0.0
26	0.0	0.0	0.0	0.0	0.0	0.0

Table A3. Estimated model parameters from Bayesian inference.

Parameter	Mean	95% CrI
β_1	13.494	-0.891, 21.695
β_2	0.114	-0.888, 1.985
σ	4.005	1.932, 5.102