

20 **Abstract**

21 As COVID-19 continues to pose significant public health threats, quantifying the 22 effectiveness of different public health interventions is crucial to inform intervention 23 strategies. Using detailed epidemiological and mobility data available for New York City 24 and comprehensive modeling accounting for under-detection, we reconstruct the COVID-25 19 transmission dynamics therein during the 2020 spring pandemic wave and estimate the 26 effectiveness of two major non-pharmaceutical interventions—lockdown-like measures 27 that reducing contact rates and universal masking. Lockdown-like measures were 28 associated with >50% transmission reduction for all age groups. Universal masking was 29 associated with a \sim 7% transmission reduction overall and up to 20% reduction for 65+ 30 vear-olds during the first month of implementation. This result suggests that face covering 31 can substantially reduce transmission when lockdown-like measures are lifted but by itself 32 may be insufficient to control SARS-CoV-2 transmission. Overall, findings support the need 33 to implement multiple interventions simultaneously to effectively mitigate COVID-19 34 spread before the majority of population can be protected through mass-vaccination. 35 36 **Keywords:** COVID-19; transmission dynamics; effectiveness of intervention; social

37 distancing; lockdown; face covering

38

39 **Introduction**

40 Since the emergence of SARS-CoV-2 in late 2019, the virus has infected over 79 million 41 people and killed over 1.75 million worldwide by the end of 2020 (as of $12/27/20$).¹ Non-42 pharmaceutical interventions such as social distancing and face covering have been the

43 main strategies to contain COVID-19 during this pandemic in 2020. In late 2020, Phase-III 44 trials for several SARS-CoV-2 vaccines showed highly promising results and were granted 45 emergency use in several countries.^{2,3} However, before these vaccines become widely 46 available to the general population (likely in mid- or late 2021), non-pharmaceutical 47 interventions will need to remain the main strategies to contain COVID-19. In addition, 48 future (re)emerging infectious disease outbreaks may need to rely on similar non-49 pharmaceutical measures. It is thus critical to understand the effectiveness of different 50 non-pharmaceutical interventions implemented during the COVID-19 pandemic waves in 51 order to inform effective future planning while balancing economic need. For instance, 52 with face covering and social distancing by closing businesses as two main interventions, 53 the more effective face covering is, the more businesses could remain open. As a simplified 54 calculation, with a basic reproductive number (R_0) of 3 and minimal immunity, a city could 55 maintain 55% business capacity while curbing epidemic growth, if its residents could 56 reduce transmission by 40% using face covering [ie, effective reproductive number $R_t = 3 \times$ 57 55% \times (1 – 40%) = 0.99 < 1]; this threshold business capacity would drop to 33% if no 58 residents used face covering.

59

60 However, assessing the effectiveness and impact of a given intervention for COVID-19 has 61 been challenging due to low infection detection rates (many asymptomatic and mild 62 infections do not seek care or receive testing),⁴ fluctuation of those infection detection 63 rates, differential disease manifestation by age group,^{5,6} and concurrent public health 64 interventions. As such, while a few studies have assessed the overall effectiveness of

65 lockdown-like measures, to date, the effectiveness of specific measures including face 66 covering *under real-world conditions* remains unclear.

67

68 To estimate the effectiveness of different non-pharmaceutical public health interventions, 69 here we thus focus on the 2020 spring COVID-19 pandemic wave in New York City (NYC), 70 the first COVID-19 epidemic center in the United States, where detailed data are also 71 available. NYC experienced widespread COVID-19 transmission citywide since early March 72 and recorded over 200,000 cases and over 21,000 COVID-19 confirmed or probable deaths 73 during the following three months. To curb this intense transmission, NY State and NYC 74 implemented multiple intervention measures, including health promotion campaigns in 75 early March, telecommuting and staggered work schedule recommendations beginning the 76 week of March 8, public schools closure starting the week of March 15,7 stay-at-home 77 orders for non-essential workers starting the week of March $22₁⁸$ and requirements for use 78 of face covering in public starting the week of April 12.9 With these overlapping and far 79 reaching public health interventions, case diagnoses and hospitalizations peaked in April, 80 and started to decline substantially in late April and May. NYC was able to begin its phased 81 re-opening of industries starting the week of June 7, 2020.

82

83 In this study, we apply a model-inference system¹⁰⁻¹² developed to support the city's 84 COVID-19 pandemic response to reconstruct the underlying transmission dynamics of 85 COVID-19 in NYC during March 1 – June 6, 2020 (i.e. prior to the city's reopening). To 86 address the aforementioned challenges, our model-inference system simultaneously 87 assimilates three sources of data: 1) confirmed COVID-19 case data, 2) COVID-19

88 associated death data (both cases and deaths are assimilated by neighborhood and age 89 group), and 3) neighborhood-level mobility data to constrain the model system. This 90 enables inference of the overall infection rate (i.e. including those not documented by 91 surveillance), estimation of key transmission characteristics $(e.g., the reproductive$ 92 number) through time, and assessment of the effectiveness of different public health 93 interventions, including social distancing and face covering, implemented over time. We 94 further incorporate these estimates to project cases and deaths in the weeks beyond our 95 study period and compare the projections to independent observations in order to evaluate 96 the accuracy of these estimates. We conclude with a discussion on the implication of our 97 findings on strategies to safely reopen economies in places COVID-19 continues to pose 98 substantial public health threats.

99

100 **Results**

101 *Overall Epidemic Trends*

102 Following diagnosis of the first case in NYC, confirmed COVID-19 cases in the entire 103 population increased nearly exponentially during the first three weeks (Fig 1B) before 104 slowing down beginning the week of March 22, 2020 when NYC implemented a stay-at-105 home order. However, case trajectories differed substantially by age group. Foremost, 106 reported cases increased with age: the case trajectory for those aged 25-44 years mirrored 107 the overall epidemic curve, those older than 45 years had higher case rates, and those 108 under 25 years had the lowest case rates (Fig 1A and Table S1). Infants (i.e. <1 year), 109 however, had higher case rates than 1-4 and 5-14 year-olds (Fig 1). In addition, the timing 110 of peak case rate was mixed. Case rates in 25-44, 45-64, and 65-74 year-olds peaked

111 earliest during the week of March 29, 2020, followed by $\lt 1$, 1-4, 15-24, and 75+ year-olds 112 with a 1-week lag; in comparison, the case rate for 5-14 year-olds fluctuated with a less 113 clear peak during the weeks of March 29 – April 26, 2020.

114

115 The epidemic trends based on diagnosed cases, however, were obscured by varying 116 infection detection rates by age and through time. COVID-19 infections are more likely to 117 manifest as symptomatic illness and/or more severe disease in individuals with underlying 118 conditions and in older adults.^{5,6} Such differential clinical characteristics by age thus lead to 119 varying healthcare seeking behaviors and infection detection rates by age. In addition, 120 testing policies varied over the course of the Spring 2020 COVID-19 pandemic in NYC. 121 During this time, testing capacity was limited at the federal, state and local levels by 122 guidelines for who should be tested (due, for example, to limited availability of test kits, 123 swabbing supplies and reagents), which required prioritizing testing for severely ill 124 patients and those highly vulnerable to severe disease. Testing capacities expanded during 125 the week of March 8, 2020;¹³ however, by late March, material shortages (including testing 126 kits and personal protective equipment) again prompted the city to restrict testing to those 127 severely ill.¹⁴ Using our model-inference system, we estimated that infection detection 128 rates increased in early March, reaching a peak of around 20% for all ages overall during 129 the week of March 15, and declined afterwards before increasing again in mid-April.^{11,12} 130

131 After accounting for infection detection rates to include undiagnosed infections, a different 132 picture of the NYC spring outbreak emerges (Fig 2 and Fig S3). Estimated infection rates 133 were highest among 25-44 and 45-64 year-olds (Fig. 2 F and G), followed by 65-74 and $75+$

134 year-olds (Fig. 2 H and I), then 5-14 and 15-24 year-olds (Fig. 2 D and E), and were lowest 135 among <1 and 1-4 year olds (Fig. 2 B and C; Table S1). Estimated infection rates in the 136 younger age groups (in particular, 5-14, 15-24, and 25-44 year-olds; Fig. 2 D-F) peaked 137 during the week of March 22, 2020, followed by the three older age groups (i.e. 45-64, 65-138 $\,$ 74, and 75+ year-olds; Fig. 2 G-I) about a week later. 139 140 Given the large uncertainties in model estimates, we verified our estimates of infection 141 rates using available serology data collected during three phases of the pandemic (i.e. 142 early-phase in March,¹⁵ mid-phase in April,¹⁶ and end of the pandemic wave in June¹⁷). 143 Overall, our estimated cumulative infection rates were in line with corresponding 144 measures from antibody tests, for all three phases of the pandemic wave (for details, see 145 the Appendix of Yang et al. 2020^{11,12})

146

147 Overall effectiveness of interventions

148 The reproductive number at time- t (R_t) measures the average number of persons an 149 infected individual infects and thus reflects underlying epidemic dynamics. The epidemic 150 expands in size if R_t is above unity and subsides otherwise. In addition, when the entire 151 population is susceptible and no interventions are in place, R_t , referred to as the basic 152 reproductive number (R_0) , reflects the transmissibility of an infection in that population. 153 Here we estimated that R_t was 2.99 [median and interquartile range (IQR) : 2.32 – 3.86; 154 Table S2] during the first week of the pandemic (i.e. the week of March 1) in NYC, similar to 155 R_0 estimates reported for other places.^{18,19} It decreased to around 2.2 during the next two 156 weeks, when NY State declared a state of emergency and public awareness and voluntary

203

- 224 using mobility data alone would be higher (longer) than those estimated by the model-
- 225 inference system additionally based on case and mortality data *for weeks when face*

226 *covering in public was mandated* as it would lead to further reductions in transmission (i.e., 227 temporality and direction of the impact). Second, while the efficacy of masking (i.e., 228 measured under ideal conditions of mask quality and correct use) likely does not vary by 229 individual, the effectiveness of masking (i.e., measured under real-world, often imperfect 230 conditions) and impact of this intervention could vary by subpopulation due, for example, 231 to different usage rates of masks; as such, we expect the predictive errors to be larger for 232 age groups with higher compliance of masking (i.e., magnitude of the impact). Our analyses 233 largely confirmed both predictions. For the first, as shown in Fig 4, transmission rates 234 predicted using a linear regression model with the observed mobility as the sole predictor 235 were higher than those estimated by the model-inference system, following the face 236 covering mandate starting the week of April 12.9 For the latter, the discrepancies in the two 237 model estimates (i.e. the gaps between the dashed and solid blue lines; Fig 4) appeared to 238 increase with age and were largest among the two elderly age groups who have been 239 reported to more frequently use masks. $22-24$ However, infants ≤ 1 year) appeared to have a 240 larger reduction than other children groups; this could have been due to transmission 241 reduction related to their sources of infection (e.g. their caretakers and healthcare settings 242 where they tended to be exposed). Similar patterns held for the effective infectious period 243 (Fig 4).

244

245 Given these observations, we further used the discrepancies in the two model estimates to 246 approximate the impact of face covering on reducing COVID-19 transmission. Combining 247 the reduction in the transmission rate and effective infectious period, we estimated that, 248 for all ages combined, face covering contributed to a 6.6% (95% CI: 0.8 – 12.4%) reduction

260

261 Retrospective projections of cases and deaths

262 NYC started phased reopening from the week of 6/7/2020, which allows industries to 263 gradually reopen per a four-phase plan. ²⁶ For instance, manufacturing industries were 264 allowed to reopen starting the week of $6/7/2020$ (dubbed "Phase 1"), whereas real estate 265 was allowed to reopen starting the week of $6/21/2020$ (dubbed "Phase 2"), and personal 266 care services were allowed starting the week of 7/6/2020 (dubbed "Phase 3"). As such, 267 population mobility has increased gradually during this time, which could lead to increased 268 transmission of unknown magnitude. Such changes also offer an opportunity to test the 269 accuracy of our estimates – should the estimated effectiveness of reducing contact rates 270 and utilizing face coverings be accurate, these estimates could be used to anticipate 271 changes in transmission in response to the changing mobility and in turn the epidemic

272 dynamics after reopening. We thus used these estimates to generate projections of cases 273 and deaths for the 8 weeks beyond our study period, and compared the projections to 274 available, independent corresponding observations. Overall, our projections 275 underestimated the total number of cases (relative error of median projections: -27% over 276 8 weeks; Fig 5A) but were able to accurately estimate the total number of deaths (relative 277 error: -2% over 8 weeks; Fig 5B). In addition, examination of age-grouped projections 278 shows that the underestimation of cases was mostly among younger age groups whose 279 case rates had increased in June (1-4, 5-14, 15-24 and 25-44 year-olds; Fig S5). These 280 recent increases in young cases may have resulted from more young adults returning to 281 work including some in service industries with high contact rates and, relatedly, sending 282 their children to childcare and/or summer camps due to a lack of caretakers at home 283 [information from NYC Department of Health and Mental Hygiene (DOHMH) community 284 investigation; unpublished]. In addition, increased risk behaviors of some young 285 individuals (e.g., large parties without physical distancing²⁵) may have also contributed to 286 the increased cases among young adults in late June.²⁷ Consistently, COVID-19 associated 287 mortality, mostly occurring among older adults continued to decrease and were accurately 288 predicted for different age groups (Fig S6).

289

290 **Discussion**

291 The spring 2020 pandemic wave in NYC, the first epidemic center in the US, provides a test 292 case to study COVID-19 epidemiological characteristics and the effectiveness of public 293 health interventions. Through comprehensive modeling, we have reconstructed the 294 transmission dynamics and estimated the effectiveness of two major interventions, social

295 distancing and mandatory face covering in public. Our results show that reducing contact 296 rates (mainly via school closures and voluntary or mandated stay-at-home measures) likely 297 contributed to the largest reduction in transmission in the population overall $\left(\sim 70\% \right)$ and 298 for most age groups (>50% for all age groups). Widespread use of face covering likely 299 contributed to an additional \sim 7% overall reduction and up to \sim 20% reduction among 65+ 300 vear-olds during the first month face covering was mandated in public places. Our findings 301 largely consolidate previous model estimates on the impact of lockdown-like 302 measures^{4,28,29} and studies on face covering in reducing COVID-19 transmission. These 303 findings provide insights that can inform COVID-19 mitigation efforts in the coming months 304 before the majority of population can be protected through mass-vaccination, as well as 305 control strategies for other (re)emerging infections in the future. 306 307 Lockdown-like measures where confinement at home is encouraged or mandated through 308 school closures, telework policies, closure of non-essential businesses, and stay-at-home 309 orders have been a major control measure to curb COVID-19 spread. In effect, such 310 measures reduce population contact rates and thus transmission. Previous modeling 311 studies estimated that lockdowns reduced COVID-19 transmission (measured by R_t) by 312 58% in Wuhan, China,⁴ 45% (95% CI: 42-49%) in Italy,²⁹ and 77% (95% CI: 76-78%) in 313 France.²⁸ Our estimate for NYC overall $\left(\sim 70\% \right)$ is consistent with these previous estimates. 314 In addition, our estimates show that reducing population contact rates effectively reduced 315 transmission across all age groups (ranging from a 51% reduction among 1-4 year-olds to

316 83% among 5-14 year-olds; Table 1). Together, these findings underscore the importance

317 of reducing contact rates through, for example, physical distancing in places with

318 continuous community transmission of COVID-19.

319

320 The use of surgical masks or cloth face coverings has been another major preventive 321 measure for COVID-19. Studies overall have shown that surgical masks could substantially 322 reduce onward transmission albeit with a large range of efficacy estimates across 323 settings.³⁰ However, it remains unclear the overall effectiveness of universal face covering 324 requirements at the population level, especially during a pandemic, due to several factors: 325 1) The overall effectiveness depends on compliance which may vary across subpopulations 326 and time; 2) Improper use of face coverings (e.g. without covering the nose and/or mouth 327 or improper handling³¹ can reduce the effectiveness of face covering; 3) Face coverings 328 are required and mostly worn in public and thus likely have a lower impact in private 329 settings, particularly in reducing household transmission; consequently, the relative impact 330 of face covering depends on the relative contribution of different sources of transmission 331 (e.g. household vs. community) at a given time and *vice versa*; and 4) Use of face coverings 332 may lead to complacency and less stringent adherence to social distancing and stay-at-333 home behaviors. Here we estimated a \sim 7% reduction in overall transmission during the 334 first month of the face covering mandate. However, the estimated effectiveness varied 335 largely across age groups with much higher effectiveness among older adults $(-20\%$ for 336 both $65-74$ and $75+$ year-olds vs. <10% for other age groups). This discrepancy was likely 337 due to the differential compliance and types of face covering used. Observational studies in 338 Wisconsin and surveys nationwide in April/May reported about 2-fold higher rates of face 339 covering usage among older adults versus younger adults and minors.²²⁻²⁴ In addition, due

340 to the shortage of surgical masks during March–May, 32 older adults at higher risk of severe 341 COVID-19 infection were more likely to use surgical masks whereas younger age groups 342 more frequently used non-medical cloth coverings, which are often less effective^{33,34} (e.g. 343 measured ultrafine filtration efficiency is \sim 50% for surgical masks vs. \sim 10-25% for T-shirt 344 and \sim 25-35% for cotton covers³⁴).

345

346 When lockdown-like measures are lifted, residents will spend more time outside their 347 homes than during the lockdown. Adjusting for the time spent outside of homes (~ 8.3) 348 hours in April 2020 vs. \sim 11.5 hours in June-July 2020 and \sim 13.5 hours pre-pandemic; NYC 349 data³⁵), universal face covering would have reduced overall transmission by \sim 9–11% (i.e., 350 6.6% multiplied by a factor of 1.4–1.6) during reopening, given the same rates of face 351 covering as in April. However, if the same effectiveness among older adults were achieved 352 among other age groups, universal face covering could reduce overall transmission by up to $353 \sim 28-32\%$ (i.e., 20% multiplied by a factor of 1.4–1.6). The implication of this latter 354 estimate is two-fold. On the one hand, it suggests that for places with high level of 355 transmission, implementing face covering *alone* is likely insufficient to lower the effective 356 reproductive number R_t to <1 in order to control the epidemic [for instance, for a city with 357 an $R_0 = 3$, the resulting R_t would be $3 \times (1-30\%) = 2.1$. This finding is consistent with the 358 observed resurgence of COVID-19 cases in NYC during fall/winter 2020 despite the 359 concurrent high usage rate of face coverings \sim 90% of survey respondents in NYC reported 360 always or frequently wearing masks in public in July 2020;³⁶ and this number was likely 361 higher during fall/winter 2020). On the other hand, our findings also suggest that 362 improving effective usage rates of face coverings, especially among younger age groups,

363 could significantly mitigate the risk of resurgence of COVID-19 infections during re-opening 364 (i.e., \sim 30% reduction without compromising economic growth). It is thus crucial for future 365 research to understand reasons for the use/non-use and selection of face coverings by age 366 group to inform strategies to increase consistent and correct mask use in settings where 367 social distancing is not possible.

368

369 It is important to note, however, that not all individuals have the same opportunities to 370 physically distance and/or adopt face coverings during a pandemic, despite government 371 mandates. For instance, over one million frontline workers in NYC (e.g., healthcare 372 workers, transportation workers, janitors, and grocery clerks, which comprise 25 percent 373 of the city's workforce) had to continue their essential work during the pandemic.³⁷ In 374 addition, data from the U.S. Bureau of Labor Statistics suggest Black and Latino 375 communities have less opportunities to work from home.³⁸ Consequently, NYC 376 neighborhoods with more frontline workers and/or Black and Latino residents tended to 377 have lower reductions in population mobility during the pandemic.³⁵ In NYC, these 378 communities also experience a number of social conditions that are thought to exacerbate 379 COVID-19, including overcrowded multigenerational households, poverty, and high 380 prevalence of chronic diseases. These communities, known to also carry a higher burden of 381 underlying health conditions, suffered greater impacts from COVID-19 and have expressed 382 fear and experiences of racialized bias when wearing a face covering. $39,40$ Further research 383 is warranted to investigate such health disparities. In addition, future policies should take 384 into account structural inequities in labor trends, overcrowded housing, and underlying

385 conditions and adopt additional preventive measures to protect those vulnerable 386 communities.

387

388 We also note there remain large uncertainties in our estimates due to several limitations. 389 First, we used population mobility as a proxy for contact rates rather than more direct 390 measures. Similar approximation and uncertainty applied to our estimates of the 391 effectiveness of face covering. Future studies are thus warranted for further assessment. 392 For instance, large population scale surveys documenting changes of the intensity and 393 pattern of contact during the pandemic could provide more accurate measures of contact 394 rates among different age groups and over time. Second, while we restricted our analysis 395 on the effectiveness of face covering to a period when masks were mandated, there remain 396 other residual confounding effects. For instance, increased awareness of COVID-19 and 397 health risk among key age groups such as the elderly may have contributed to further 398 reductions of transmission through other precautions in addition to face covering; this may 399 have led to an overestimation of the effectiveness of face covering for those age groups. 400 Third, here we focused on estimating the effectiveness of interventions in the general 401 population without segregating key settings with intense transmission (e.g., long-term care 402 facilities). Future studies should assess the impact of interventions targeting such high-risk 403 settings. Lastly, our estimates here were largely based on the first wave of the pandemic 404 and may not fully capture subsequent changes in awareness and perception of COVID-19 405 and related behavioral adjustment during later waves. However, we have also used a 406 similar methodology to estimate the effectiveness of reducing contact rates and face 407 covering under different city reopening schedules and generated long-term projections for

408 NYC; the projections generated during June 2020 for the period of June 2020 – May 2021 409 have been consistent with observations up to the end of 2020 (i.e., at the time of this 410 writing; see the projected resurgence and second wave in Yang et al.⁴¹ and comparison 411 with available data in Fig S7). These results thus support the robustness of our estimates 412 here.

413

414 Our study also has several strengths. In particular, our estimates were based on 415 comprehensive model-inference incorporating multiple data streams and further evaluated 416 using model projections. Our results thus provide an assessment of two major public heath 417 interventions (reducing contact rates and face covering) at the population level where the 418 overall effectiveness depends on multiple factors in addition to the efficacy of a given 419 intervention. Altogether, our estimates support the need for multiple interventions 420 (including reducing contact rates by, e.g., restricting occupancy, universal face covering, 421 and, albeit not studied here, testing, contact tracing, isolation and timely treatment of 422 cases) in order to effectively mitigate the spread of COVID-19 as it continues to pose 423 threats to public health.

424

425 **Methods**

426 *Data*

427 COVID-19 cases included all laboratory-confirmed cases by week of diagnosis reported to 428 the NYC DOHMH. Mortality data by week of death combined confirmed and probable 429 COVID-19-associated deaths. Confirmed COVID-19-associated deaths were defined as those 430 occurring in persons with laboratory-confirmed SARS-CoV-2 infection, and probable

454 The epidemic model used in this study was described in Yang et al. 2020^{11,12} Briefly, the

455 model simulated intra- and inter neighborhood transmission of COVID-19 using a

456 susceptible-exposed-infectious-removed (SEIR) network model:

457
\n
$$
\begin{cases}\n\frac{dS_i}{dt} = -S_i \sum_{j=1}^{j=42} b_j \beta_{city} c_{ij} I_j / N_j \\
\frac{dE_i}{dt} = S_i \sum_{j=1}^{j=42} b_j \beta_{city} c_{ij} I_j / N_j - \frac{E_i}{Z} \\
\frac{dI_i}{dt} = \frac{E_i}{Z} - \frac{I_i}{D} \\
\frac{dR_i}{dt} = \frac{I_i}{D}\n\end{cases}
$$
\n(Fqn 1)

458

459 where S_i , E_i , I_i , R_i , and N_i are the numbers of susceptible, exposed (but not yet infectious), 460 infectious, and removed (either recovered or deceased) individuals and the total 461 population, respectively, from a given age group in neighborhood *i*. Note that due to model 462 complexity and a lack of information for parameterizing interactions among age groups, we 463 modeled each age group separately (i.e., combining all sources of transmission to each age 464 group; see further detail on parameter estimation below); as such, Eqn 1 describes the 465 spatial transmission across neighborhoods without interactions among age groups. β_{city} is 466 the citywide transmission rate, which incorporated seasonal variation as observed for 467 OC43, a beta-coronavirus in humans from the same genus as SARS-CoV-2.¹² To allow 468 differential transmission in each neighborhood, we included a multiplicative factor, b_i , to 469 scale neighborhood local transmission rates. Z and D are the latency and infectious 470 periods, respectively (Table S4).

471

472 The contact rates (c_{ii}) in each neighborhood over time and connectivity among

473 neighborhoods were computed based on mobility data. The model also accounted for

 482 and state variables (e.g., number of susceptible and infectious individuals in each 483 neighborhood) for each week, we ran the network-model stochastically with a daily time 484 step in conjunction with the ensemble adjustment Kalman filter (EAKF)⁴⁵ and fit to weekly 485 case and mortality data from the week starting March 1 to the week ending June 6, 2020. 486 The posterior distribution of each model parameter/variable was updated for that week at 487 the same time.⁴⁵ This parameter estimation process was done separately for each of the 488 eight age groups (i.e. <1, 1-4, 5-14, 15-24, 25-44, 45-64, 65-74, and 75+ years). To include 489 transmission from other age groups, we used measured intra and inter-group contacts 490 from the POLYMOD study²¹ to compute the total number of contacts made with each age 491 group and adjusted the *prior* range of the transmission rate (β_{city}) for each age group 492 accordingly. The posterior estimate was computed based on cases and mortality data for 493 each group, which included all sources of infection. Thus, the estimated transmission rate 494 for each age group nevertheless included all sources of transmission. To account for 495 stochasticity in model initiation, we ran the parameter estimation process independently 496 10 times. Results for each age group were combined from these 10 runs (each with 500

503

504 Estimating the effectiveness of reducing contact rates

505 The R_t estimates from the model-inference system capture changes in transmission due to 506 various interventions, i.e., the overall effectiveness of all implemented interventions. To 507 separately estimate the effectiveness of interventions that reduce contact rates, we used 508 human mobility as a measure of population contact rate to estimate the changes in R_t in 509 response to changing population contact rates. Specifically, we regressed the R_t estimates 510 from the full model-inference system on the mobility data: $R_t = a_0 + a_1 M_{\text{ave},t}$ (Eqn 2), 511 where M_{avet} is the mean of all intra-neighborhood mobility at week-*t*. We then computed 512 the effectiveness of reducing contact rate based on $\widehat{R_t}$, the R_t estimate from this regression 513 model solely based on the observed mobility. That is, the reduction in R_t by week-*t*, likely 514 $\,$ due to reducing contact rate, was computed as $(\widehat R_t-\widehat R_0)/\widehat R_0$ (Eqn 3). To test the robustness 515 of our method, we performed the same analysis for individual UHF neighborhoods $(n = 42)$. 516

517 **Estimating the effectiveness of face covering/masking**

518 In addition to changes in population contact rates, face covering/masking was another

 519 major control measure implemented beginning the week of April 12, 2020 when NYC

520 mandated residents wear face coverings in public places. To estimate the effectiveness of 521 face covering, we first estimated the changes in transmission rate and effective infectious 522 period, two model parameters determining R_t , due to changes in mobility (as opposed to 523 masking) using regression models similar to Eqn 2. Specifically, we regressed the estimated 524 citywide transmission rate (or effective infectious period) from the full model-inference s 525 system on average mobility: $Y_t = a_0 + a_1 M_{avet}$ (Eqn 4), where Y_t is β_{city} or *D*. We then 526 computed the relative reduction in transmission rate (or effective infectious period) due to 527 face covering as $\eta_Y = E((Y_t - \widehat{Y}_t)/Y_t)$, where $E(\cdot)$ gives the mean over the relevant 528 timeframe (here we estimated two timeframes, i.e., 1 month following the mandate and 529 over 8 weeks up to $6/6/2020$. Combining both reductions, we computed the effectiveness 530 of face covering as $\eta = 1 - (1 - \eta_B)(1 - \eta_D)$. Of note, while mechanistically face coverings 531 act primarily by reducing the probability of transmission (i.e., transmission rate), here we 532 included both the potential impact on the transmission rate (η_{β}) and effective infectious 533 period (η_D) , mainly because the multiplicative relationship of the two variables with R_t 534 makes it challenging to separate the two effects. Nevertheless, reductions in the infectious 535 period via face covering are possible. A recent study on animals showed that masking could 536 reduce the severity of infection;⁴⁷ if persons with milder infection experience shorter 537 duration of viral shedding (there is some evidence for this, e.g., from He et al.⁴⁸), milder 538 symptoms in individuals infected while wearing face covering could lead to shorter 539 infectious period of these individuals.

540

541 Projections of cases and deaths

542 To evaluate the accuracy of model estimates, in particular, the effectiveness of transmission 543 reduction by reducing contact rates and use of face covering, we tested if these estimates 544 along with the model could generate accurate predictions of cases and deaths for 8 weeks 545 beyond the study period (i.e. from the week of 6/7/2020 to the week of 7/26/2020). We 546 first projected the citywide transmission rate and infectious period based on observed 547 mobility using Eqn 4; these estimates thus accounted for changes due to changes in contact 548 rates. To incorporate the reduction in transmission by face covering, we further reduced 549 the projected city transmission rate by a factor of $1 - \eta_{\beta} p_{out}$ and the infectious period by a 550 factor of $1 - \eta_D p_{out}$, where p_{out} is a factor to adjust for time spent outside of the home 551 during each week. To reflect longer-term usage rates of face covering, we used η_B and η_D 552 estimated during the entire 8 weeks face covering was required (i.e. from the week of 553 $4/12/2020$ to the week of $5/31/2020$. Finally, we used estimates of population 554 susceptibility and infection rates at the end of the week of $5/31/2020$ to model initial 555 conditions and integrated the SEIR network model forward stochastically for 8 weeks using 556 the projected transmission rate and infectious period.

557

558 **Acknowledgments:**

559 This study was supported by the National Institute of Allergy and Infectious Diseases

560 (AI145883), the National Science Foundation Rapid Response Research Program (RAPID;

561 2027369), and the NYC DOHMH. We thank the NYC DOHMH Bureau of Vital Statistics team

562 and EpiData team; in particular, Mary Huynh, Sharon K. Greene, Alice Yeung, Anne Fine,

563 Miranda S. Moore and Kevin Guerra, for data management and provision. We thank Jennifer

564 Brite at NYC DOHMH for coordinating discussions on modeling COVID-19 in NYC and Alana

590 3. Pharmacy Times. Pfizer, BioNTech Granted Authorization for COVID-19 Vaccine in 591 the European Union. 2020. https://www.pharmacytimes.com/news/pfizer-biontech-592 granted-authorization-for-covid-19-vaccine-in-the-european-union (accessed 12/23/20. 593 4. Li R, Pei S, Chen B, et al. Substantial undocumented infection facilitates the rapid 594 dissemination of novel coronavirus (SARS-CoV-2). *Science* 2020; 368(6490): 489-93. 595 5. The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. The 596 epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-597 19)—China, 2020. *China CDC Weekly* 2020; 2(8): 113-22. 598 6. CDC COVID-19 Response Team. Preliminary estimates of the prevalence of selected 599 underlying health conditions among patients with coronavirus disease 2019—United 600 States, February 12–March 28, 2020. *Morbidity and Mortality Weekly Report* 2020; **69**(13): 601 382. 602 7. New York City Office of the Mayor. New York City to Close All School Buildings and 603 Transition to Remote Learning. $3/15/2020$ 2020. https://www1.nyc.gov/office-of-the-604 mayor/news/151-20/new-york-city-close-all-school-buildings-transition-remote-learning. 605 8. New York State Department of Health. New York State on PAUSE. 2020. 606 https://coronavirus.health.ny.gov/new-york-state-pause (accessed 5/16/2020 2020). 607 9. New York city Office of the Mayor. Transcript: Mayor de Blasio Holds Media 608 Availability on COVID-19. $4/12/2020$ 2020. https://www1.nyc.gov/office-of-the-609 mayor/news/249-20/transcript-mayor-de-blasio-holds-media-availability-covid-19. 610 10. Yang W, Kandula S, Shaman J. Projections of COVID19 Epidemic Outcomes and 611 Healthcare Demands for New York City (NYC). 2020. https://github.com/wan-612 yang/COLUMBIA-COVID19-PROJECTIONS-FOR-NYC (accessed 5/12/2020 2020). 613 11. Yang W, Kandula S, Huynh M, et al. Estimating the infection fatality risk of COVID-19 614 in New York City, March 1-May 16, 2020. *medRxiv* 2020: 2020.06.27.20141689. 615 12. Yang W, Kandula S, Huynh M, et al. Estimating the infection-fatality risk of SARS- 616 CoV-2 in New York City during the spring 2020 pandemic wave: a model-based analysis. 617 *The Lancet Infectious Diseases*. 618 13. Governor.ny.gov. After Weeks of Demanding Approval, Governor Cuomo Announces 619 FDA Gives New York State Authority to Conduct All COVID-19 Testing at Public and Private 620 Labs. March 13, 2020 2020. https://www.governor.ny.gov/news/after-weeks-demanding-

- 621 approval-governor-cuomo-announces-fda-gives-new-york-state-authority (accessed
- 622 6/22/2020 2020).
- 623 14. Johnson CY, Sun LH, McGinley L. In hard-hit areas, testing restricted to health care
- 624 workers, hospital patients. March 21, 2020 2020.
- 625 https://www.washingtonpost.com/health/2020/03/21/coronavirus-testing-
- 626 strategyshift/ (accessed $6/22/2020$ 2020).
- 627 15. Havers FP, Reed C, Lim T, et al. Seroprevalence of Antibodies to SARS-CoV-2 in 10
- 628 Sites in the United States, March 23-May 12, 2020. *Jama Intern Med* 2020.
- 629 16. www.governor.ny.gov. Amid Ongoing COVID-19 Pandemic, Governor Cuomo
- 630 Announces Results of Completed Antibody Testing Study of 15,000 People Showing 12.3
- 631 Percent of Population Has COVID-19 Antibodies. MAY 2, 2020 2020.
- 632 https://www.governor.ny.gov/news/amid-ongoing-covid-19-pandemic-governor-cuomo-
- 633 announces-results-completed-antibody-testing (accessed 5/12/2020 2020).
- 634 17. Government NYS. Video, Audio, Photos & Rush Transcript: Governor Cuomo
- 635 Announces U.S. Open to Be Held Without Fans from August 31st to September 13th.
- 636 6/16/2020 2020. https://www.governor.ny.gov/news/video-audio-photos-rush-
- 637 transcript-governor-cuomo-announces-us-open-be-held-without-fans-august.
- 638 18. Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of Novel
- 639 Coronavirus-Infected Pneumonia. *N Engl J Med* 2020; **382**(13): 1199-207.
- 640 19. Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and
- 641 international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling
- 642 study. *Lancet* 2020.
- 643 20. Goldbaum C. Coronavirus in N.Y.: Riders Ditch the Subway as Fear Spreads.
- 644 3/13/2020 2020. https://www.nytimes.com/2020/03/13/nyregion/coronavirus-subway-
- 645 public-transit.html (accessed $6/22$ 2020).
- 646 21. Mossong J, Hens N, Jit M, et al. Social contacts and mixing patterns relevant to the
- 647 spread of infectious diseases. *PLoS Med* 2008; **5**(3): e74.
- 648 22. Arp NL, Nguyen TH, Graham Linck EJ, et al. Use of face coverings by the public
- 649 during the COVID-19 pandemic: an observational study. *medRxiv* 2020:
- 650 2020.06.09.20126946.

- 651 23. Haischer MH, Beilfuss R, Hart MR, et al. Who is wearing a mask? Gender-, age-, and
- 652 location-related differences during the COVID-19 pandemic. *medRxiv* 2020:
- 653 2020.07.13.20152736.
- 654 24. Ritter Z, Brenan M. New April Guidelines Boost Perceived Efficacy of Face Masks.
- 655 2020. https://news.gallup.com/poll/310400/new-april-guidelines-boost-perceived-
- 656 efficacy-face-masks.aspx (accessed $5/13/2020$.
- 657 25. Barone V, Marino J. Hundreds pack St. Marks Place to drink, party despite
- 658 coronavirus. 2020. https://nypost.com/2020/06/12/hundreds-pack-st-marks-place-to-
- 659 drink-despite-coronavirus/ (accessed 7/14 2020).
- 660 26. New York State Government. Reopening New York City. 2020.
- 661 https://forward.ny.gov/reopening-new-york-city (accessed 7/13 2020).
- 662 27. Durkin E. New York City sees uptick in coronavirus cases among young adults. 2020.
- 663 https://www.politico.com/states/new-york/albany/story/2020/07/13/new-york-city-
- 664 sees-uptick-in-coronavirus-cases-among-young-adults-1300223 (accessed 7/14 2020).
- 665 28. Salje H, Tran Kiem C, Lefrancq N, et al. Estimating the burden of SARS-CoV-2 in 666 France. *Science* 2020; **369**(6500): 208-11.
- 667 29. Gatto M, Bertuzzo E, Mari L, et al. Spread and dynamics of the COVID-19 epidemic in
- 668 Italy: Effects of emergency containment measures. *Proc Natl Acad Sci U S A* 2020; 117(19): 669 10484-91.
- 670 30. Liang M, Gao L, Cheng C, et al. Efficacy of face mask in preventing respiratory virus
- 671 transmission: A systematic review and meta-analysis. *Travel Med Infect Dis* 2020: 101751.
- 672 31. Parker-Pope T. How NOT to Wear a Mask. $4/8/2020$ 2020.
- 673 https://www.nytimes.com/2020/04/08/well/live/coronavirus-face-mask-mistakes.html.
- 674 32. Bernstein L, Safarpour A. Mask shortage for most health-care workers extended into
- 675 May, Post-Ipsos poll shows. $5/20/2020$ 2020.
- 676 https://www.washingtonpost.com/health/mask-shortage-for-most-health-care-workers-
- 677 extended-into-may-post-ipsos-poll-shows/2020/05/20/1ddbe588-9a21-11ea-ac72-
- 678 3841fcc9b35f_story.html.
- 679 33. Fischer EP, Fischer MC, Grass D, Henrion I, Warren WS, Westman E. Low-cost
- 680 measurement of face mask efficacy for filtering expelled droplets during speech. *Science*
- 681 *Advances* 2020; **6**(36): eabd3083.

- 682 34. O'Kelly E, Pirog S, Ward J, Clarkson PJ. Ability of Fabric Facemasks Materials to Filter
- 683 Ultrafine Particles at Coughing Velocity for Home Made and Fabric Face Mask Creation.
- 684 *medRxiv* 2020: 2020.04.14.20065375.
- 685 35. SafeGraph. Weekly Patterns: Foot Traffic Data To Understand The COVID-19
- 686 Pandemic. 2020. https://www.safegraph.com/weekly-foot-traffic-patterns.
- 687 36. New York Times. Mask-Wearing Survey Data. 2020.
- 688 https://github.com/nytimes/covid-19-data/tree/master/mask-use (accessed
- 689 12/22/2020.
- 690 37. Stringer SM. New York City's Frontline Workers. March 2020 2020.
- 691 https://comptroller.nyc.gov/reports/new-york-citys-frontline-workers/.
- 692 38. Gould E, Shierholz H. Not everybody can work from home: Black and Hispanic
- 693 workers are much less likely to be able to telework. $3/19/2020$ 2020.
- 694 https://www.epi.org/blog/black-and-hispanic-workers-are-much-less-likely-to-be-able-
- 695 to-work-from-
- 696 home/#:~:text=Less%20than%20one%20in%20five,race%20and%20ethnicity%2C%202
- 697 017–
- 698 2018&text=The%20data%20below%20can%20be%20saved%20or%20copied%20directl
- 699 y%20into%20Excel.&text=The%20next%20figure%20illustrates%20the,who%20can%20
- 700 telework%20by%20wage. (accessed 8/17 2020).
- 701 39. Taylor DB. For Black Men, Fear That Masks Will Invite Racial Profiling. 2020.
- 702 https://www.nytimes.com/2020/04/14/us/coronavirus-masks-racism-african-
- 703 americans.html.
- 704 40. Ruiz NG, Horowitz JM, Tamir C. Many Black and Asian Americans Say They Have
- 705 Experienced Discrimination Amid the COVID-19 Outbreak. 7/1/2020 2020.
- 706 https://www.pewsocialtrends.org/2020/07/01/many-black-and-asian-americans-say-
- 707 they-have-experienced-discrimination-amid-the-covid-19-outbreak/.
- 708 41. Yang W, Kandula S, Shaman J. Simulating Epidemic Outcomes under Different
- 709 Occupancy and Rollback Policies. Posted online on 6/30/2020. https://github.com/wan-
- 710 yang/re-opening analysis/blob/master/report3 occupancy.pdf.
- 711 42. New York City Department of Health and Mental Hygiene (DOHMH) COVID-19
- 712 Response Team. Preliminary Estimate of Excess Mortality During the COVID-19 Outbreak

- 713 New York City, March 11–May 2, 2020. *MMWR Morbidity and mortality weekly report*
- 714 2020: 69: 603-5.
- 715 43. New York City Department of Health and Mental Hygiene. NYC UHF 42
- 716 Neighborhoods. http://a816-dohbesp.nyc.gov/IndicatorPublic/EPHTPDF/uhf42.pdf.
- 717 44. Lasry A, Kidder D, Hast M, et al. Timing of Community Mitigation and Changes in
- 718 Reported COVID-19 and Community Mobility Four U.S. Metropolitan Areas, February 26-
- 719 April 1, 2020. *MMWR Morbidity and mortality weekly report* 2020; 69(15): 451-7.
- 720 45. Anderson JL. An Ensemble Adjustment Kalman Filter for Data Assimilation. Mon
- 721 *Weather Rev* 2001; **129**(12): 2884-903.
- 722 $46.$ Heffernan JM, Smith RJ, Wahl LM. Perspectives on the basic reproductive ratio. *J R*
- 723 *Soc Interface* 2005; **2**(4): 281-93.
- 724 $\,$ 47. Chan JF, Yuan S, Zhang AJ, et al. Surgical mask partition reduces the risk of non-
- 725 contact transmission in a golden Syrian hamster model for Coronavirus Disease 2019
- 726 (COVID-19). *Clinical infectious diseases : an official publication of the Infectious Diseases*
- 727 Society of America 2020.
- 728 48. He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility 729 of COVID-19. *Nat Med* 2020; **26**(5): 672-5.
- 730

731 **Table**

735 and voluntary or mandated stay-at-home measures.

736

738 **Figures**

- 739 **Fig 1.** Epidemic dynamics. Reported laboratory confirmed cases (A) and cumulative cases
- 740 (B) per 100,000 population by week of diagnosis for all ages overall and by age group.

all <1 1−4 5−14 15−24 25−44 45−64 65−74 75+

742 **Fig 2.** Estimated changes in the effective reproductive number and infection rates. Blue 743 lines show the estimated effective reproductive number (R_t) for each week; surrounding 744 areas show the 50% and 95% CrIs. Superimposed boxes (right y-axis) show estimated 745 infection rates by week: median (thick vertical lines), 50% CrIs (box edges), and 95% CrIs 746 (whiskers).

757

759 Fig 4. Effectiveness of face covering in reducing the transmission rate and infectious 760 period. Solid lines show the estimated transmission rate (in blue, left y-axis) and infectious 761 period (in red, right y-axis) using the model-inference system incorporating interventions 762 including face covering. Surrounding areas show the 50% CrIs of model estimates. Dashed 763 lines show corresponding estimates from a linear regression model with mobility as the 764 sole predictor (i.e without accounting for face covering).

766 Fig 5. Projections of COVID-19 cases and deaths eight weeks beyond the study period. Blue 767 dots show confirmed cases by week of diagnosis and deaths by week of death, as observed 768 by the surveillance system. Blue lines show model median estimates; surrounding shades 769 show 50% and 90% CrIs. Orange lines show model projected median weekly cases and 770 deaths; surrounding shades show 50% and 90% CIs of the projection.

Supplemental Tables and Figures

Supplemental Tables

Table S1. Total number of reported cases and estimated cumulative infection rate by the week of 5/31/20. Case rate was computed as the number of total cases divided by the population size of the corresponding age group. Estimated infection rate was estimated by model-inference system, normalized to the corresponding population size; numbers are median (and 95% CrIs).

	Age Groups								
date	all	≤ 1	$1 - 4$	$5-14$	15-24	25-44	45-64	65-74	$75+$
3/1/20		2.99 (2.32, 3.86) 1.52 (1.19, 1.96)	1.56 (1.22, 2.02)	3.15 (2.47, 4.09)	2.47 (1.93, 3.22)	4(3.1, 5.15)	2.97(2.3, 3.86)	1.77 (1.37, 2.29) 1.75 (1.35, 2.27)	
3/8/20		2.25 (1.64, 2.98) 1.43 (1.11, 1.85)	1.28(1, 1.7)	2.35(1.74, 3.18)	2.02 (1.53, 2.64)	2.46 (1.57, 3.48)	2.49 (1.97, 3.12)	1.98 (1.55, 2.44)	1.84 (1.48, 2.29)
	$3/15/20$ 2.14 (1.62, 2.74) 1.63 (1.32, 2)		1.17(0.92, 1.48)	1.52(1.13, 2.06)	2.38(1.86, 3)	1.45 (0.97, 2.03)	2.99(2.3, 3.68)	2.84 (2.33, 3.51)	2.87 (2.33, 3.52)
	3/22/20 1.37 (1.08, 1.68) 1.59 (1.35, 1.89)		1.05(0.85, 1.29)	0.99(0.74, 1.3)	1.29(1.03, 1.57)	1(0.76, 1.26)	1.37(1.06, 1.71)	2.58 (2.15, 3.04)	2.52 (2.06, 3.02)
	3/29/20 0.93 (0.73, 1.13)	1.22 (0.97, 1.44)	1.06(0.82, 1.3)	0.88(0.66, 1.14)	0.96(0.75, 1.15)	0.7(0.52,0.88)	0.89(0.71, 1.06)	1.43 (1.17, 1.65)	1.46 (1.27, 1.66)
4/5/20	0.63(0.5, 0.75)	0.74(0.6, 0.88)	0.84(0.65, 1.03)	0.53(0.41, 0.68)	0.77(0.59, 0.91)	0.44(0.33, 0.56)	0.62 (0.5, 0.73)	0.9(0.79, 1.02)	0.9(0.78, 1.02)
	$4/12/20$ 0.56 (0.45, 0.67)	0.67(0.54, 0.79)	0.69(0.53, 0.84)	0.5(0.38, 0.62)	0.72(0.57, 0.83)	$0.44\,(0.34,0.55)$	0.57(0.47, 0.67)	0.63(0.55, 0.71)	0.65(0.57, 0.74)
	$4/19/20$ 0.59 (0.48, 0.7)	0.67(0.54, 0.8)	0.68(0.54, 0.81)	0.57(0.44, 0.7)	0.81(0.66, 0.93)	$0.55\,(0.42,0.68)$	0.49(0.41, 0.57)	0.6(0.52, 0.68)	0.66 (0.58, 0.74)
	$4/26/20$ 0.62 (0.5, 0.73)	0.62 (0.5, 0.75)	0.77(0.62, 0.91)	0.54(0.42, 0.67)	0.75(0.61, 0.86)	$0.58\,(0.45,0.72)$	0.6(0.5, 0.71)	0.58(0.5, 0.67)	0.63(0.56, 0.71)
5/3/20	0.66(0.54, 0.77)	0.7(0.56, 0.84)	0.78(0.62, 0.91)	0.56(0.44, 0.67)	0.72(0.61, 0.8)	0.68(0.54, 0.82)	0.68(0.56, 0.78)	0.6(0.52, 0.67)	0.55(0.48, 0.63)
	$5/10/20$ 0.82 (0.67, 0.95)	0.71(0.56, 0.86)	0.75(0.6, 0.88)	0.6(0.47, 0.73)	0.88(0.73, 0.98)	0.9(0.72, 1.06)	0.88(0.74, 0.99)	0.73(0.64, 0.81)	0.65(0.57, 0.74)
	$5/17/20$ 1.03 (0.83, 1.18)	$0.82\ (0.65, 0.99)$	0.8(0.65, 0.93)	$0.73(0.57, 0.88)$ 1.1 $(0.89, 1.24)$		1.21(0.95, 1.42)	1.1(0.91, 1.24)	0.78(0.69, 0.87)	0.81(0.72, 0.92)
	5/24/20 0.89 (0.73, 1.03)	$0.78\,(0.61,0.95)$	0.79(0.63, 0.91)	0.69(0.53, 0.83)	0.94(0.78, 1.06)	0.96(0.76, 1.13)	0.95(0.78, 1.08)	0.79(0.7, 0.88)	0.89(0.78, 0.99)
			5/31/20 0.79 (0.66, 0.91) 0.71 (0.55, 0.87) 0.71 (0.56, 0.83) 0.7 (0.55, 0.86)			$0.85(0.72, 0.95)$ $0.79(0.64, 0.93)$ $0.77(0.65, 0.89)$		0.72(0.64, 0.79)	1.04(0.93, 1.15)

Table S2. Estimated reproductive number by week and age group. Numbers are median (and interquartile range) of the posterior estimates.

Table S3. Correlation of key epidemiological parameters with population mobility during the week of March 1 – the week of May 31, 2020.

Table S4. Prior ranges for main model parameters and variables. The spatial, temporal, and age resolution of each parameter or variable, estimated in the model-inference system, is specified in the column "Resolution". Note posterior parameter estimates can extend outside the specified prior ranges. Note this is the same as Table S1 in Yang et al.(1)

Reference:

- 1. W. Yang, S. Kandula, M. Huynh, S. K. Greene, G. Van Wye, W. Li, H. T. Chan, E. McGibbon, A. Yeung, D. Olson, A. Fine, J. Shaman, Estimating the infection fatality risk of COVID-19 in New York City, March 1-May 16, 2020. *medRxiv*, 2020.2006.2027.20141689 (2020).
- 2. New York City Department of Health and Mental Hygiene.
- 3. J. Mossong, N. Hens, M. Jit, P. Beutels, K. Auranen, R. Mikolajczyk, M. Massari, S. Salmaso, G. S. Tomba, J. Wallinga, J. Heijne, M. Sadkowska-Todys, M. Rosinska, W. J. Edmunds, Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS Med* **5**, e74 (2008).
- 4. O. Li, X. Guan, P. Wu, X. Wang, L. Zhou, Y. Tong, R. Ren, K. S. M. Leung, E. H. Y. Lau, J. Y. Wong, X. Xing, N. Xiang, Y. Wu, C. Li, Q. Chen, D. Li, T. Liu, J. Zhao, M. Liu, W. Tu, C. Chen, L. Jin, R. Yang, Q. Wang, S. Zhou, R. Wang, H. Liu, Y. Luo, Y. Liu, G. Shao, H. Li, Z. Tao, Y. Yang, Z. Deng, B. Liu, Z. Ma, Y. Zhang, G. Shi, T. T. Y. Lam, J. T. Wu, G. F. Gao, B. J. Cowling, B. Yang, G. M. Leung, Z. Feng, Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *New Engl J Med.* (2020).
- 5. J. T. Wu, K. Leung, G. M. Leung, Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet*, (2020).
- 6. R. Li, S. Pei, B. Chen, Y. Song, T. Zhang, W. Yang, J. Shaman, Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science* **368**, 489-493 (2020).

- 7. J. Zhang, M. Litvinova, W. Wang, Y. Wang, X. Deng, X. Chen, M. Li, W. Zheng, L. Yi, X. Chen, Q. Wu, Y. Liang, X. Wang, J. Yang, K. Sun, I. M. Longini, Jr., M. E. Halloran, P. Wu, B. J. Cowling, S. Merler, C. Viboud, A. Vespignani, M. Ajelli, H. Yu, Evolving epidemiology and transmission dynamics of coronavirus disease 2019 outside Hubei province, China: a descriptive and modelling study. The Lancet. Infectious *diseases*, (2020).
- 8. W. Yang, J. Shaman, A simple modification for improving inference of non-linear dynamical systems. *arXiv*, 1403.6804 (2014).
- 9. R. Verity, L. C. Okell, I. Dorigatti, P. Winskill, C. Whittaker, N. Imai, G. Cuomo-Dannenburg, H. Thompson, P. G. T. Walker, H. Fu, A. Dighe, J. T. Griffin, M. Baguelin, S. Bhatia, A. Boonyasiri, A. Cori, Z. Cucunuba, R. FitzJohn, K. Gaythorpe, W. Green, A. Hamlet, W. Hinsley, D. Laydon, G. Nedjati-Gilani, S. Riley, S. van Elsland, E. Volz, H. Wang, Y. Wang, X. Xi, C. A. Donnelly, A. C. Ghani, N. M. Ferguson, Estimates of the severity of coronavirus disease 2019: a model-based analysis. The Lancet. Infectious *diseases*, (2020).

Supplemental Figures

Fig S1. Model fits of reported confirmed COVID-19 cases. Blue dots show reported number of cases by age group and week of diagnosis. Boxes show the model fitted weekly number of cases by age group. Box edges, thick middle lines, and whiskers show the $2.5th$, $25th$, $50th$, 75th, and 97.5th percentiles of model estimates.

Fig S2. Model fits of reported COVID-19 associated deaths. Blue dots show reported weekly number of deaths by age group. Boxes show the model fitted weekly number of deaths by age group. Box edges, thick middle lines, and whiskers show the $2.5th$, $25th$, $50th$, $75th$, and 97.5th percentiles of model estimates. Note that deaths among <25 year-olds are combined due to low counts.

Fig S3. Estimated case rates and infection rates by age group. Blue dots show confirmed case rates by age group and week of diagnosis. Blue boxes (left y-axis) show the model fitted weekly case rates and grey boxes (right y-axis) show the model estimated weekly infection rates by age group. Box edges, thick middle lines, and whiskers show the $2.5th$, 25th, 50th, 75th, and 97.5th percentiles of model estimates.

Fig S4. Sensitivity analysis on the effectiveness of reducing contact rate by neighborhood. There are 42 United Hospital Fund (UHF) neighborhoods in NYC. (A) shows the changes in human mobility during the pandemic by neighborhood (each colored line). The reductions were substantial in all neighborhoods but to varying degrees. (B) shows the estimated R_t (combining all ages) for each week and neighborhood (each colored line). Note these estimates did not account for changes in susceptibility so as to restrict to changes due to interventions. (C) shows the adjusted r^2 of linear regression model fitting the mobility data (A) to the R_t estimates (B), for each neighborhood, per Eqn 2 in the main text. Adjusted r^2 for most neighborhoods was >0.9. (D) shows the estimated reduction in R_t by the Week of April 12, 2020 based on the Eqn 3 in the main text, for each neighborhood. The histogram is based on the median estimated reduction in R_t .

Fig S5. Projections of COVID-19 cases by age group eight weeks beyond the study period. Blue dots show observed confirmed cases by week of diagnosis (those after the Week of 5/31/2020 were not used in the model). Blue lines show model median estimates; surrounding shades show 50% and 90% CrIs. Orange lines show model projected median weekly cases and deaths; surrounding shades show 50% and 90% CIs of the projection.

Fig S6. Projections of COVID-19 associated deaths by age group eight weeks beyond the study period. Blue dots show observed weekly deaths (those after the Week of 5/31/2020 were not used in the model). Blue lines show model median estimates; surrounding shades show 50% and 90% CrIs. Orange lines show model projected median weekly cases and deaths; surrounding shades show 50% and 90% CIs of the projection.

Fig S7. Comparing long term projection with available observations. The projections on COVID-19 cases, hospitalizations and deaths were generated on June 30, 2020, using case and mortality data up to June 26, 2020. These projections were generated at the time using a simpler model without age grouping but otherwise the same methodology presented in the study. Results were posted online on June 30, 2020 (https://github.com/wan-yang/reopening analysis/tree/master/test3_occupancy). Various policy scenarios were tested and here we used the one labeled "sce2fix_asIs" for the "intervention" identifier, which has been the closest to implemented interventions thus far (main settings were capping capacity for all industries including schools at 50%; no shutdown). For comparison with observations thus far, we used data published by the New York City Department of Health and Mental Hygiene, as of Dec 27, 2020 (https://raw.githubusercontent.com/nychealth/coronavirusdata/master/trends/data-by-day.csv). Red line and shaded area show our projections (median and interquartile range) and blue dots show corresponding observations from the New York City Department of Health and Mental Hygiene. Note that estimated infectiondetection rates using PCR tests during June – Dec 2020 have been similar to those in June. However, hospitalization rates and estimated infection-fatality-risk in months after June 2020 have been lower than that during the spring wave; these lower hospitalization rates and mortality risks contribute to the lower observed numbers than our projections.

