

Piotr T. Chruściel, Sebastian J. Szybka

On the lag between deaths and infections in the first phase of the Covid-19 pandemic
supplementary material

Introduction

We present here the time series and fits used for the analysis in the main accompanying paper. The analysis is concerned with the first 180 days of the epidemic, using data available on the John Hopkins University server on December 6, 2020. The best fits, and the data of the end of the first phase of the epidemic, are obtained as described in <https://www.medrxiv.org/content/early/2020/08/30/2020.08.24.20181214>.

List of countries and states used in our analysis

a) global

```
Out[1]= {{"Austria"}, {"Belgium"}, {"Denmark"}, {"France"}, {"Germany"}, {"Hungary"}, {"Ireland"}, {"Italy"}, {"Netherlands"}, {"Poland"}, {"Portugal"}, {"Romania"}, {"Spain"}, {"Sweden"}, {"United Kingdom"}, {"US"}, {"Afghanistan"}, {"Algeria"}, {"Argentina"}, {"Armenia"}, {"Bangladesh"}, {"Bolivia"}, {"Brazil"}, {"Chile"}, {"Colombia"}, {"Dominican Republic"}, {"Ecuador"}, {"Egypt"}, {"Guatemala"}, {"Honduras"}, {"India"}, {"Indonesia"}, {"Iran"}, {"Iraq"}, {"Japan"}, {"Mexico"}, {"Moldova"}, {"Nigeria"}, {"Pakistan"}, {"Panama"}, {"Peru"}, {"Philippines"}, {"Russia"}, {"Saudi Arabia"}, {"South Africa"}, {"Sudan"}, {"Switzerland"}, {"Turkey"}, {"Ukraine"}, {"Hubei, China"}, {"Ontario, Canada"}, {"Quebec, Canada"}}
```

b) US

```
Out[1]= {"Maricopa, Arizona"}, {"Los Angeles, California"}, {"Riverside, California"}, {"Fairfield, Connecticut"}, {"Hartford, Connecticut"}, {"New Haven, Connecticut"}, {"District of Columbia, District of Columbia"}, {"Miami-Dade, Florida"}, {"Palm Beach, Florida"}, {"Cook, Illinois"}, {"Marion, Indiana"}, {"Orleans, Louisiana"}, {"Baltimore, Maryland"}, {"Montgomery, Maryland"}, {"Prince George's, Maryland"}, {"Bristol, Massachusetts"}, {"Essex, Massachusetts"}, {"Hampden, Massachusetts"}, {"Middlesex, Massachusetts"}, {"Norfolk, Massachusetts"}, {"Plymouth, Massachusetts"}, {"Suffolk, Massachusetts"}, {"Worcester, Massachusetts"}, {"Macomb, Michigan"}, {"Oakland, Michigan"}, {"Wayne, Michigan"}, {"Hennepin, Minnesota"}, {"St. Louis, Missouri"}, {"Clark, Nevada"}, {"Bergen, New Jersey"}, {"Camden, New Jersey"}, {"Essex, New Jersey"}, {"Hudson, New Jersey"}, {"Mercer, New Jersey"}, {"Middlesex, New Jersey"}, {"Monmouth, New Jersey"}, {"Morris, New Jersey"}, {"Ocean, New Jersey"}, {"Passaic, New Jersey"}, {"Somerset, New Jersey"}, {"Union, New Jersey"}, {"Bronx, New York"}, {"Erie, New York"}, {"Kings, New York"}, {"Nassau, New York"}, {"New York, New York"}, {"Queens, New York"}, {"Richmond, New York"}, {"Rockland, New York"}, {"Suffolk, New York"}, {"Westchester, New York"}, {"Bucks, Pennsylvania"}, {"Delaware, Pennsylvania"}, {"Montgomery, Pennsylvania"}, {"Philadelphia, Pennsylvania"}, {"Providence, Rhode Island"}, {"Harris, Texas"}, {"Fairfax, Virginia"}, {"King, Washington"}
```

Plots, fits and integral averages, not including US counties


Austria


lag from integral overlap between 12 and 13 days, maximum overlap on day 13

lag from integral overlap, using data averaged over 7 days, between 11 and 14 days, maximum overlap on day 12

lag from fits to the function I: 14.1083 days

lag from fits to the function I (averaged data): 14.1546 days






Parameters of fits to cases data.

The best fit ends on Wed 13 May 2020.


α	β	k	τ	σ		
0.322 ± 0.0071	0.0705 ± 0.00793	16200 ± 75	28.7 ± 0.234	22.3 ± 1.18		
badness	α	β	k	τ	σ	quality
2.1%	error	2.2%	11.%	0.46%	0.81%	5.3% 12.6%



Parameters of fits to averaged cases data.

The best fit ends on Tue 12 May 2020.


α	β	k	τ	σ		
0.31 ± 0.00351	0.0736 ± 0.00376	16100 ± 41.7	28.6 ± 0.142	22.7 ± 0.633		
badness	α	β	k	τ	σ	quality
0.65%	error	1.1%	5.1%	0.26%	0.5%	2.8% 6.09%



Parameters of fits to death data.

The best fit ends on Wed 27 May 2020.


α	β	k	τ	σ		
0.409 ± 0.0283	0.113 ± 0.00206	642 ± 1.54	26.6 ± 0.136	11.1 ± 0.365		
badness	α	β	k	τ	σ	quality
1.8%	error	6.9%	1.8%	0.24%	0.51%	3.3% 14.6%



Parameters of fits to averaged death data.

The best fit ends on Wed 27 May 2020.

α	β	k	τ	σ			
0.39 ± 0.0139	0.113 ± 0.00112	$641. \pm 0.878$	28.5 ± 0.0742	12.7 ± 0.206			
badness	α	β	k	τ	σ	quality	
1.%	error	3.6%	1.%	0.14%	0.26%	1.6%	7.62%




Belgium

lag from integral overlap between 6 and 7 days, maximum overlap on day 6

lag from integral overlap, using data averaged over 7 days, between 3 and 9 days, maximum overlap on day 6

lag from fits to the function I: 4.76359 days

lag from fits to the function I (averaged data): 3.14838 days




Parameters of fits to cases data.

The best fit ends on Sat 11 Jul 2020.

α	β	k	τ	σ
0.188 ± 0.00417	0.0473 ± 0.00503	$62200. \pm 165.$	33.6 ± 0.461	33.6 ± 1.14

badness	α	β	k	τ	σ	quality
2.6%	error	2.2%	11%	0.27%	1.4%	3.4% 12.%




Parameters of fits to averaged cases data.

The best fit ends on Thu 14 May 2020.

α	β	k	τ	σ
0.309 ± 0.0109	0.0965 ± 0.00143	$55800. \pm 187.$	44.6 ± 0.0882	25.5 ± 0.317

badness	α	β	k	τ	σ	quality
0.65%	error	3.5%	1.5%	0.34%	0.2%	1.2% 7.44%




Parameters of fits to death data.

The best fit ends on Fri 7 Aug 2020.

α	β	k	τ	σ
0.25 ± 0.00353	0.0648 ± 0.00106	$9820. \pm 6.9$	30.7 ± 0.451	$29. \pm 0.205$

badness	α	β	k	τ	σ	quality
1.6%	error	1.4%	1.6%	0.07%	1.5%	0.71% 6.86%




Parameters of fits to averaged death data.

The best fit ends on Wed 29 Jul 2020.

α	β	k	τ	σ
0.246 ± 0.00195	0.0659 ± 0.00245	9800 ± 4.29	29.7 ± 0.125	32 ± 0.255

badness	α	β	k	τ	σ	quality
0.82%	error	0.79%	3.7%	0.044%	0.42%	0.8% 3.71%




Denmark

lag from integral overlap between 0 and 1 days, maximum overlap on day 0

lag from integral overlap, using data averaged over 7 days, between -1 and 3 days, maximum overlap on day 1

lag from fits to the function I: -0.98422 days

lag from fits to the function I (averaged data): -0.883999 days




Parameters of fits to cases data.

The best fit ends on Wed 29 Jul 2020.


α	β	k	τ	σ
0.142 ± 0.00374	0.029 ± 0.00457	13800 ± 85.8	33.6 ± 0.559	33.6 ± 1.54

badness	α	β	k	τ	σ	quality
2.4%	error	2.6%	16.%	0.62%	1.7%	4.6% 14.3%

Denmark



Denmark




Parameters of fits to averaged cases data.

The best fit ends on Tue 28 Jul 2020.


α	β	k	τ	σ
0.141 ± 0.00323	0.0299 ± 0.00389	13700 ± 75.3	35.4 ± 0.519	35.4 ± 1.39

badness	α	β	k	τ	σ	quality
1.7%	error	2.3%	13.%	0.55%	1.5%	3.9% 12.%

Denmark



Denmark




Parameters of fits to death data.

The best fit ends on Wed 29 Jul 2020.

α	β	k	τ	σ
0.261 ± 0.0082	0.0628 ± 0.00109	610 ± 0.821	29.5 ± 0.327	17 ± 0.298


badness	α	β	k	τ	σ	quality
2.5%	error	3.1%	1.7%	0.13%	1.1%	1.8% 10.3%



Parameters of fits to averaged death data.

The best fit ends on Tue 28 Jul 2020.

α	β	k	τ	σ			
0.254 ± 0.00672	0.0628 ± 0.00095	$609. \pm 0.726$	30.5 ± 0.289	17.8 ± 0.269			
badness	error	α	β	k	τ	σ	quality
2.%		2.7%	1.5%	0.12%	0.95%	1.5%	8.77%




France

lag from integral overlap between 3 and 3 days, maximum overlap on day 3

lag from integral overlap, using data averaged over 7 days, between 2 and 4 days, maximum overlap on day 3

lag from fits to the function I: 4.01184 days

lag from fits to the function I (averaged data): 7.40324 days




Parameters of fits to cases data.

The best fit ends on Mon 27 Jul 2020.

α	β	k	τ	σ
0.217 ± 0.0162	0.0546 ± 0.0203	196000 ± 987	67.8 ± 1.26	67.9 ± 3.18

badness	α	β	k	τ	σ	quality
7.4%	error	7.5%	37.3%	0.5%	1.9%	4.7%




Parameters of fits to averaged cases data.

The best fit ends on Thu 9 Apr 2020.

α	β	k	τ	σ
0.482 ± 0.0313	0.175 ± 0.000988	101000 ± 493	69 ± 0.0597	43.9 ± 0.372

badness	α	β	k	τ	σ	quality
0.56%	error	6.5%	0.56%	0.49%	0.087%	0.85%




Parameters of fits to death data.

The best fit ends on Tue 11 Aug 2020.

α	β	k	τ	σ
0.242 ± 0.00333	0.0553 ± 0.00426	30100 ± 24.5	39.8 ± 0.177	38.5 ± 0.527


badness	α	β	k	τ	σ	quality
2.2%	error	1.4%	7.7%	0.082%	0.44%	1.4%



Parameters of fits to averaged death data.

The best fit ends on Tue 14 Jul 2020.

α	β	k	τ	σ			
0.243 ± 0.00199	0.0591 ± 0.0025	29900 ± 19.5	42.2 ± 0.116	42.8 ± 0.288			
badness	α	β	k	τ	σ	quality	
0.65%	error	0.82%	4.2%	0.065%	0.28%	0.67%	3.31%




Germany

lag from integral overlap between 13 and 13 days, maximum overlap on day 13

lag from integral overlap, using data averaged over 7 days, between 11 and 16 days, maximum overlap on day 13

lag from fits to the function I: 14.8609 days

lag from fits to the function I (averaged data): 16.5692 days




Parameters of fits to cases data.

The best fit ends on Mon 15 Jun 2020.

α	β	k	τ	σ
0.241 \pm 0.003	0.0521 \pm 0.00373	190000. \pm 355.	59.6 \pm 0.161	59.6 \pm 0.445

badness	α	β	k	τ	σ	quality
1.2%	error	1.2%	7.2%	0.19%	0.27%	0.75%




Parameters of fits to averaged cases data.

The best fit ends on Wed 20 May 2020.

α	β	k	τ	σ
0.247 \pm 0.00185	0.0625 \pm 0.00211	184000. \pm 381.	60.5 \pm 0.129	63.1 \pm 0.233

badness	α	β	k	τ	σ	quality
0.46%	error	0.75%	3.4%	0.21%	0.21%	0.37%




Parameters of fits to death data.

The best fit ends on Thu 13 Aug 2020.

α	β	k	τ	σ
0.191 \pm 0.00351	0.0481 \pm 0.00428	9160. \pm 12.8	32.4 \pm 0.371	29.5 \pm 1.08

badness	α	β	k	τ	σ	quality
2.9%	error	1.8%	8.9%	0.14%	1.1%	3.7%




Parameters of fits to averaged death data.

The best fit ends on Sat 9 May 2020.

α	β	k	τ	σ
0.348 ± 0.0111	0.108 ± 0.00147	8000 ± 36.6	38.9 ± 0.0722	19.9 ± 0.249

badness	α	β	k	τ	σ	quality
0.71%	error	3.2%	1.4%	0.46%	0.19%	1.3% 7.15%




Hungary

lag from integral overlap between 0 and 7 days, maximum overlap on day 7

lag from integral overlap, using data averaged over 7 days, between 5 and 10 days, maximum overlap on day 8

lag from fits to the function I: 4.00293 days

lag from fits to the function I (averaged data): 4.20852 days





Parameters of fits to cases data.

The best fit ends on Thu 16 Jul 2020.

α	β	k	τ	σ
0.162 ± 0.00386	0.0449 ± 0.00452	4290 ± 13.8	36.2 ± 0.685	39.5 ± 1.29

badness	α	β	k	τ	σ	quality
2.1%	error	2.4%	10.%	0.32%	1.9%	3.3% 12.1%





Parameters of fits to averaged cases data.

The best fit ends on Fri 17 Jul 2020.

α	β	k	τ	σ
0.161 ± 0.00279	0.0457 ± 0.00326	4280 ± 9.96	35.9 ± 0.514	39.9 ± 0.909

badness	α	β	k	τ	σ	quality
1.1%	error	1.7%	7.1%	0.23%	1.4%	2.3% 8.34%




Parameters of fits to death data.

The best fit ends on Sun 9 Aug 2020.

α	β	k	τ	σ
0.209 ± 0.00383	0.0531 ± 0.000672	602 ± 0.647	37.6 ± 0.283	24.2 ± 0.249


badness	α	β	k	τ	σ	quality
2.2%	error	1.8%	1.3%	0.11%	0.75%	1.% 7.14%



Parameters of fits to averaged death data.

The best fit ends on Tue 11 Aug 2020.

α	β	k	τ	σ			
0.205 ± 0.00262	0.0529 ± 0.000482	602 ± 0.464	40.5 ± 0.208	27.1 ± 0.184			
badness	α	β	k	τ	σ	quality	
0.77%	error	1.3%	0.91%	0.077%	0.51%	0.68%	4.23%




Ireland

lag from integral overlap between 14 and 14 days, maximum overlap on day 14

lag from integral overlap, using data averaged over 7 days, between 7 and 10 days, maximum overlap on day 9

lag from fits to the function I: 5.58788 days

lag from fits to the function I (averaged data): 5.4144 days





Parameters of fits to cases data.

The best fit ends on Sun 2 Aug 2020.

α	β	k	τ	σ
0.202 ± 0.00738	0.0709 ± 0.00822	$25600. \pm 41.7$	35.9 ± 1.44	35.9 ± 2.51

badness	α	β	k	τ	σ	quality
4.7%	error	3.6%	12.%	0.16%	4.%	7.%





Parameters of fits to averaged cases data.

The best fit ends on Thu 9 Apr 2020.

α	β	k	τ	σ
0.368 ± 0.00693	0.104 ± 0.00761	$16900. \pm 1470.$	19.8 ± 0.245	44.2 ± 1.46

badness	α	β	k	τ	σ	quality
0.34%	error	1.9%	7.3%	8.7%	1.2%	3.3%




Parameters of fits to death data.

The best fit ends on Tue 6 Oct 2020.

α	β	k	τ	σ
0.22 ± 0.00913	0.0596 ± 0.00313	$1770. \pm 2.75$	32.3 ± 1.8	32.2 ± 0.766


badness	α	β	k	τ	σ	quality
5.4%	error	4.2%	5.2%	0.16%	5.6%	2.4%



Parameters of fits to averaged death data.

The best fit ends on Sat 26 Sep 2020.

α	β	k	τ	σ			
0.216 ± 0.00741	0.0603 ± 0.00913	1770 ± 2.4	33.1 ± 0.689	33.1 ± 1.57			
badness	α	β	k	τ	σ	quality	
3.9%	error	3.4%	15.%	0.14%	2.1%	4.7%	17.8%




Italy

lag from integral overlap between 1 and 7 days, maximum overlap on day 6

lag from integral overlap, using data averaged over 7 days, between 1 and 6 days, maximum overlap on day 4

lag from fits to the function I: 4.43773 days

lag from fits to the function I (averaged data): 3.81535 days




Parameters of fits to cases data.

The best fit ends on Tue 16 Jun 2020.

α	β	k	τ	σ
0.252 ± 0.00293	0.0636 ± 0.000406	$238000. \pm 178.$	62.6 ± 0.0932	46.8 ± 0.118

badness	α	β	k	τ	σ	quality
0.77%	error	1.2%	0.64%	0.075%	0.15%	0.25%




Parameters of fits to averaged cases data.

The best fit ends on Tue 16 Jun 2020.

α	β	k	τ	σ
0.246 ± 0.00209	0.0636 ± 0.000308	$238000. \pm 138.$	62.6 ± 0.0699	46.6 ± 0.0916

badness	α	β	k	τ	σ	quality
0.35%	error	0.85%	0.48%	0.058%	0.11%	0.2%




Parameters of fits to death data.

The best fit ends on Sat 1 Aug 2020.

α	β	k	τ	σ
0.233 ± 0.00232	0.0555 ± 0.000302	$35100. \pm 15.7$	44.5 ± 0.113	30.1 ± 0.105


badness	α	β	k	τ	σ	quality
0.76%	error	1.%	0.54%	0.045%	0.25%	0.35%



Parameters of fits to averaged death data.

The best fit ends on Wed 10 Jun 2020.

α	β	k	τ	σ			
0.241 ± 0.00176	0.0591 ± 0.00031	34700 ± 28.4	45.9 ± 0.0632	30.2 ± 0.0832			
badness	α	β	k	τ	σ	quality	
0.36%	error	0.73%	0.52%	0.082%	0.14%	0.28%	2.11%




Netherlands

lag from integral overlap between 5 and 6 days, maximum overlap on day 5

lag from integral overlap, using data averaged over 7 days, between 2 and 7 days, maximum overlap on day 4

lag from fits to the function I: -2.95931 days

lag from fits to the function I (averaged data): -3.4936 days





Parameters of fits to cases data.

The best fit ends on Wed 20 May 2020.

α	β	k	τ	σ
0.314 ± 0.0148	0.0938 ± 0.00133	45100 ± 115	41.1 ± 0.104	20.7 ± 0.35

badness	α	β	k	τ	σ	quality
1.2%	error	4.7%	1.4%	0.26%	0.25%	1.7% 9.52%





Parameters of fits to averaged cases data.

The best fit ends on Tue 19 May 2020.

α	β	k	τ	σ
0.309 ± 0.0103	0.0944 ± 0.000979	44900 ± 93.3	$44. \pm 0.0708$	23.2 ± 0.258

badness	α	β	k	τ	σ	quality
0.63%	error	3.3%	1.%	0.21%	0.16%	1.1% 6.48%




Parameters of fits to death data.

The best fit ends on Tue 11 Aug 2020.

α	β	k	τ	σ
0.291 ± 0.00649	0.0762 ± 0.000622	6130 ± 3.44	37.3 ± 0.129	21.6 ± 0.179


badness	α	β	k	τ	σ	quality
1.8%	error	2.2%	0.82%	0.056%	0.35%	0.83% 6.12%



Parameters of fits to averaged death data.

The best fit ends on Sat 30 May 2020.

α	β	k	τ	σ			
0.317 ± 0.00344	0.0829 ± 0.000395	6030 ± 5.35	39.5 ± 0.0359	22.2 ± 0.0817			
badness	α	β	k	τ	σ	quality	
0.4%	error	1.1%	0.48%	0.089%	0.091%	0.37%	2.51%




Poland

lag from integral overlap between 5 and 11 days, maximum overlap on day 10

lag from integral overlap, using data averaged over 7 days, between 5 and 14 days, maximum overlap on day 9

lag from fits to the function I: 11.0418 days

lag from fits to the function I (averaged data): 10.6938 days




Parameters of fits to cases data.

The best fit ends on Fri 17 Jul 2020.

α	β	k	τ	σ
0.191 ± 0.00406	0.0324 ± 0.000236	47400 ± 225	86.4 ± 0.329	28.6 ± 0.188

badness	α	β	k	τ	σ	quality
1.2%	error	2.1%	0.73%	0.47%	0.38%	0.66%




Parameters of fits to averaged cases data.

The best fit ends on Sun 19 Jul 2020.

α	β	k	τ	σ
0.188 ± 0.0032	0.0324 ± 0.000195	47400 ± 188	88.3 ± 0.274	30.5 ± 0.154

badness	α	β	k	τ	σ	quality
0.84%	error	1.7%	0.6%	0.4%	0.31%	0.5%




Parameters of fits to death data.

The best fit ends on Thu 13 Aug 2020.

α	β	k	τ	σ
0.157 ± 0.00414	0.0238 ± 0.000519	2110 ± 20.5	77 ± 0.787	34.1 ± 0.312


badness	α	β	k	τ	σ	quality
1.7%	error	2.6%	2.2%	0.97%	1.%	0.91%



Parameters of fits to averaged death data.

The best fit ends on Wed 12 Aug 2020.

α	β	k	τ	σ			
0.158 ± 0.00365	0.0245 ± 0.000477	$2070. \pm 17.9$	76.6 ± 0.674	34.8 ± 0.276			
badness	α	β	k	τ	σ	quality	
1.3%	error	2.3%	1.9%	0.87%	0.88%	0.79%	8.06%




Portugal

lag from integral overlap between 3 and 7 days, maximum overlap on day 5

lag from integral overlap, using data averaged over 7 days, between 3 and 9 days, maximum overlap on day 5

lag from fits to the function I: 10.2502 days

lag from fits to the function I (averaged data): 10.7209 days




Parameters of fits to cases data.

The best fit ends on Fri 4 Sep 2020.


α	β	k	τ	σ
0.191 ± 0.0055	0.0164 ± 0.000275	$73300. \pm 813.$	$103. \pm 1.47$	32.2 ± 0.217

badness	α	β	k	τ	σ	quality
1.8%	error	2.9%	1.7%	1.1%	1.4%	0.67% 9.61%

Portugal



Portugal




Parameters of fits to averaged cases data.

The best fit ends on Sat 9 May 2020.


α	β	k	τ	σ
0.323 ± 0.00697	0.0893 ± 0.00145	$28500. \pm 121.$	39.5 ± 0.0824	24.7 ± 0.211

badness	α	β	k	τ	σ	quality
0.69%	error	2.2%	1.6%	0.42%	0.21%	0.85% 5.96%

Portugal



Portugal




Parameters of fits to death data.

The best fit ends on Fri 15 May 2020.

α	β	k	τ	σ
0.351 ± 0.0133	0.0921 ± 0.00135	$1270. \pm 5.48$	30.3 ± 0.0916	10.9 ± 0.24


badness	α	β	k	τ	σ	quality
1.1%	error	3.8%	1.5%	0.43%	0.3%	2.2% 9.29%



Parameters of fits to averaged death data.

The best fit ends on Thu 23 Apr 2020.

α	β	k	τ	σ			
0.451 ± 0.00599	0.121 ± 0.000926	1030 ± 6.08	27.3 ± 0.101	9.15 ± 0.0747			
badness	α	β	k	τ	σ	quality	
0.22%	error	1.3%	0.77%	0.59%	0.37%	0.82%	4.09%




Romania

lag from integral overlap between 16 and 16 days, maximum overlap on day 16

lag from integral overlap, using data averaged over 7 days, between 4 and 19 days, maximum overlap on day 13

lag from fits to the function I: -3.24463 days

lag from fits to the function I (averaged data): 19.8699 days




Parameters of fits to cases data.

The best fit ends on Wed 3 Jun 2020.


α	β	k	τ	σ
0.231 ± 0.00568	0.0581 ± 0.000842	$21600 \pm 103.$	57.9 ± 0.141	32 ± 0.266

badness	α	β	k	τ	σ	quality
0.98%	error	2.5%	1.5%	0.48%	0.24%	0.83% 6.44%

Romania



Romania




Parameters of fits to averaged cases data.

The best fit ends on Sat 30 May 2020.


α	β	k	τ	σ
0.237 ± 0.00338	0.0608 ± 0.000543	21100 ± 71.5	59.2 ± 0.0966	33.2 ± 0.157

badness	α	β	k	τ	σ	quality
0.49%	error	1.4%	0.89%	0.34%	0.16%	0.47% 3.78%

Romania



Romania




Parameters of fits to death data.

The best fit ends on Sat 6 Jun 2020.

α	β	k	τ	σ
0.348 ± 0.0166	0.072 ± 0.000663	1400 ± 4.77	39.1 ± 0.116	10.2 ± 0.24


badness	α	β	k	τ	σ	quality
1.1%	error	4.8%	0.92%	0.34%	0.3%	2.4% 9.85%



Parameters of fits to averaged death data.

The best fit ends on Wed 20 May 2020.

α	β	k	τ	σ			
0.31 ± 0.00551	0.0666 ± 0.000587	1490 ± 10.4	43.1 ± 0.224	12.6 ± 0.114			
badness	α	β	k	τ	σ	quality	
0.42%	error	1.8%	0.88%	0.7%	0.52%	0.9%	5.2%




Spain

lag from integral overlap between 0 and 0 days, maximum overlap on day 0

lag from integral overlap, using data averaged over 7 days, between 1 and 5 days, maximum overlap on day 3

lag from fits to the function I: -0.415911 days

lag from fits to the function I (averaged data): -0.309878 days





Parameters of fits to cases data.

The best fit ends on Fri 26 Jun 2020.

α	β	k	τ	σ
0.228 ± 0.00532	0.0471 ± 0.00646	$248000. \pm 855.$	49.2 ± 0.312	42.6 ± 1.36

badness	α	β	k	τ	σ	quality
4.5%	error	2.3%	14.%	0.34%	0.63%	3.2% 13.1%





Parameters of fits to averaged cases data.

The best fit ends on Mon 29 Jun 2020.

α	β	k	τ	σ
0.221 ± 0.00392	0.0459 ± 0.00471	$249000. \pm 689.$	52.3 ± 0.25	44.9 ± 1.13

badness	α	β	k	τ	σ	quality
2.4%	error	1.8%	10.%	0.28%	0.48%	2.5% 8.99%




Parameters of fits to death data.

The best fit ends on Thu 21 May 2020.

α	β	k	τ	σ
0.315 ± 0.00418	0.0683 ± 0.000853	$29300. \pm 71.8$	34.4 ± 0.0959	23.6 ± 0.111


badness	α	β	k	τ	σ	quality
1.2%	error	1.3%	1.2%	0.24%	0.28%	0.47% 4.8%



Parameters of fits to averaged death data.

The best fit ends on Fri 22 May 2020.

α	β	k	τ	σ			
0.3 ± 0.00168	0.0671 ± 0.000397	29400 ± 35.7	35.3 ± 0.0437	24.6 ± 0.0529			
badness	α	β	k	τ	σ	quality	
0.26%	error	0.56%	0.59%	0.12%	0.12%	0.22%	1.88%




Sweden

lag from integral overlap between -9 and -2 days, maximum overlap on day -6

lag from integral overlap, using data averaged over 7 days, between -11 and 0 days, maximum overlap on day -5

lag from fits to the function I: -16.8232 days

lag from fits to the function I (averaged data): -17.9024 days





Parameters of fits to cases data.

The best fit ends on Wed 27 May 2020.

α	β	k	τ	σ
0.147 ± 0.00528	0.0389 ± 0.00623	$55500. \pm 2450.$	34.2 ± 0.988	75.8 ± 2.22

badness	α	β	k	τ	σ	quality
1.%	error	3.6%	16.%	4.4%	2.9%	2.9% 18.1%





Parameters of fits to averaged cases data.

The best fit ends on Sat 30 May 2020.

α	β	k	τ	σ
0.144 ± 0.00273	0.0375 ± 0.00318	$57600. \pm 1560.$	37.6 ± 0.539	80.7 ± 1.42

badness	α	β	k	τ	σ	quality
0.46%	error	1.9%	8.5%	2.7%	1.4%	1.8% 9.91%




Parameters of fits to death data.

The best fit ends on Mon 28 Sep 2020.

α	β	k	τ	σ
0.206 ± 0.0027	0.0432 ± 0.000212	$5870. \pm 2.92$	48.7 ± 0.107	25.2 ± 0.126


badness	α	β	k	τ	σ	quality
0.92%	error	1.3%	0.49%	0.05%	0.22%	0.5% 3.49%



Parameters of fits to averaged death data.

The best fit ends on Wed 3 Jun 2020.

α	β	k	τ	σ			
0.238 ± 0.00234	0.0548 ± 0.000455	5320 ± 16.6	46.7 ± 0.0943	23.6 ± 0.109			
badness	α	β	k	τ	σ	quality	
0.36%	error	0.98%	0.83%	0.31%	0.2%	0.46%	3.15%




United Kingdom

lag from integral overlap between 1 and 1 days, maximum overlap on day 1

lag from integral overlap, using data averaged over 7 days, between 0 and 6 days, maximum overlap on day 3

lag from fits to the function I: 0.996832 days

lag from fits to the function I (averaged data): -1.64298 days




Parameters of fits to cases data.

The best fit ends on Wed 10 Jun 2020.

α	β	k	τ	σ
0.236 ± 0.00435	0.0625 ± 0.00049	$278000. \pm 508.$	82.3 ± 0.0682	56.9 ± 0.184

badness	α	β	k	τ	σ	quality
1.%	error	1.8%	0.78%	0.18%	0.083%	0.32%




Parameters of fits to averaged cases data.

The best fit ends on Wed 10 Jun 2020.

α	β	k	τ	σ
0.232 ± 0.00304	0.0625 ± 0.000371	$278000. \pm 405.$	82.3 ± 0.0499	56.6 ± 0.138

badness	α	β	k	τ	σ	quality
0.54%	error	1.3%	0.59%	0.15%	0.061%	0.24%




Parameters of fits to death data.

The best fit ends on Fri 4 Sep 2020.

α	β	k	τ	σ
0.222 ± 0.00263	0.0522 ± 0.00041	$41400. \pm 21.5$	42.9 ± 0.198	32.1 ± 0.138


badness	α	β	k	τ	σ	quality
1.5%	error	1.2%	0.79%	0.052%	0.46%	0.43%



Parameters of fits to averaged death data.

The best fit ends on Fri 15 May 2020.

α	β	k	τ	σ			
0.274 ± 0.00152	0.0779 ± 0.00185	37100 ± 78.6	28.8 ± 0.0828	43.9 ± 0.0354			
badness	α	β	k	τ	σ	quality	
0.18%	error	0.55%	2.4%	0.21%	0.29%	0.081%	1.87%




US

lag from integral overlap between -1 and 7 days, maximum overlap on day 6

lag from integral overlap, using data averaged over 7 days, between 0 and 7 days, maximum overlap on day 4

lag from fits to the function I: 6.72005 days

lag from fits to the function I (averaged data): 6.4673 days




Parameters of fits to cases data.

The best fit ends on Thu 14 May 2020.

α	β	k	τ	σ
0.271 ± 0.0032	0.0606 ± 0.000644	$1860000. \pm 12400.$	92.6 ± 0.221	66.7 ± 0.112

badness	α	β	k	τ	σ	quality
0.46%	error	1.2%	1.1%	0.67%	0.24%	0.17%




Parameters of fits to averaged cases data.

The best fit ends on Sun 17 May 2020.

α	β	k	τ	σ
0.262 ± 0.00158	0.0596 ± 0.000342	$1880000. \pm 6810.$	93.9 ± 0.122	67.7 ± 0.0621

badness	α	β	k	τ	σ	quality
0.24%	error	0.6%	0.57%	0.36%	0.13%	0.092%




Parameters of fits to death data.

The best fit ends on Mon 1 Jun 2020.

α	β	k	τ	σ
0.251 ± 0.00516	0.0597 ± 0.000935	$122000. \pm 669.$	57.2 ± 0.151	35.2 ± 0.214


badness	α	β	k	τ	σ	quality
0.8%	error	2.1%	1.6%	0.55%	0.26%	0.61%



Parameters of fits to averaged death data.

The best fit ends on Sun 31 May 2020.

α	β	k	τ	σ			
0.25 ± 0.002	0.0609 ± 0.000406	$120000.0 \pm 303.$	59.8 ± 0.0677	37.8 ± 0.0897			
badness	error	α	β	k	τ	σ	quality
0.29%	error	0.8%	0.67%	0.25%	0.11%	0.24%	2.36%



Integral averages, not including US counties


Afghanistan

no clear lag in the first six months of the pandemic from integral overlap


no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

no meaningful fits to the function I available


no meaningful fits to the function I available (averaged data)




averaged new cases, Afghanistan



averaged new deaths, Afghanistan



no clear lag in the first six months of the pandemic



Algeria


lag from integral overlap between 1 and 7 days, maximum overlap on day 3

lag from integral overlap, using data averaged over 7 days, between 3 and 8 days, maximum overlap on day 5


no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)


new cases, Algeria




new deaths, Algeria




lag 3 in [1,7]




averaged new cases, Algeria



averaged new deaths, Algeria



lag 5 in [3,8]



Argentina


no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days


no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)


new cases, Argentina




new deaths, Argentina




no clear lag in the first six months of the pandemic




averaged new cases, Argentina



averaged new deaths, Argentina



no clear lag in the first six months of the pandemic



Armenia


no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days


no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)


new cases, Armenia




new deaths, Armenia




no clear lag in the first six months of the pandemic




averaged new cases, Armenia



averaged new deaths, Armenia



no clear lag in the first six months of the pandemic



Bangladesh


no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days


no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)


new cases, Bangladesh




new deaths, Bangladesh




no clear lag in the first six months of the pandemic




averaged new cases, Bangladesh



averaged new deaths, Bangladesh



no clear lag in the first six months of the pandemic




Bolivia

no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




Brazil

no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




Chile


no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

no meaningful fits to the function I available


no meaningful fits to the function I available (averaged data)






Colombia

no clear lag in the first six months of the pandemic from integral overlap
no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days
no meaningful fits to the function I available
no meaningful fits to the function I available (averaged data)




Dominican Republic

no clear lag in the first six months of the pandemic from integral overlap
no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days
no meaningful fits to the function I available
no meaningful fits to the function I available (averaged data)




Ecuador

lag from integral overlap between 16 and 16 days, maximum overlap on day 16
 lag from integral overlap, using data averaged over 7 days, between 18 and 18 days, maximum overlap on day 18
 no meaningful fits to the function I available
 no meaningful fits to the function I available (averaged data)




Egypt

no clear lag in the first six months of the pandemic from integral overlap
 no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days
 no meaningful fits to the function I available
 no meaningful fits to the function I available (averaged data)



Guatemala

no clear lag in the first six months of the pandemic from integral overlap
 no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days
 no meaningful fits to the function I available
 no meaningful fits to the function I available (averaged data)




Honduras

lag from integral overlap between 20 and 20 days, maximum overlap on day 20

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




India


no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

no meaningful fits to the function I available


no meaningful fits to the function I available (averaged data)






Indonesia

no clear lag in the first six months of the pandemic from integral overlap
no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days
no meaningful fits to the function I available
no meaningful fits to the function I available (averaged data)



Iran

no clear lag in the first six months of the pandemic from integral overlap
no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days
no meaningful fits to the function I available
no meaningful fits to the function I available (averaged data)



Iraq


no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days


no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)


new cases, Iraq




new deaths, Iraq




no clear lag in the first six months of the pandemic




averaged new cases, Iraq



averaged new deaths, Iraq



no clear lag in the first six months of the pandemic



Japan


lag from integral overlap between 11 and 16 days, maximum overlap on day 14

lag from integral overlap, using data averaged over 7 days, between 12 and 17 days, maximum overlap on day 14


no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)


new cases, Japan




new deaths, Japan




lag 14 in [11,16]




averaged new cases, Japan



averaged new deaths, Japan



lag 14 in [12,17]




Mexico

no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




Moldova

lag from integral overlap between 20 and 20 days, maximum overlap on day 20

lag from integral overlap, using data averaged over 7 days, between 21 and 26 days, maximum overlap on day 23

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




Nigeria

lag from integral overlap between 21 and 21 days, maximum overlap on day 21


no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

no meaningful fits to the function I available


no meaningful fits to the function I available (averaged data)




averaged new cases, Nigeria



averaged new deaths, Nigeria



no clear lag in the first six months of the pandemic



Pakistan


lag from integral overlap between 10 and 21 days, maximum overlap on day 10

lag from integral overlap, using data averaged over 7 days, between 9 and 16 days, maximum overlap on day 11


no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)


new cases, Pakistan




new deaths, Pakistan




lag 10 in [10,21]




averaged new cases, Pakistan



averaged new deaths, Pakistan



lag 11 in [9,16]



Panama


no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days


no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)


new cases, Panama




new deaths, Panama




no clear lag in the first six months of the pandemic




averaged new cases, Panama



averaged new deaths, Panama




no clear lag in the first six months of the pandemic




Peru

no clear lag in the first six months of the pandemic from integral overlap
 no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days
 no meaningful fits to the function I available
 no meaningful fits to the function I available (averaged data)




Philippines

no clear lag in the first six months of the pandemic from integral overlap
 no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days
 no meaningful fits to the function I available
 no meaningful fits to the function I available (averaged data)



Russia

no clear lag in the first six months of the pandemic from integral overlap
 no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days
 no meaningful fits to the function I available
 no meaningful fits to the function I available (averaged data)




Saudi Arabia

no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




South Africa


no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)






Sudan

lag from integral overlap between 0 and 0 days, maximum overlap on day 0

lag from integral overlap, using data averaged over 7 days, between 7 and 11 days, maximum overlap on day 8

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




Switzerland

lag from integral overlap between 11 and 11 days, maximum overlap on day 11

lag from integral overlap, using data averaged over 7 days, between 9 and 12 days, maximum overlap on day 10

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




Turkey

lag from integral overlap between 2 and 6 days, maximum overlap on day 5

lag from integral overlap, using data averaged over 7 days, between 2 and 7 days, maximum overlap on day 4

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




Ukraine

no clear lag in the first six months of the pandemic from integral overlap

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




Hubei, China

lag from integral overlap between 0 and 0 days, maximum overlap on day 0

lag from integral overlap, using data averaged over 7 days, between 3 and 6 days, maximum overlap on day 5

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




Ontario, Canada

lag from integral overlap between 0 and 0 days, maximum overlap on day 0

lag from integral overlap, using data averaged over 7 days, between 3 and 11 days, maximum overlap on day 7

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)




Quebec, Canada


lag from integral overlap between 11 and 14 days, maximum overlap on day 11

lag from integral overlap, using data averaged over 7 days, between 7 and 14 days, maximum overlap on day 11

no meaningful fits to the function I available

no meaningful fits to the function I available (averaged data)






Plots, fits and integral averages for US counties (ordered as on the JHU server)

Maricopa, Arizona

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days


lag from fits to the function I (averaged data): -1.31252 days



Parameters of fits to averaged cases data.

The best fit ends on Wed 15 Apr 2020.


α	β	k	τ	σ			
0.485 ± 0.0115	0.157 ± 0.00312	2530 ± 26.3	33.6 ± 0.112	21.3 ± 0.2			
badness	α	β	k	τ	σ	quality	
0.35%	error	2.4%	2%	1%	0.33%	0.94%	7.02%



Parameters of fits to averaged death data.

The best fit ends on Thu 7 May 2020.


α	β	k	τ	σ			
0.586 ± 0.123	0.0991 ± 0.00472	$230 \pm 12.$	35.3 ± 1.14	9.94 ± 0.587			
badness	α	β	k	τ	σ	quality	
2.8%	error	21%	4.8%	5.2%	3.2%	5.9%	42.8%



Los Angeles, California

lag from integral overlap, using data averaged over 7 days, between -3 and 28 days, maximum overlap on day 4


lag from fits to the function I (averaged data): 21.5503 days



Parameters of fits to averaged cases data.

The best fit ends on Wed 20 May 2020.


α	β	k	τ	σ
0.396 ± 0.0349	0.0744 ± 0.0012	$50000. \pm 772.$	63.1 ± 0.465	27.5 ± 0.379
badness	α	β	k	τ
1.7%	error	8.8%	1.6%	1.5%



Parameters of fits to averaged death data.

The best fit ends on Wed 8 Jul 2020.


α	β	k	τ	σ
0.175 ± 0.00184	0.036 ± 0.000298	$4300. \pm 19.3$	$71. \pm 0.248$	30.9 ± 0.136
badness	α	β	k	τ
0.55%	error	1.%	0.83%	0.45%



Riverside, California

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days


lag from fits to the function I (averaged data): 11.8012 days



Parameters of fits to averaged cases data.

The best fit ends on Wed 6 May 2020.


α	β	k	τ	σ
0.322 ± 0.0132	0.104 ± 0.00271	$5220. \pm 62.5$	42.8 ± 0.191	23.2 ± 0.444
<hr/>				
badness	α	β	k	τ
0.96%	error	4.1%	2.6%	1.2%
0.45%				
1.9%				
11.2%				



Parameters of fits to averaged death data.

The best fit ends on Thu 2 Jul 2020.


α	β	k	τ	σ
0.132 ± 0.00275	0.0296 ± 0.00318	$603. \pm 14.7$	35.3 ± 0.568	70.5 ± 1.5
<hr/>				
badness	α	β	k	τ
0.96%	error	2.1%	11.%	2.4%
1.6%				
2.1%				
10.9%				



Fairfield, Connecticut

lag from integral overlap, using data averaged over 7 days, between 4 and 12 days, maximum overlap on day 9


lag from fits to the function I (averaged data): 6.90919 days



Parameters of fits to averaged cases data.

The best fit ends on Tue 21 Jul 2020.


α	β	k	τ	σ
0.202 ± 0.00548	0.0536 ± 0.00126	$17000. \pm 35.1$	38.2 ± 0.537	26.6 ± 0.447
badness	α	β	k	τ
2.5%	error	2.7%	2.3%	0.21%



Parameters of fits to averaged death data.

The best fit ends on Mon 20 Jul 2020.


α	β	k	τ	σ
0.219 ± 0.00177	0.0609 ± 0.000403	$1400. \pm 0.759$	37.6 ± 0.141	26.3 ± 0.128
badness	α	β	k	τ
0.46%	error	0.81%	0.66%	0.054%



Hartford, Connecticut

lag from integral overlap, using data averaged over 7 days, between 0 and 5 days, maximum overlap on day 3


lag from fits to the function I (averaged data): 4.14423 days



Parameters of fits to averaged cases data.

The best fit ends on Wed 22 Jul 2020.


α	β	k	τ	σ			
0.25 ± 0.0107	0.0605 ± 0.000763	12000 ± 23.6	50 ± 0.152	24.6 ± 0.348			
badness	error	α	β	k	τ	σ	quality
1.7%		4.3%	1.3%	0.2%	0.3%	1.4%	9.13%



Parameters of fits to averaged death data.

The best fit ends on Fri 17 Jul 2020.


α	β	k	τ	σ			
0.274 ± 0.00561	0.0652 ± 0.000705	1400 ± 1.77	38.4 ± 0.149	23.6 ± 0.178			
badness	error	α	β	k	τ	σ	quality
1.1%		2%	1.1%	0.13%	0.39%	0.75%	5.51%



New Haven, Connecticut

lag from integral overlap, using data averaged over 7 days, between 4 and 9 days, maximum overlap on day 6

lag from fits to the function I (averaged data): 5.79625 days




Parameters of fits to averaged cases data.

The best fit ends on Sat 11 Jul 2020.

α	β	k	τ	σ
0.247 ± 0.00281	0.0593 ± 0.000413	12600 ± 10.8	42.8 ± 0.103	27.7 ± 0.117
badness	α	β	k	τ
0.63%	error	1.1%	0.7%	0.086%

σ	quality
0.42%	3.21%




Parameters of fits to averaged death data.

The best fit ends on Fri 24 Jul 2020.

α	β	k	τ	σ
0.269 ± 0.00375	0.0683 ± 0.000323	1090 ± 0.617	39.2 ± 0.0612	19 ± 0.113
badness	α	β	k	τ
0.55%	error	1.4%	0.47%	0.056%


σ	quality
0.59%	3.22%



District of Columbia, District of Columbia

lag from integral overlap, using data averaged over 7 days, between 1 and 8 days, maximum overlap on day 4


lag from fits to the function I (averaged data): -4.96504 days



Parameters of fits to averaged cases data.

The best fit ends on Mon 29 Jun 2020.

α	β	k	τ	σ
0.25 ± 0.00712	0.0622 ± 0.000359	10500 ± 17.1	53.8 ± 0.0657	19.3 ± 0.22
<hr/>				
badness	error	α	β	k
0.88%	2.8%	0.58%	0.16%	0.12%
				σ
				1.1%
				5.73%




Parameters of fits to averaged death data.


The best fit ends on Wed 19 Aug 2020.

α	β	k	τ	σ
0.146 ± 0.00156	0.0366 ± 0.00187	604 ± 0.981	33.3 ± 0.28	36.5 ± 0.593
<hr/>				
badness	error	α	β	k
0.78%	1.1%	5.1%	0.16%	0.84%
				σ
				1.6%
				5.47%

District of Columbia, District of Columbia



District of Columbia, District of Columbia



Miami-Dade, Florida




no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days

lag from fits to the function I (averaged data): 12.7839 days

new cases, Miami-Dade, Florida

new deaths, Miami-Dade, Florida

no clear lag in the first six months of the pandemic





Parameters of fits to averaged cases data.

The best fit ends on Tue 5 May 2020.

α	β	k	τ	σ
0.344 ± 0.00441	0.0861 ± 0.00551	$15000. \pm 91.4$	20.1 ± 0.126	33.9 ± 0.109
badness	α	β	k	τ
0.44%	error	1.3%	6.4%	0.61%
σ	quality			
0.32%	4.59%			

Miami-Dade, Florida


Miami-Dade, Florida



Parameters of fits to averaged death data.

The best fit ends on Tue 23 Jun 2020.


α	β	k	τ	σ
0.231 ± 0.00848	0.0438 ± 0.00113	$1030. \pm 13.4$	48.5 ± 0.594	17.1 ± 0.325
badness	α	β	k	τ
1.5%	error	3.7%	2.6%	1.3%
σ	quality			
1.9%	12.2%			



Palm Beach, Florida

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days


lag from fits to the function I (averaged data): 2.79552 days



Parameters of fits to averaged cases data.

The best fit ends on Mon 4 May 2020.


α	β	k	τ	σ
0.372 ± 0.0117	0.0865 ± 0.00214	$3850 \pm 55.$	35.9 ± 0.328	17.1 ± 0.216
badness	α	β	k	τ
0.71%	error	3.1%	2.5%	1.4%
				0.91%
				1.3%
				9.95%



Parameters of fits to averaged death data.

The best fit ends on Thu 28 May 2020.


α	β	k	τ	σ
0.352 ± 0.00711	0.053 ± 0.000551	440 ± 4.13	46.9 ± 0.382	13.2 ± 0.0959
badness	α	β	k	τ
0.6%	error	2.%	1.%	0.94%
				0.82%
				0.72%
				6.14%



Cook, Illinois

lag from integral overlap, using data averaged over 7 days, between 2 and 10 days, maximum overlap on day 6


lag from fits to the function I (averaged data): 24.7331 days



Parameters of fits to averaged cases data.

The best fit ends on Tue 21 Apr 2020.


α	β	k	τ	σ
0.364 ± 0.00482	0.118 ± 0.00122	$32800 \pm 311.$	77.9 ± 0.156	57.4 ± 0.139
badness	α	β	k	τ
0.22%	error	1.3%	1.%	0.95%
			0.2%	0.24%
				3.97%



Parameters of fits to averaged death data.

The best fit ends on Mon 8 Jun 2020.


α	β	k	τ	σ
0.275 ± 0.00601	0.0674 ± 0.000436	4460 ± 14.7	52.6 ± 0.107	19.8 ± 0.156
badness	α	β	k	τ
0.54%	error	2.2%	0.65%	0.33%
			0.2%	0.79%
				4.69%



Marion, Indiana

lag from integral overlap, using data averaged over 7 days, between 0 and 12 days, maximum overlap on day 9

lag from fits to the function I (averaged data): 16.7342 days



Parameters of fits to averaged cases data.


The best fit ends on Mon 29 Jun 2020.

α	β	k	τ	σ
0.347 ± 0.0103	0.0587 ± 0.000287	11500 ± 17.7	53.8 ± 0.0722	21.5 ± 0.141
badness	error	α	β	k
1.1%		3.%	0.49%	0.15%

badness	error	α	β	k	τ	σ	quality
1.1%		3.%	0.49%	0.15%	0.13%	0.66%	5.45%

Marion, Indiana

Marion, Indiana




Parameters of fits to averaged death data.

The best fit ends on Fri 14 Aug 2020.

α	β	k	τ	σ
0.198 ± 0.00795	0.0488 ± 0.000938	724 ± 1.69	46.6 ± 0.347	24.3 ± 0.479
badness	error	α	β	k
2.6%		4.%	1.9%	0.23%


2.6% error 4.% 1.9% 0.23% 0.74% 2.% 11.5%



Orleans, Louisiana

lag from integral overlap, using data averaged over 7 days, between 2 and 11 days, maximum overlap on day 7

lag from fits to the function I (averaged data): 6.7595 days




Parameters of fits to averaged cases data.

The best fit ends on Fri 27 Nov 2020.

α	β	k	τ	σ
0.309 ± 0.0238	0.00834 ± 0.000299	$21300 \pm 758.$	163 ± 9.07	23.6 ± 0.304

badness	α	β	k	τ	σ	quality
4.5%	error	7.7%	3.6%	3.6%	5.6%	1.3% 26.3%




Parameters of fits to averaged death data.

The best fit ends on Wed 10 Jun 2020.

α	β	k	τ	σ
0.349 ± 0.011	0.0912 ± 0.000797	513 ± 0.627	30.3 ± 0.0763	12.3 ± 0.188


badness	α	β	k	τ	σ	quality
1.1%	error	3.1%	0.87%	0.12%	0.25%	1.5% 7.01%



Baltimore, Maryland

lag from integral overlap, using data averaged over 7 days, between 1 and 19 days, maximum overlap on day 14


lag from fits to the function I (averaged data): 29.6195 days



Parameters of fits to averaged cases data.

The best fit ends on Sat 4 Jul 2020.


α	β	k	τ	σ			
0.245 ± 0.00963	0.05 ± 0.000473	$8690. \pm 34.4$	64.1 ± 0.184	24.2 ± 0.286			
badness	α	β	k	τ	σ	quality	
1.3%	error	3.9%	0.95%	0.4%	0.29%	1.2%	8.06%



Parameters of fits to averaged death data.

The best fit ends on Wed 28 Oct 2020.


α	β	k	τ	σ			
0.0993 ± 0.00145	0.0159 ± 0.00184	$705. \pm 4.11$	38.5 ± 0.343	47.7 ± 0.692			
badness	α	β	k	τ	σ	quality	
1.5%	error	1.5%	12%	0.58%	0.89%	1.5%	7.26%



Montgomery, Maryland

lag from integral overlap, using data averaged over 7 days, between -7 and 0 days, maximum overlap on day -4


lag from fits to the function I (averaged data): -15.057 days



Parameters of fits to averaged cases data.

The best fit ends on Thu 18 Jun 2020.


α	β	k	τ	σ
0.217 ± 0.00739	0.0592 ± 0.00049	16000 ± 70.4	72.7 ± 0.162	29.6 ± 0.319
badness	α	β	k	σ
0.73%	error	3.4%	0.83%	0.44%



Parameters of fits to averaged death data.

The best fit ends on Wed 22 Jul 2020.


α	β	k	τ	σ
0.19 ± 0.00384	0.0558 ± 0.000825	779 ± 1.39	39.8 ± 0.268	21.6 ± 0.373
badness	α	β	k	σ
0.98%	error	2%	1.5%	0.18%



Prince George's, Maryland

lag from integral overlap, using data averaged over 7 days, between -4 and 4 days, maximum overlap on day -1


lag from fits to the function I (averaged data): -10.4462 days



Parameters of fits to averaged cases data.

The best fit ends on Thu 2 Jul 2020.


α	β	k	τ	σ
0.23 \pm 0.00836	0.0658 \pm 0.000471	19100. \pm 36.5	63.7 \pm 0.0725	27.7 \pm 0.327
<hr/>				
badness	α	β	k	σ
0.9%	error	3.6%	0.72%	0.19%
0.11%			1.2%	6.74%



Parameters of fits to averaged death data.

The best fit ends on Sat 1 Aug 2020.


α	β	k	τ	σ
0.167 \pm 0.00339	0.047 \pm 0.000759	745. \pm 1.59	48.8 \pm 0.314	29. \pm 0.411
<hr/>				
badness	α	β	k	σ
1.%	error	2.%	1.6%	0.21%
0.64%			1.4%	6.95%



Bristol, Massachusetts

lag from integral overlap, using data averaged over 7 days, between 9 and 19 days, maximum overlap on day 14


lag from fits to the function I (averaged data): -0.459238 days



Parameters of fits to averaged cases data.

The best fit ends on Mon 29 Jun 2020.


α	β	k	τ	σ
0.24 ± 0.0052	0.0722 ± 0.000278	$8040. \pm 6.09$	50.2 ± 0.0372	17.2 ± 0.19
badness	error	α	β	k
0.51%		2.2%	0.39%	0.076%



Parameters of fits to averaged death data.

The best fit ends on Fri 21 Aug 2020.


α	β	k	τ	σ
0.188 ± 0.00494	0.0477 ± 0.000444	$643. \pm 0.918$	50.1 ± 0.143	19.6 ± 0.303
badness	error	α	β	k
1.1%		2.6%	0.93%	0.14%



Essex, Massachusetts

lag from integral overlap, using data averaged over 7 days, between 9 and 15 days, maximum overlap on day 11


lag from fits to the function I (averaged data): 11.5304 days



Parameters of fits to averaged cases data.

The best fit ends on Fri 24 Apr 2020.


α	β	k	τ	σ			
0.442 ± 0.0094	0.119 ± 0.000556	$10800. \pm 65.5$	40.8 ± 0.105	13.3 ± 0.0945			
badness	error	α	β	k	τ	σ	quality
0.25%		2.1%	0.47%	0.61%	0.26%	0.71%	4.42%



Parameters of fits to averaged death data.

The best fit ends on Mon 27 Jul 2020.


α	β	k	τ	σ			
0.171 ± 0.00329	0.0491 ± 0.00407	$1170. \pm 2.57$	26.1 ± 0.471	37.7 ± 0.541			
badness	error	α	β	k	τ	σ	quality
1.3%		1.9%	8.3%	0.22%	1.8%	1.4%	8.75%



Hampden, Massachusetts

lag from integral overlap, using data averaged over 7 days, between 6 and 12 days, maximum overlap on day 10


lag from fits to the function I (averaged data): 0.825176 days



Parameters of fits to averaged cases data.

The best fit ends on Wed 6 May 2020.


α	β	k	τ	σ
0.355 ± 0.00445	0.0825 ± 0.000553	5610 ± 26.2	38.7 ± 0.119	14.6 ± 0.0771
badness	α	β	k	σ
0.35%	error	1.3%	0.67%	0.47%



Parameters of fits to averaged death data.

The best fit ends on Mon 3 Aug 2020.


α	β	k	τ	σ
0.211 ± 0.00464	0.0461 ± 0.00091	696 ± 1.51	30.9 ± 0.418	20.7 ± 0.274
badness	α	β	k	σ
1.6%	error	2.2%	2%	0.22%



Middlesex, Massachusetts

lag from integral overlap, using data averaged over 7 days, between 9 and 13 days, maximum overlap on day 11


lag from fits to the function I (averaged data): 10.2116 days



Parameters of fits to averaged cases data.

The best fit ends on Thu 9 Jul 2020.


α	β	k	τ	σ
0.165 ± 0.00157	0.0512 ± 0.00173	23000 ± 29.1	39.8 ± 0.349	41.3 ± 0.637
badness	α	β	k	τ
0.55%	error	0.95%	3.4%	0.13%



Parameters of fits to averaged death data.

The best fit ends on Mon 3 Aug 2020.


α	β	k	τ	σ
0.18 ± 0.00212	0.0378 ± 0.00261	2010 ± 4.62	32.1 ± 0.199	32.1 ± 0.551
badness	α	β	k	τ
0.99%	error	1.2%	6.9%	0.23%



Norfolk, Massachusetts

lag from integral overlap, using data averaged over 7 days, between 9 and 14 days, maximum overlap on day 12


lag from fits to the function I (averaged data): 9.99908 days



Parameters of fits to averaged cases data.

The best fit ends on Tue 30 Jun 2020.


α	β	k	τ	σ
0.176 ± 0.00138	0.0563 ± 0.00151	8460 ± 8.22	35.2 ± 0.29	36.9 ± 0.504
badness	α	β	k	τ
0.43%	error	0.78%	2.7%	0.097%



Parameters of fits to averaged death data.

The best fit ends on Thu 23 Jul 2020.


α	β	k	τ	σ
0.207 ± 0.00227	0.0416 ± 0.00288	987 ± 2.01	27.7 ± 0.148	29.9 ± 0.37
badness	α	β	k	τ
1.1%	error	1.1%	6.9%	0.2%



Plymouth, Massachusetts

lag from integral overlap, using data averaged over 7 days, between 8 and 12 days, maximum overlap on day 10


lag from fits to the function I (averaged data): 8.7179 days



Parameters of fits to averaged cases data.

The best fit ends on Tue 30 Jun 2020.


α	β	k	τ	σ
0.207 ± 0.00429	0.0735 ± 0.000759	8320 ± 7.56	41.5 ± 0.21	21.4 ± 0.394
badness	α	β	k	τ
0.85%	error	2.1%	1.9%	0.091%



Parameters of fits to averaged death data.

The best fit ends on Sun 23 Aug 2020.


α	β	k	τ	σ
0.178 ± 0.00356	0.0382 ± 0.000795	732 ± 1.72	33 ± 0.526	23.5 ± 0.303
badness	α	β	k	τ
1.5%	error	2.0%	2.1%	0.23%



Suffolk, Massachusetts

lag from integral overlap, using data averaged over 7 days, between 8 and 13 days, maximum overlap on day 10


lag from fits to the function I (averaged data): 15.7431 days



Parameters of fits to averaged cases data.

The best fit ends on Mon 13 Apr 2020.


α	β	k	τ	σ			
0.418 ± 0.0109	0.15 ± 0.00118	$9940. \pm 107.$	$40. \pm 0.139$	15.5 ± 0.157			
badness	α	β	k	τ	σ	quality	
0.14%	error	2.6%	0.79%	1.1%	0.35%	1.9%	5.97%



Parameters of fits to averaged death data.

The best fit ends on Fri 7 Aug 2020.


α	β	k	τ	σ			
0.189 ± 0.00214	0.0323 ± 0.00266	$1090. \pm 2.96$	31.5 ± 0.15	$24. \pm 0.696$			
badness	α	β	k	τ	σ	quality	
1.1%	error	1.1%	8.2%	0.27%	0.48%	2.9%	6.91%



Worcester, Massachusetts

lag from integral overlap, using data averaged over 7 days, between 9 and 16 days, maximum overlap on day 13


lag from fits to the function I (averaged data): 7.79107 days



Parameters of fits to averaged cases data.

The best fit ends on Thu 28 May 2020.


α	β	k	τ	σ
0.225 ± 0.00685	0.0735 ± 0.000801	12400 ± 57.8	55.2 ± 0.113	22 ± 0.353
<hr/>				
badness	α	β	k	τ
0.53%	error	3.1%	1.1%	0.46%
				0.2%
				1.6%
				6.95%



Parameters of fits to averaged death data.

The best fit ends on Fri 7 Aug 2020.


α	β	k	τ	σ
0.14 ± 0.00216	0.0439 ± 0.00231	999 ± 2.36	37.7 ± 0.714	37.8 ± 1.38
<hr/>				
badness	α	β	k	τ
0.94%	error	1.5%	5.3%	0.24%
				1.9%
				3.7%
				9.44%



Macomb, Michigan

lag from integral overlap, using data averaged over 7 days, between 6 and 9 days, maximum overlap on day 8


lag from fits to the function I (averaged data): 9.2049 days



Parameters of fits to averaged cases data.

The best fit ends on Sat 30 May 2020.


α	β	k	τ	σ
0.295 ± 0.00539	0.0726 ± 0.0013	6890 ± 19.6	29.8 ± 0.174	19.8 ± 0.192
badness	α	β	k	τ
0.76%	error	1.8%	1.8%	0.28%



Parameters of fits to averaged death data.

The best fit ends on Sun 9 Aug 2020.


α	β	k	τ	σ
0.217 ± 0.00243	0.0458 ± 0.00316	942 ± 0.937	21.1 ± 0.141	23.6 ± 0.355
badness	α	β	k	τ
0.85%	error	1.1%	6.9%	0.1%



Oakland, Michigan

lag from integral overlap, using data averaged over 7 days, between 3 and 9 days, maximum overlap on day 6


lag from fits to the function I (averaged data): 3.53659 days



Parameters of fits to averaged cases data.

The best fit ends on Thu 14 May 2020.


α	β	k	τ	σ			
0.253 ± 0.0032	0.0508 ± 0.00348	$10500. \pm 127.$	23.3 ± 0.168	29.3 ± 0.24			
badness	α	β	k	τ	σ	quality	
0.53%	error	1.3%	6.9%	1.2%	0.72%	0.82%	5.38%



Parameters of fits to averaged death data.

The best fit ends on Wed 3 Jun 2020.


α	β	k	τ	σ			
0.329 ± 0.00601	0.0795 ± 0.000571	$1060. \pm 1.77$	32.2 ± 0.0504	13.1 ± 0.118			
badness	α	β	k	τ	σ	quality	
0.61%	error	1.8%	0.72%	0.17%	0.16%	0.9%	4.39%



Wayne, Michigan

lag from integral overlap, using data averaged over 7 days, between 7 and 10 days, maximum overlap on day 9

lag from fits to the function I (averaged data): 15.0953 days




Parameters of fits to averaged cases data.

The best fit ends on Wed 24 Jun 2020.

α	β	k	τ	σ
0.296 ± 0.00392	0.05 ± 0.000601	22600 ± 43.3	30 ± 0.156	21.8 ± 0.0933
badness	α	β	k	τ
0.94%	error	1.3%	1.2%	0.19%

σ	quality
0.43%	4.6%




Parameters of fits to averaged death data.

The best fit ends on Thu 21 May 2020.

α	β	k	τ	σ
0.368 ± 0.0147	0.103 ± 0.00134	2470 ± 8.3	36.2 ± 0.0723	17.5 ± 0.244
badness	α	β	k	τ
0.95%	error	4.%	1.3%	0.34%


σ	quality
1.4%	8.18%



Hennepin, Minnesota

lag from integral overlap, using data averaged over 7 days, between -13 and -8 days, maximum overlap on day -10


lag from fits to the function I (averaged data): -11.2948 days



Parameters of fits to averaged cases data.

The best fit ends on Fri 16 Oct 2020.


α	β	k	τ	σ			
0.132 ± 0.00585	0.0168 ± 0.00033	$43100. \pm 882.$	$167. \pm 2.61$	56.8 ± 0.521			
badness	error	α	β	k	τ	σ	quality
2.1%		4.4%	2.9%	2.9%	1.6%	0.92%	13.%



Parameters of fits to averaged death data.

The best fit ends on Fri 7 Aug 2020.


α	β	k	τ	σ			
0.21 ± 0.00611	0.0554 ± 0.000794	$822. \pm 1.46$	46.6 ± 0.232	25.5 ± 0.352			
badness	error	α	β	k	τ	σ	quality
1.5%		2.9%	1.4%	0.18%	0.5%	1.4%	7.85%



St. Louis, Missouri

lag from integral overlap, using data averaged over 7 days, between 11 and 16 days, maximum overlap on day 14

lag from fits to the function I (averaged data): 23.5652 days




Parameters of fits to averaged cases data.

The best fit ends on Wed 27 May 2020.

α	β	k	τ	σ
0.32 ± 0.00768	0.0587 ± 0.000829	5440 ± 36.4	44 ± 0.232	20.2 ± 0.145

badness	α	β	k	τ	σ	quality
0.8%	error	2.4%	1.4%	0.67%	0.53%	0.72%




Parameters of fits to averaged death data.

The best fit ends on Tue 24 Nov 2020.

α	β	k	τ	σ
0.13 ± 0.00251	0.01 ± 0.00348	1150 ± 14.1	34.2 ± 0.216	105 ± 2.46


badness	α	β	k	τ	σ	quality
2.2%	error	1.9%	35.%	1.2%	0.63%	2.4%



Clark, Nevada

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days


lag from fits to the function I (averaged data): 7.83028 days



Parameters of fits to averaged cases data.

The best fit ends on Sun 17 May 2020.


α	β	k	τ	σ			
0.307 ± 0.0072	0.0573 ± 0.00127	6470 ± 87.2	46.8 ± 0.476	23.3 ± 0.171			
badness	α	β	k	τ	σ	quality	
0.82%	error	2.3%	2.2%	1.3%	1.%	0.73%	8.49%



Parameters of fits to averaged death data.

The best fit ends on Mon 15 Jun 2020.


α	β	k	τ	σ			
0.264 ± 0.00735	0.0589 ± 0.000678	400 ± 1.5	43.7 ± 0.127	16.9 ± 0.213			
badness	α	β	k	τ	σ	quality	
1.2%	error	2.8%	1.2%	0.38%	0.29%	1.3%	7.05%



Bergen, New Jersey

lag from integral overlap, using data averaged over 7 days, between 7 and 11 days, maximum overlap on day 9


lag from fits to the function I (averaged data): 5.729 days



Parameters of fits to averaged cases data.

The best fit ends on Tue 26 May 2020.


α	β	k	τ	σ
0.333 ± 0.00831	0.0916 ± 0.00157	17900 ± 40.2	37.2 ± 0.167	26.2 ± 0.231
badness	α	β	k	τ
0.94%	error	2.5%	1.7%	0.22%
			0.45%	0.88%
			quality	6.7%



Parameters of fits to averaged death data.

The best fit ends on Tue 9 Jun 2020.


α	β	k	τ	σ
0.346 ± 0.0122	0.0806 ± 0.00125	1640 ± 4.72	38.6 ± 0.118	22.7 ± 0.222
badness	α	β	k	τ
1.5%	error	3.5%	1.6%	0.29%
			0.31%	0.98%
			quality	8.11%



Camden, New Jersey

lag from integral overlap, using data averaged over 7 days, between 8 and 16 days, maximum overlap on day 11


lag from fits to the function I (averaged data): 11.0318 days



Parameters of fits to averaged cases data.

The best fit ends on Mon 29 Jun 2020.


α	β	k	τ	σ			
0.225 ± 0.00482	0.0649 ± 0.000552	7390 ± 11.3	50.6 ± 0.0883	25.5 ± 0.239			
badness	α	β	k	τ	σ	quality	
0.92%	error	2.1%	0.85%	0.15%	0.17%	0.94%	5.18%



Parameters of fits to averaged death data.

The best fit ends on Fri 5 Jun 2020.


α	β	k	τ	σ			
0.331 ± 0.0218	0.0795 ± 0.00132	379 ± 2.8	39.8 ± 0.207	12.2 ± 0.377			
badness	α	β	k	τ	σ	quality	
1.3%	error	6.6%	1.7%	0.74%	0.52%	3.1%	13.9%



Essex, New Jersey

lag from integral overlap, using data averaged over 7 days, between 5 and 9 days, maximum overlap on day 7


lag from fits to the function I (averaged data): 8.98589 days



Parameters of fits to averaged cases data.

The best fit ends on Mon 18 May 2020.


α	β	k	τ	σ			
0.321 ± 0.00359	0.0884 ± 0.00072	17100 ± 31.6	35 ± 0.0462	20.8 ± 0.108			
badness	α	β	k	τ	σ	quality	
0.3%	error	1.1%	0.82%	0.18%	0.13%	0.52%	3.07%



Parameters of fits to averaged death data.

The best fit ends on Fri 29 May 2020.


α	β	k	τ	σ			
0.366 ± 0.0108	0.0962 ± 0.00107	1660 ± 3.95	33.9 ± 0.0668	17 ± 0.182			
badness	α	β	k	τ	σ	quality	
0.81%	error	3.%	1.1%	0.24%	0.2%	1.1%	6.38%



Hudson, New Jersey

lag from integral overlap, using data averaged over 7 days, between 7 and 11 days, maximum overlap on day 9


lag from fits to the function I (averaged data): 8.35023 days



Parameters of fits to averaged cases data.

The best fit ends on Wed 27 May 2020.


α	β	k	τ	σ			
0.35 ± 0.00958	0.0884 ± 0.00106	$19600. \pm 51.1$	$39. \pm 0.0752$	22.4 ± 0.179			
badness	error	α	β	k	τ	σ	quality
0.88%		2.7%	1.2%	0.26%	0.19%	0.8%	6.07%



Parameters of fits to averaged death data.

The best fit ends on Thu 26 Nov 2020.


α	β	k	τ	σ			
0.261 ± 0.011	0.0365 ± 0.000674	$1530. \pm 2.41$	34.4 ± 0.512	21.6 ± 0.27			
badness	error	α	β	k	τ	σ	quality
3.6%		4.2%	1.8%	0.16%	1.5%	1.3%	12.6%



Mercer, New Jersey

lag from integral overlap, using data averaged over 7 days, between 4 and 11 days, maximum overlap on day 8

lag from fits to the function I (averaged data): 10.9576 days




Parameters of fits to averaged cases data.

The best fit ends on Mon 27 Jul 2020.

α	β	k	τ	σ
0.191 ± 0.0054	0.0572 ± 0.000924	$7880. \pm 12.4$	46.7 ± 0.331	26.9 ± 0.473
badness	error	α	β	k
1.4%		2.8%	1.6%	0.16%


badness	error	α	β	k	τ	σ	quality
1.4%		2.8%	1.6%	0.16%	0.71%	1.8%	8.51%



Parameters of fits to averaged death data.

The best fit ends on Wed 24 Jun 2020.


α	β	k	τ	σ
0.278 ± 0.0142	0.0775 ± 0.00114	$536. \pm 1.52$	34.3 ± 0.123	10.5 ± 0.387
badness	error	α	β	k
1.4%		5.1%	1.5%	0.28%



Middlesex, New Jersey

lag from integral overlap, using data averaged over 7 days, between 6 and 11 days, maximum overlap on day 9


lag from fits to the function I (averaged data): 2.45389 days



Parameters of fits to averaged cases data.

The best fit ends on Sun 31 May 2020.


α	β	k	τ	σ			
0.334 ± 0.0129	0.0859 ± 0.0012	16000 ± 49.1	40.6 ± 0.0942	21.6 ± 0.253			
badness	error	α	β	k	τ	σ	quality
0.92%		3.9%	1.4%	0.31%	0.23%	1.2%	7.88%



Parameters of fits to averaged death data.

The best fit ends on Mon 27 Apr 2020.


α	β	k	τ	σ			
0.59 ± 0.0252	0.141 ± 0.00277	534 ± 7.47	21.8 ± 0.211	6.74 ± 0.152			
badness	error	α	β	k	τ	σ	quality
0.53%		4.3%	2%	1.4%	0.97%	2.3%	11.4%



Monmouth, New Jersey

lag from integral overlap, using data averaged over 7 days, between 9 and 12 days, maximum overlap on day 10


lag from fits to the function I (averaged data): 10.405 days



Parameters of fits to averaged cases data.

The best fit ends on Sat 4 Jul 2020.


α	β	k	τ	σ
0.25 ± 0.00399	0.0421 ± 0.000532	$9520. \pm 25.5$	40.6 ± 0.146	25.1 ± 0.128
badness	α	β	k	τ
1.3%	error	1.6%	1.3%	0.27%



Parameters of fits to averaged death data.

The best fit ends on Wed 24 Jun 2020.


α	β	k	τ	σ
0.216 ± 0.00687	0.0477 ± 0.00116	$762. \pm 5.71$	47.1 ± 0.255	22.9 ± 0.361
badness	α	β	k	τ
1.4%	error	3.2%	2.4%	0.75%



Morris, New Jersey

lag from integral overlap, using data averaged over 7 days, between 6 and 10 days, maximum overlap on day 8


lag from fits to the function I (averaged data): 4.2849 days



Parameters of fits to averaged cases data.

The best fit ends on Wed 27 May 2020.


α	β	k	τ	σ
0.313 ± 0.00587	0.0834 ± 0.00126	$6370. \pm 14.4$	31.2 ± 0.141	20.2 ± 0.19
badness	α	β	k	τ
0.7%	error	1.9%	1.5%	0.23%



Parameters of fits to averaged death data.

The best fit ends on Sun 14 Jun 2020.


α	β	k	τ	σ
0.37 ± 0.0107	0.0846 ± 0.000685	$638. \pm 0.98$	30.1 ± 0.0623	11.4 ± 0.149
badness	α	β	k	τ
0.98%	error	2.9%	0.81%	0.15%



Ocean, New Jersey

lag from integral overlap, using data averaged over 7 days, between 10 and 19 days, maximum overlap on day 14


lag from fits to the function I (averaged data): 0.709819 days



Parameters of fits to averaged cases data.

The best fit ends on Mon 25 May 2020.


α	β	k	τ	σ
0.345 ± 0.00802	0.074 ± 0.000925	$8820. \pm 32.7$	34.8 ± 0.0932	17.1 ± 0.144
badness	α	β	k	τ
0.71%	error	2.3%	1.2%	0.37%



Parameters of fits to averaged death data.

The best fit ends on Sun 17 May 2020.


α	β	k	τ	σ
0.551 ± 0.0169	0.0866 ± 0.000512	$750. \pm 3.77$	36.2 ± 0.133	8.93 ± 0.0891
badness	α	β	k	τ
0.57%	error	3.1%	0.59%	0.5%



Passaic, New Jersey

lag from integral overlap, using data averaged over 7 days, between 7 and 12 days, maximum overlap on day 8


lag from fits to the function I (averaged data): 41.9186 days



Parameters of fits to averaged cases data.

The best fit ends on Wed 3 Jun 2020.


α	β	k	τ	σ
0.449 ± 0.0286	0.0978 ± 0.00113	16200 ± 40.9	38.3 ± 0.0843	18.6 ± 0.259
badness	α	β	k	τ
1.3%	error	6.4%	1.2%	0.25%
			0.22%	0.22%
			1.4%	1.4%
			10.7%	10.7%



Parameters of fits to averaged death data.

The best fit ends on Tue 1 Dec 2020.


α	β	k	τ	σ
0.195 ± 0.00973	0.0346 ± 0.00102	1270 ± 2.83	37.2 ± 0.981	25.7 ± 0.505
badness	α	β	k	τ
5.8%	error	5.%	3.%	0.22%
			2.6%	2.6%
			2.%	2.%
			18.6%	18.6%



Somerset, New Jersey

lag from integral overlap, using data averaged over 7 days, between 6 and 10 days, maximum overlap on day 8

lag from fits to the function I (averaged data): 7.75209 days




Parameters of fits to averaged cases data.

The best fit ends on Thu 11 Jun 2020.

α	β	k	τ	σ
0.267 ± 0.00972	0.0777 ± 0.0016	4720 ± 13.3	36.4 ± 0.235	20.3 ± 0.413
badness	α	β	k	τ
1.2%	error	3.6%	2.1%	0.28%

badness	α	β	k	τ	σ	quality	
1.2%	error	3.6%	2.1%	0.28%	0.65%	2.9%	9.85%




Parameters of fits to averaged death data.

The best fit ends on Wed 24 Jun 2020.

α	β	k	τ	σ
0.226 ± 0.00658	0.0652 ± 0.0018	450 ± 1.47	30 ± 0.441	18 ± 0.494
badness	α	β	k	τ
1.5%	error	2.9%	2.8%	0.33%


badness	α	β	k	τ	σ	quality	
1.5%	error	2.9%	2.8%	0.33%	1.5%	2.8%	11.7%



Union, New Jersey

lag from integral overlap, using data averaged over 7 days, between 6 and 12 days, maximum overlap on day 9

lag from fits to the function I (averaged data): 10.757 days




Parameters of fits to averaged cases data.

The best fit ends on Sun 17 May 2020.

α	β	k	τ	σ
0.37 ± 0.0121	0.11 ± 0.00149	$14900. \pm 40.4$	34.2 ± 0.0761	18.7 ± 0.231

badness	α	β	k	τ	σ	quality
0.92%	error	3.3%	1.4%	0.27%	0.22%	1.2% 7.27%




Parameters of fits to averaged death data.

The best fit ends on Sat 9 May 2020.

α	β	k	τ	σ
0.498 ± 0.0209	0.122 ± 0.00124	$941. \pm 5.57$	27.3 ± 0.108	7.35 ± 0.159


badness	α	β	k	τ	σ	quality
0.85%	error	4.2%	1.%	0.59%	0.4%	2.2% 9.21%



Bronx, New York

lag from integral overlap, using data averaged over 7 days, between 0 and 4 days, maximum overlap on day 2


lag from fits to the function I (averaged data): -0.999217 days



Parameters of fits to averaged cases data.

The best fit ends on Tue 14 Apr 2020.


α	β	k	τ	σ
0.502 ± 0.00497	0.139 ± 0.00104	$32900. \pm 251.$	31.7 ± 0.111	15.1 ± 0.0592
badness	α	β	k	τ
0.11%	error	0.99%	0.75%	0.76%
0.35%			0.35%	0.39%
				3.35%



Parameters of fits to averaged death data.

The best fit ends on Sun 17 May 2020.


α	β	k	τ	σ
0.306 ± 0.004	0.0867 ± 0.0049	$3650. \pm 9.76$	19.2 ± 0.18	28.1 ± 0.137
badness	α	β	k	τ
0.43%	error	1.3%	5.7%	0.27%
			0.94%	0.49%
				4.77%



Erie, New York

lag from integral overlap, using data averaged over 7 days, between 5 and 12 days, maximum overlap on day 10

lag from fits to the function I (averaged data): 20.463 days




Parameters of fits to averaged cases data.

The best fit ends on Thu 2 Jul 2020.

α	β	k	τ	σ
0.279 ± 0.0145	0.0545 ± 0.000574	7550 ± 26.2	53.7 ± 0.161	18.4 ± 0.325
badness	error	α	β	k
1.6%		5.2%	1.1%	0.35%

badness	error	α	β	k	τ	σ	quality
1.6%		5.2%	1.1%	0.35%	0.3%	1.8%	10.3%




Parameters of fits to averaged death data.

The best fit ends on Sun 13 Sep 2020.

α	β	k	τ	σ
0.177 ± 0.00385	0.0488 ± 0.000537	678 ± 0.618	44.1 ± 0.273	22.7 ± 0.337
badness	error	α	β	k
1.6%		2.2%	1.1%	0.091%


badness	error	α	β	k	τ	σ	quality
1.6%		2.2%	1.1%	0.091%	0.62%	1.5%	7.04%



Kings, New York

lag from integral overlap, using data averaged over 7 days, between 1 and 6 days, maximum overlap on day 3


lag from fits to the function I (averaged data): 5.82379 days



Parameters of fits to averaged cases data.

The best fit ends on Thu 9 Apr 2020.


α	β	k	τ	σ
0.712 ± 0.0122	0.186 ± 0.00116	$29300 \pm 182.$	29.1 ± 0.0703	14.9 ± 0.0502
badness	error	α	β	k
0.2%		1.7%	0.63%	0.62%
				0.24%
				0.34%
				3.74%



Parameters of fits to averaged death data.

The best fit ends on Wed 22 Apr 2020.


α	β	k	τ	σ
0.408 ± 0.00428	0.148 ± 0.00124	$4400 \pm 14.$	26.9 ± 0.0285	13.9 ± 0.128
badness	error	α	β	k
0.14%		1.%	0.84%	0.32%
				0.11%
				0.92%
				3.37%



Nassau, New York

lag from integral overlap, using data averaged over 7 days, between 5 and 8 days, maximum overlap on day 6


lag from fits to the function I (averaged data): 1.68729 days



Parameters of fits to averaged cases data.

The best fit ends on Sun 19 Apr 2020.


α	β	k	τ	σ
0.541 ± 0.0134	0.157 ± 0.000906	$33900. \pm 124.$	33.8 ± 0.0507	15.9 ± 0.0943
badness	error	α	β	k
0.27%		2.5%	0.58%	0.37%



Parameters of fits to averaged death data.

The best fit ends on Thu 8 Oct 2020.


α	β	k	τ	σ
0.441 ± 0.0105	0.0822 ± 0.000781	$2190. \pm 1.11$	30.1 ± 0.136	21.6 ± 0.103
badness	error	α	β	k
1.9%		2.4%	0.95%	0.051%



New York, New York

lag from integral overlap, using data averaged over 7 days, between 3 and 8 days, maximum overlap on day 5


lag from fits to the function I (averaged data): 1.14256 days



Parameters of fits to averaged cases data.

The best fit ends on Tue 14 Apr 2020.


α	β	k	τ	σ			
0.513 ± 0.00465	0.117 ± 0.000892	$20500. \pm 149.$	33.7 ± 0.127	17.1 ± 0.045			
badness	α	β	k	τ	σ	quality	
0.17%	error	0.91%	0.76%	0.72%	0.38%	0.26%	3.2%



Parameters of fits to averaged death data.

The best fit ends on Sun 17 May 2020.


α	β	k	τ	σ			
0.293 ± 0.00373	0.0825 ± 0.0046	$2320. \pm 6.7$	17.5 ± 0.175	28.4 ± 0.106			
badness	α	β	k	τ	σ	quality	
0.4%	error	1.3%	5.6%	0.29%	1.%	0.37%	4.66%



Queens, New York

lag from integral overlap, using data averaged over 7 days, between 2 and 7 days, maximum overlap on day 5

lag from fits to the function I (averaged data): 2.65965 days




Parameters of fits to averaged cases data.

The best fit ends on Sat 11 Apr 2020.

α	β	k	τ	σ
0.575 ± 0.0065	0.156 ± 0.00122	$41700 \pm 321.$	30.6 ± 0.1	15.8 ± 0.0549

badness	α	β	k	τ	σ	quality
0.16%	error	1.1%	0.78%	0.77%	0.33%	0.35%




Parameters of fits to averaged death data.

The best fit ends on Sun 17 May 2020.

α	β	k	τ	σ
0.295 ± 0.00252	0.0847 ± 0.00305	5640 ± 10.7	21.4 ± 0.126	30.7 ± 0.0906


badness	α	β	k	τ	σ	quality
0.24%	error	0.85%	3.6%	0.19%	0.59%	0.3%



Richmond, New York

lag from integral overlap, using data averaged over 7 days, between -1 and 4 days, maximum overlap on day 2


lag from fits to the function I (averaged data): -3.52476 days



Parameters of fits to averaged cases data.

The best fit ends on Thu 14 May 2020.


α	β	k	τ	σ
0.471 ± 0.0236	0.12 ± 0.00104	13000 ± 27.7	32.3 ± 0.0512	13.7 ± 0.208
badness	α	β	k	τ
0.95%	error	5.%	0.87%	0.21%
			0.16%	1.5%
				8.71%



Parameters of fits to averaged death data.

The best fit ends on Sun 17 May 2020.


α	β	k	τ	σ
0.256 ± 0.00445	0.0572 ± 0.00489	890 ± 10.3	19.3 ± 0.254	24.6 ± 0.256
badness	α	β	k	τ
0.69%	error	1.7%	8.6%	1.2%
			1.3%	1.0%
				7.18%



Rockland, New York

lag from integral overlap, using data averaged over 7 days, between 7 and 10 days, maximum overlap on day 8


lag from fits to the function I (averaged data): 3.1231 days



Parameters of fits to averaged cases data.

The best fit ends on Sun 12 Apr 2020.


α	β	k	τ	σ
0.654 ± 0.012	0.205 ± 0.00111	8760 ± 31.5	29.2 ± 0.0361	15.4 ± 0.0634
<hr/>				
badness	error	α	β	k
0.17%		1.8%	0.54%	0.36%
				0.12%
				0.41%
				3.43%



Parameters of fits to averaged death data.

The best fit ends on Thu 14 May 2020.


α	β	k	τ	σ
0.548 ± 0.021	0.119 ± 0.00168	601 ± 2.32	27 ± 0.0709	14.2 ± 0.137
<hr/>				
badness	error	α	β	k
1.1%		3.8%	1.4%	0.39%
				0.26%
				0.96%
				7.96%



Suffolk, New York

lag from integral overlap, using data averaged over 7 days, between 6 and 13 days, maximum overlap on day 11


lag from fits to the function I (averaged data): 0.186019 days



Parameters of fits to averaged cases data.

The best fit ends on Fri 10 Apr 2020.


α	β	k	τ	σ
0.555 ± 0.0108	0.175 ± 0.0017	33000 ± 526 .	31.5 ± 0.173	13.4 ± 0.0972
badness	α	β	k	τ
0.19%	error	1.9%	0.97%	1.6%



Parameters of fits to averaged death data.

The best fit ends on Sun 15 Nov 2020.


α	β	k	τ	σ
0.42 ± 0.0244	0.0806 ± 0.00072	2000 ± 1.33	41.4 ± 0.104	21.6 ± 0.241
badness	α	β	k	τ
3.5%	error	5.8%	0.89%	0.066%



Westchester, New York

lag from integral overlap, using data averaged over 7 days, between 12 and 14 days, maximum overlap on day 13


lag from fits to the function I (averaged data): 13.9184 days



Parameters of fits to averaged cases data.

The best fit ends on Thu 21 May 2020.

α	β	k	τ	σ			
0.472 ± 0.00843	0.0899 ± 0.000437	33100 ± 39.7	39.7 ± 0.0354	22.7 ± 0.0666			
badness	α	β	k	τ	σ	quality	
0.5%	error	1.8%	0.49%	0.12%	0.089%	0.29%	3.27%




Parameters of fits to averaged death data.

The best fit ends on Tue 30 Jun 2020.

α	β	k	τ	σ			
0.459 ± 0.00873	0.0778 ± 0.000573	1420 ± 1.33	24.6 ± 0.0754	13.6 ± 0.0735			
badness	α	β	k	τ	σ	quality	
1.1%	error	1.9%	0.74%	0.094%	0.31%	0.54%	4.58%


Westchester, New York



Bucks, Pennsylvania

lag from integral overlap, using data averaged over 7 days, between 7 and 13 days, maximum overlap on day 11


lag from fits to the function I (averaged data): 4.31689 days



Parameters of fits to averaged cases data.

The best fit ends on Thu 11 Jun 2020.


α	β	k	τ	σ
0.274 ± 0.00761	0.0763 ± 0.000525	5490 ± 9.95	48.9 ± 0.0564	21 ± 0.211
badness	α	β	k	τ
0.83%	error	2.8%	0.69%	0.18%
			0.12%	1.0%
				5.6%



Parameters of fits to averaged death data.

The best fit ends on Sun 6 Sep 2020.


α	β	k	τ	σ
0.169 ± 0.00337	0.0535 ± 0.00388	582 ± 0.663	29.1 ± 0.713	29.1 ± 1.41
badness	α	β	k	τ
1.7%	error	2.0%	7.3%	0.11%
			2.5%	4.8%
				12.9%



Delaware, Pennsylvania

lag from integral overlap, using data averaged over 7 days, between 7 and 16 days, maximum overlap on day 12


lag from fits to the function I (averaged data): 18.386 days



Parameters of fits to averaged cases data.

The best fit ends on Wed 27 May 2020.


α	β	k	τ	σ
0.247 ± 0.00436	0.0578 ± 0.00086	$7870. \pm 67.5$	53.5 ± 0.292	$26. \pm 0.189$
badness	error	α	β	k
0.6%		1.8%	1.5%	0.86%



Parameters of fits to averaged death data.

The best fit ends on Sat 15 Aug 2020.


α	β	k	τ	σ
0.171 ± 0.00484	0.0416 ± 0.00585	$700. \pm 2.63$	33.4 ± 0.616	33.4 ± 1.56
badness	error	α	β	k
2.4%		2.8%	14.%	0.38%



Montgomery, Pennsylvania

lag from integral overlap, using data averaged over 7 days, between 8 and 12 days, maximum overlap on day 10

lag from fits to the function I (averaged data): 17.4783 days




Parameters of fits to averaged cases data.

The best fit ends on Thu 9 Jul 2020.

α	β	k	τ	σ
0.213 ± 0.00369	0.0442 ± 0.000372	$9100. \pm 21.2$	$59. \pm 0.105$	29.5 ± 0.166

badness	α	β	k	τ	σ	quality
0.82%	error	1.7%	0.84%	0.23%	0.18%	0.56%




Parameters of fits to averaged death data.

The best fit ends on Sat 10 Oct 2020.

α	β	k	τ	σ
0.173 ± 0.00379	0.0357 ± 0.00494	$876. \pm 1.46$	32.6 ± 0.344	32.6 ± 1.02


badness	α	β	k	τ	σ	quality
3.1%	error	2.2%	14.%	0.17%	1.1%	3.1% 12.%



Philadelphia, Pennsylvania

lag from integral overlap, using data averaged over 7 days, between 8 and 12 days, maximum overlap on day 10


lag from fits to the function I (averaged data): -3.30921 days



Parameters of fits to averaged cases data.

The best fit ends on Mon 25 May 2020.


α	β	k	τ	σ			
0.35 ± 0.00916	0.0788 ± 0.0007	$23000. \pm 83.$	44.7 ± 0.0972	21.1 ± 0.145			
badness	α	β	k	τ	σ	quality	
0.68%	error	2.6%	0.89%	0.36%	0.22%	0.69%	5.45%



Parameters of fits to averaged death data.

The best fit ends on Mon 20 Jul 2020.


α	β	k	τ	σ			
0.273 ± 0.0174	0.0628 ± 0.00089	$1660. \pm 4.51$	45.3 ± 0.154	$17. \pm 0.441$			
badness	α	β	k	τ	σ	quality	
2.4%	error	6.4%	1.4%	0.27%	0.34%	2.6%	13.4%



Providence, Rhode Island

lag from integral overlap, using data averaged over 7 days, between 10 and 10 days, maximum overlap on day 10


lag from fits to the function I (averaged data): 3.85265 days



Parameters of fits to averaged cases data.

The best fit ends on Sun 19 Jul 2020.


α	β	k	τ	σ
0.21 ± 0.00743	0.0462 ± 0.00146	$13700. \pm 69.4$	60.7 ± 0.462	46.2 ± 0.445
badness	error	α	β	k
3.4%		3.5%	3.2%	0.51%



Parameters of fits to averaged death data.

The best fit ends on Tue 1 Sep 2020.


α	β	k	τ	σ
0.239 ± 0.025	0.0565 ± 0.00125	$826. \pm 2.45$	48.3 ± 0.315	18.2 ± 0.834
badness	error	α	β	k
4.8%		10.%	2.2%	0.3%



Harris, Texas

no clear lag in the first six months of the pandemic from integral overlap using data averaged over 7 days


lag from fits to the function I (averaged data): 8.16663 days



Parameters of fits to averaged cases data.

The best fit ends on Sat 6 Jun 2020.


α	β	k	τ	σ
0.288 ± 0.00647	0.0285 ± 0.000644	39500 ± 2920	116 ± 4.37	32.9 ± 0.121
badness	α	β	k	τ
1.1%	error	2.2%	2.3%	7.4%
				3.8%
				0.37%
				17.1%



Parameters of fits to averaged death data.

The best fit ends on Tue 5 May 2020.


α	β	k	τ	σ
0.314 ± 0.00716	0.088 ± 0.00199	262 ± 3.31	30.9 ± 0.261	11.9 ± 0.259
badness	α	β	k	τ
0.5%	error	2.3%	2.3%	1.3%
				0.84%
				2.2%
				9.33%



Fairfax, Virginia

lag from integral overlap, using data averaged over 7 days, between -7 and 0 days, maximum overlap on day -2

lag from fits to the function I (averaged data): -9.76437 days




Parameters of fits to averaged cases data.

The best fit ends on Mon 25 May 2020.

α	β	k	τ	σ
0.213 ± 0.00666	0.0745 ± 0.000944	$12800 \pm 134.$	65.8 ± 0.274	26.2 ± 0.425
badness	α	β	k	τ
0.46%	error	3.1%	1.3%	1.%


badness	error	α	β	k	τ	σ	quality
0.46%	error	3.1%	1.3%	1.%	0.42%	1.6%	7.94%



Parameters of fits to averaged death data.

The best fit ends on Sun 30 Aug 2020.


α	β	k	τ	σ
0.174 ± 0.00558	0.036 ± 0.00112	551 ± 2.27	46.8 ± 0.7	34 ± 0.469
badness	α	β	k	τ
2.5%	error	3.2%	3.1%	0.41%



King, Washington

lag from integral overlap, using data averaged over 7 days, between -2 and 2 days, maximum overlap on day 0


lag from fits to the function I (averaged data): 5.00642 days



Parameters of fits to averaged cases data.

The best fit ends on Tue 16 Jun 2020.

α	β	k	τ	σ			
0.171 ± 0.00302	0.036 ± 0.00373	9540 ± 72.7	31.2 ± 0.317	47 ± 0.278			
badness	α	β	k	τ	σ	quality	
0.94%	error	1.8%	10.%	0.76%	1.%	0.59%	6.75%



Parameters of fits to averaged death data.

The best fit ends on Mon 6 Jul 2020.

α	β	k	τ	σ			
0.141 ± 0.00328	0.0437 ± 0.00385	624 ± 2.02	28 ± 0.871	39.6 ± 0.997			
badness	α	β	k	τ	σ	quality	
1.4%	error	2.3%	8.8%	0.32%	3.1%	2.5%	11.9%

