

## SUPPLEMENTARY TABLE

**Table S1: Cox regression survival analysis for the whole cohort, CTX+ST and ST only groups**

Whole cohort		Univariate analysis			Multivariate analysis		
Covariates		HR	95% CI	p-value	HR	95% CI	p-value
Age		1.02	0.99-1.05	0.109			
DM	Ref = absent	2.09	1.03-4.26	0.042			
Asthma	Ref = absent	4.32	0.59-31.71	0.150			
CKD	Ref = absent	8.10	2.49-26.42	0.001	6.29	1.32-29.93	0.021
CAD	Ref = absent	2.08	0.74-5.89	0.166			
BMI		0.88	0.75-1.04	0.136			
CRP day 0		1.02	1.009-1.03	<0.001			
CXR/CT		9.19	3.08-27.44	<0.001			
NLR day 0		1.032	0.99-1.07	0.064			
R/R		1.19	1.12-1.29	<0.001	1.19	1.08-1.31	0.001
Temp		1.28	1.01-1.63	0.041			
CTX+ST group		Univariate analysis			Multivariate analysis		
Covariates		HR	95% CI	p-value	HR	95% CI	p-value
Age		1.06	1.02-1.11	0.009	1.09	1.01-1.185	0.027
DM	Ref = absent	2.41	0.91-6.39	0.078			
Asthma	Ref = absent	7.64	1.01-57.88	0.049	562.67	8.01-39509.33	0.004
CKD	Ref = absent	24.56	2.41-250.42	0.007			
CAD	Ref = absent	2.93	0.82-10.38	0.095			
CRP at day 0		1.03	1.01-1.04	0.001			
NLR at day 0		1.05	1.01-1.09	0.022			
R/R		1.34	1.19-1.50	<0.001	1.37	1.16-1.63	<0.001
ST only group		Univariate analysis			Multivariate analysis		
Covariates		HR	95% CI	p-value	HR	95% CI	p-value
Male sex	Ref = absent	2.29	0.66-7.95	0.191			
Age		0.96	0.91-1.01	0.132			
CRP at day 0		1.02	0.99-1.04	0.088			
CXR/CT		12.73	2.99-54.17	0.001			
R/R		1.20	1.02-1.41	0.028			
Temp		2.61	1.43-4.77	0.002	2.68	1.09-6.54	0.031

Definition of abbreviations: SD = Standard Deviation; CTX = Cotrimoxazole; ST = Standard Therapy; HR = Hazard Ratio; CI = Confidence Interval; DM = Diabetes Mellitus; CKD = Chronic Kidney Disease; CAD = Coronary Artery Disease; BMI = Body Mass Index; CRP = C-reactive protein; NLR = Neutrophil-Lymphocyte ratio; CXR = Chest x-ray; R/R = Respiratory rate; Temp = Temperature