

List of supplementary tables

Supplementary Table 1: Sample demographics and descriptive variables (weighted)	2
Supplementary Table 2a: Regression analyses of co-occurrence of sleep disorders and psychotic experiences at baseline (T0) – weighting method A	3
Supplementary Table 2b: Regression analyses of co-occurrence of sleep disorders and psychotic experiences at baseline (T0) – weighting method B	4
Supplementary Table 3a: Regression analyses of whether sleep disorders at baseline predict onset of psychotic experiences at 12 months – weighting method A	5
Supplementary Table 3b: Regression analyses of whether sleep disorders at baseline predict onset of psychotic experiences at 12 months – weighting method B	6
Supplementary Table 4a: Regression analyses of whether sleep disorders at baseline predict persistence of psychotic experiences at 12 months – weighting method A	7
Supplementary Table 4b: Regression analyses of whether sleep disorders at baseline predict persistence of psychotic experiences at 12 months – weighting method B	8
RQ4a: Regression analysis of whether remission of sleep problems predicts remission of psychotic experiences (weighted method A):	9
RQ4a: Regression analysis of whether remission of sleep problems predicts remission of psychotic experiences (weighted method B):	9

Supplementary Table 1: Sample demographics and descriptive variables (weighted)

Descriptive statistics	Baseline	12-month follow-up
	n = 11830	n = 4910
Age (months) – mean (SE)	119.2 (0.3)	132.2 (0.5)
Gender – male % (SE)	51.13 (0.1)	51.04 (0.9)
Ethnicity % (SE)		
White	52.34 (5.6)	59.48 (6.3)
Black	13.31 (2.5)	7.61 (1.6)
Hispanic	23.83 (6.2)	22.50 (6.7)
Asian	3.61 (0.9)	4.02 (0.9)
Other	6.89 (1.0)	6.40 (1.3)
Socioeconomic status scale – mean (SE)	0.56 (0.1)	0.47 (0.1)
Child IQ – mean (SE)	9.74 (0.1)	10.03 (0.1)
Neighbourhood deprivation percentile – mean (SE)	43.49 (3.8)	40.5 (3.9)
Family conflict scale – mean (SE)	2.52 (0.1)	2.45 (0.06)
Stimulant medication prescribed – % (SE)	6.41 (<0.0)	7.00 (<0.0)
PQ-BC total – mean (SE)	2.74 (0.3)	1.82 (0.2)
SDSC total – mean (SE)	36.78 (0.3)	36.59 (0.3)

Derived variable counts

Sleep disorders present (≥ 39 cut off on SDSC) – % (SE)	31.71 (1.5)	29.15 (1.6)
Sleep disorders new onset (<i>absent at baseline, present at follow up</i>) – % (SE)		9.56 (0.1)
Sleep disorders persisting (<i>present at both baseline and follow up</i>) – % (SE)		19.50 (1.6)
Sleep disorders remitting (<i>present at baseline, absent at follow up</i>) – % (SE)		10.48 (0.8)
Psychotic experiences present (≥ 1 on PQ-BC total) – % (SE)	63.08 (2.6)	48.25 (2.8)
Psychotic experiences new onset – % (SE)		9.24 (0.9)
Psychotic experiences persisting – % (SE)		29.01 (3.0)
Psychotic experiences remitting – % (SE)		24.51 (1.1)

Cross tabulation (sleep disorders vs psychotic experiences, % prevalence)

		Psychotic experience onset	
		No	Yes
Onset (absent at baseline, present at follow up)	Sleep disorder onset	No	82.10
		Yes	8.66
		Psychotic experience persisting	
		No	51.17
Persistence (present at baseline and follow up)	Sleep disorder persisting	Yes	29.33
		No	9.84
		Psychotic experience remitting	
		No	21.81
Remission (present at baseline, absent at follow up)	Sleep disorder remission	Yes	2.71
		No	67.70

SDSC = Sleep Disorder Scale for Children; PQBC = Prodromal Questionnaire – Brief Child version; SE = standard error;

Supplementary Table 2a: Regression analyses of co-occurrence of sleep disorders and psychotic experiences at baseline (T0) – weighting method A

Model (outcome)	Parameters	B	95% CI	p-value	AIC
Model 1a (PQBC total T0) ^a	SDSC total T0	0.051	0.04, 0.06	<0.001	26047
Model (outcome)	Parameters	Odds	95% CI	p-value	AIC
Model 1b (PQBC cut-off T0) ^b	SDSC total T0	1.034	1.02, 1.04	<0.001	6053
Model 1c (PQBC cut-off T0) ^b	SDSC cut off T0	1.614	1.37, 1.9	<0.001	6067
Model (outcome)	Parameters	B	95% CI	p-value	AIC
Model 1d (PQBC total T0) ^a	SDSC total T0	0.040	0.03, 0.05	<0.001	24325
	Gender	-0.248	-0.45, -0.05	0.014	
	Ethnicity	0.055	-0.03, 0.14	0.220	
	Socioeconomic status	0.171	0.07, 0.28	0.001	
	Neighbourhood deprivation	0.011	0.01, 0.02	<0.001	
	IQ	-0.053	-0.09, -0.02	0.003	
	Family conflict	0.011	-0.04, 0.07	0.688	
	Stimulant medication	0.581	0.18, 0.98	0.004	
Model (outcome)	Parameters	Odds	95% CI	p-value	AIC
Model 1e (PQBC cut-off T0) ^b	SDSC total T0	1.026	1.02, 1.04	<0.001	5641
	Gender	0.835	0.72, 0.97	0.021	
	Ethnicity	1.032	0.96, 1.11	0.402	
	Socioeconomic status	1.155	1.06, 1.26	0.001	
	Neighbourhood deprivation	1.010	1.01, 1.01	<0.001	
	IQ	0.942	0.92, 0.97	<0.001	
	Family conflict	1.004	0.96, 1.05	0.852	
	Stimulant medication	1.208	0.88, 1.66	0.249	
Model 1f (PQBC cut off T0) ^b	SDSC cut off T0	1.470	1.23, 1.75	<0.001	5647
	Gender	0.834	0.72, 0.97	0.020	
	Ethnicity	1.034	0.96, 1.11	0.378	
	Socioeconomic status	1.167	1.07, 1.27	<0.001	
	Neighbourhood deprivation	1.010	1.01, 1.01	<0.001	
	IQ	0.942	0.92, 0.97	<0.001	
	Family conflict	1.010	0.97, 1.05	0.638	
	Stimulant medication	1.237	0.90, 1.70	0.192	

^a = linear regression; ^b = logistic regression; SDSC = Sleep Disorder Scale for Children; PQBC = Prodromal Questionnaire – Brief Child version; B = standardised beta; CI = confidence intervals; AIC = Akaike information criterion.

Supplementary Table 2b: Regression analyses of co-occurrence of sleep disorders and psychotic experiences at baseline (T0) – weighting method B

Model (outcome)	Parameters	B	95% CI	p-value	AIC
Model 1a (PQBC total T0) ^a	SDSC total	0.051	0.04, 0.06	<0.001	26054
Model (outcome)	Parameters	Odds	95% CI	p-value	AIC
Model 1b (PQBC cut-off T0) ^b	SDSC total	1.029	1.02, 1.04	<0.001	5027
Model 1c (PQBC cut-off T0) ^b	SDSC cut off	1.501	1.29, 1.75	<0.001	5039
Model (outcome)	Parameters	B	95% CI	p-value	AIC
Model 1d (PQBC total T0) ^a	SDSC total	0.040	0.03, 0.05	<0.001	24333
	Gender	-0.247	-0.45, -0.05	0.015	
	Ethnicity	0.057	-0.03, 0.15	0.207	
	Socioeconomic status	0.170	0.07, 0.27	0.001	
	Neighbourhood deprivation	0.011	0.01, 0.02	<0.001	
	IQ	-0.053	-0.09, -0.02	0.003	
	Family	0.011	-0.04, 0.07	0.694	
	Stimulant medication	0.582	0.18, 0.98	0.004	
Model (outcome)	Parameters	Odds	95% CI	p-value	AIC
Model 1e (PQBC cut-off T0) ^b	SDSC total	1.022	1.01, 1.03	<0.001	4687
	Gender	0.876	0.76, 1.01	0.066	
	Ethnicity	1.037	0.97, 1.11	0.296	
	Socioeconomic status	1.147	1.06, 1.24	0.001	
	Neighbourhood deprivation	1.008	1.00, 1.01	<0.001	
	IQ	0.952	0.93, 0.98	<0.001	
	Family	1.002	0.96, 1.04	0.910	
	Stimulant medication	1.151	0.86, 1.55	0.353	
Model 1f (PQBC cut off T0) ^b	SDSC cut off	1.374	1.17, 1.62	<0.001	4693
	Gender	0.875	0.76, 1.01	0.063	
	Ethnicity	1.038	0.97, 1.11	0.280	
	Socioeconomic status	1.157	1.07, 1.25	<0.001	
	Neighbourhood deprivation	1.008	1.00, 1.01	<0.001	
	IQ	0.952	0.93, 0.98	<0.001	
	Family	1.008	0.97, 1.05	0.693	
	Stimulant medication	1.180	0.88, 1.59	0.274	

^a = linear regression; ^b = logistic regression; SDSC = Sleep Disorder Scale for Children; PQBC = Prodromal Questionnaire – Brief Child version; B = standardised beta; CI = confidence intervals; AIC = Akaike information criterion.

Supplementary Table 3a: Regression analyses of whether sleep disorders at baseline predict onset of psychotic experiences at 12 months – weighting method A

Model (outcome)	Parameters	Odds	95% CI	p-value	AIC
Model 2a (PQBC cut off T1)	SDSC cut off T0	1.719	1.47, 2.01	<0.001	6303
Model 2b (PQBC cut off T1)	SDSC cut off T0	1.482	1.24, 1.77	<0.001	6294
	SDSC cut off T1	1.346	1.12, 1.61	0.001	
Model 2c (PQBC cut off T1)	SDSC cut off T0	1.511	1.29, 1.77	<0.001	5826
	Gender	0.808	0.7, 0.93	0.003	
	Ethnicity	1.066	1.00, 1.13	0.041	
	Socioeconomic status	1.231	1.14, 1.33	<0.001	
	Neighbourhood deprivation	1.009	1.01, 1.01	<0.001	
	IQ	0.969	0.95, 0.99	0.013	
	Family conflict	0.994	0.96, 1.03	0.749	
	Stimulant medication	1.702	1.28, 2.26	<0.001	
Model 2d (PQBC cut off T1)	SDSC cut off T0	1.318	1.32, 1.32	<0.001	5819
	SDSC cut off T1	1.328	1.33, 1.33	<0.001	
	Gender	0.798	0.80, 0.80	<0.001	
	Ethnicity	1.064	1.06, 1.07	<0.001	
	Socioeconomic status	1.220	1.22, 1.22	<0.001	
	Neighbourhood deprivation	1.009	1.01, 1.01	<0.001	
	IQ	0.969	0.97, 0.97	<0.001	
	Family conflict	0.992	0.99, 0.99	<0.001	
	Stimulant medication	1.677	1.67, 1.68	<0.001	

SDSC = Sleep Disorder Scale for Children; PQBC = Prodromal Questionnaire – Brief Child version; CI = confidence intervals; AIC = Akaike information criterion

Supplementary Table 3b: Regression analyses of whether sleep disorders at baseline predict onset of psychotic experiences at 12 months – weighting method B

Model (outcome)	Parameters	Odds	95% CI	p-value	AIC
Model 2a (PQBC cut off T1)	SDSC cut off T0	1.568	1.36, 1.81	<0.001	5226
Model 2b (PQBC cut off T1)	SDSC cut off T0	1.391	1.18, 1.64	<0.001	5220
	SDSC cut off T1	1.268	1.07, 1.50	0.006	
Model 2c (PQBC cut off T1)	SDSC cut off T0	1.384	1.18, 1.62	<0.001	4836
	Gender	0.862	0.75, 0.99	0.035	
	Ethnicity	1.072	1.00, 1.15	0.039	
	Socioeconomic status	1.214	1.13, 1.30	<0.001	
	Neighbourhood deprivation	1.007	1.00, 1.01	<0.001	
	IQ	0.975	0.95, 1.00	0.049	
	Family conflict	0.998	0.96, 1.04	0.906	
	Stimulant medication	1.583	1.19, 2.10	0.001	
Model 2d (PQBC cut off T1)	SDSC cut off T0	1.248	1.04, 1.49	0.015	4832
	SDSC cut off T1	1.242	1.04, 1.48	0.017	
	Gender	0.857	0.75, 0.98	0.029	
	Ethnicity	1.071	1.00, 1.14	0.042	
	Socioeconomic status	1.205	1.12, 1.30	<0.001	
	Neighbourhood deprivation	1.007	1.00, 1.01	<0.001	
	IQ	0.975	0.95, 1.00	0.045	
	Family conflict	0.995	0.96, 1.03	0.798	
	Stimulant medication	1.546	1.16, 2.05	0.003	

SDSC = Sleep Disorder Scale for Children; PQBC = Prodromal Questionnaire – Brief Child version; CI = confidence intervals; AIC = Akaike information criterion

Supplementary Table 4a: Regression analyses of whether sleep disorders at baseline predict persistence of psychotic experiences at 12 months – weighting method A

Model (outcome)	Parameters	Odds	95% CI	p-value	AIC
Model 3a (PQBC persist)	SDSC cut off T0	1.841	1.57, 2.16	<0.001	5996
Model 3b (PQBC persist)	SDSC persist	2.058	1.71, 2.48	<0.001	5994
Model 3c (PQBC persist)	SDSC cut off T0	1.598	1.60, 1.60	<0.001	5537
	Gender	0.808	0.81, 0.81	<0.001	
	Ethnicity	1.022	1.02, 1.02	<0.001	
	Socioeconomic status	1.220	1.22, 1.22	<0.001	
	Neighbourhood deprivation	1.009	1.01, 1.01	<0.001	
	IQ	0.960	0.96, 0.96	<0.001	
	Family conflict	1.002	1.00, 1.00	<0.001	
	Stimulant medication	1.508	1.51, 1.51	<0.001	
Model 3d (PQBC persist)	SDSC persist	1.795	1.46, 2.20	<0.001	5532
	Gender	0.813	0.69, 0.95	0.010	
	Ethnicity	1.025	0.95, 1.10	0.517	
	Socioeconomic status	1.217	1.12, 1.32	<0.001	
	Neighbourhood deprivation	1.010	1.01, 1.01	<0.001	
	IQ	0.963	0.94, 0.99	0.008	
	Family conflict	1.004	0.96, 1.05	0.844	
	Stimulant medication	1.567	1.15, 2.14	0.005	

persist = above cut off at T0 and T1; SDSC = Sleep Disorder Scale for Children; PQBC = Prodromal Questionnaire – Brief Child version; CI = confidence intervals; AIC = Akaike information criterion.

Supplementary Table 4b: Regression analyses of whether sleep disorders at baseline predict persistence of psychotic experiences at 12 months – weighting method B

Model (outcome)	Parameters	Odds	95% CI	p-value	AIC
Model 3a (PQBC persist)	SDSC cut off T0	1.671	1.44, 1.94	<0.001	4989
Model 3b (PQBC persist)	SDSC persist	1.828	1.54, 2.17	<0.001	4987
Model 3c (PQBC persist)	SDSC cut off T0	1.473	1.25, 1.73	<0.001	4609
	Gender	0.862	0.75, 1.00	0.043	
	Ethnicity	1.042	0.97, 1.12	0.237	
	Socioeconomic status	1.208	1.13, 1.30	<0.001	
	Neighbourhood deprivation	1.007	1.00, 1.01	<0.001	
	IQ	0.969	0.94, 0.99	0.017	
	Family conflict	1.006	0.97, 1.05	0.744	
	Stimulant medication	1.407	1.06, 1.86	0.017	
Model 3d (PQBC persist)	SDSC persist	1.598	1.33, 1.92	<0.001	4321
	Gender	0.862	0.75, 0.99	0.042	
	Ethnicity	1.043	0.97, 1.12	0.232	
	Socioeconomic status	1.203	1.12, 1.29	<0.001	
	Neighbourhood deprivation	1.007	1.00, 1.01	<0.001	
	IQ	0.969	0.94, 0.99	0.016	
	Family conflict	1.007	0.97, 1.05	0.717	
	Stimulant medication	1.406	1.06, 1.86	0.018	

persist = above cut off at T0 and T1; SDSC = Sleep Disorder Scale for Children; PQBC = Prodromal Questionnaire – Brief Child version; CI = confidence intervals; AIC = Akaike information criterion.

RQ4a: Regression analysis of whether remission of sleep problems predicts remission of psychotic experiences (weighted method A):

Remission of sleep disorder symptoms was not a significant predictor of remission of psychotic experiences (OR=1.041, p=0.588, 95% CI 0.80, 1.35, p=0.766, AIC=5241) in the uncontrolled model, therefore the analysis with control variables was not carried out.

RQ4a: Regression analysis of whether remission of sleep problems predicts remission of psychotic experiences (weighted method B):

Remission of sleep disorder symptoms was not a significant predictor of remission of psychotic experiences (OR=1.011, p=0.588, 95% CI 0.79, 1.29, p=0.929) in the uncontrolled model, therefore the analysis with control variables was not carried out.