

Psychological Legacies of Intergenerational Trauma under South African Apartheid: Prenatal Stress Predicts Increased Psychiatric Morbidity during Late Adolescence and Early Adulthood in Soweto, South Africa

Andrew Wooyoung Kim¹⁻³, Rihlat Said Mohamed⁴, Shane A. Norris^{2,5-6}, Linda M. Richter⁶, Christopher W. Kuzawa^{1,7}

¹ Center for Global Health, Massachusetts General Hospital, Harvard Medical School, Boston, MA, United States

² SAMRC/Wits Developmental Pathways for Health Research Unit, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

³ Department of Global Health and Population, Harvard T. H. Chan School of Public Health, Boston, Massachusetts, USA

⁴ Department of Archaeology, University of Cambridge, Cambridge, UK

⁵ Global Health Research Institute, School of Human Development and Health, University of Southampton, UK

⁶ DSI-NRF Centre of Excellence in Human Development, University of the Witwatersrand, Johannesburg, South Africa

⁷ Department of Anthropology, Northwestern University, Evanston, IL, United States

⁸ Institute for Policy Research, Northwestern University, Evanston, IL, United States

ABSTRACT

Background: South Africa's rates of psychiatric morbidity are among the highest in sub-Saharan Africa and are foregrounded by the country's long history of political violence during apartheid. Recent evidence suggests that maternal trauma during gestation may intergenerationally impact the developing fetus and elevate the future child's risk for psychiatric disease. We aim to evaluate the intergenerational effects of prenatal stress experienced during apartheid on late adolescent psychiatric morbidity and also assess the potential ameliorative effects of prenatal social support.

Method: Participants (n = 1051) come from Birth-to-Twenty, a longitudinal birth cohort study in Soweto-Johannesburg, South Africa's largest peri-urban township which was the epicenter of violent repression and resistance during the final years of the apartheid regime. Pregnant women were prospectively enrolled in 1990 and completed questionnaires assessing social experiences, and their children's psychiatric morbidity were assessed at age 17-18.

Results: Full data were available from 304 mother-child pairs in 2007-8. Pregnant women who experienced worse traumatic stress in 1990 had children who exhibited greater psychiatric morbidity during late adolescence. This relationship was only significant, however, among children born to younger mothers and adolescents who experienced greater concurrent stress. Social support did not buffer the long-term impacts of prenatal stress on future psychiatric morbidity.

Conclusion: Greater prenatal stress predicted adverse psychiatric outcomes among children born to younger mothers and adolescents who experienced greater concurrent stress. Our findings suggest that prenatal stress may affect adolescent mental health, have stress-sensitizing effects, and represent possible intergenerational effects of trauma experienced under apartheid in this sample.

INTRODUCTION

South Africa's rates of mental, neurological, and substance use disorders are among the highest in sub-Saharan Africa. In 2016, the estimated 12-month prevalence for any psychiatric disorder was 16.2%, or approximately 9.1 million individuals (GBDCN 2017). Despite these elevated rates of psychiatric morbidity, access to mental health treatment is poor: only 27% of patients living with severe mental illness are expected to receive treatment (Herman et al. 2009). The current state of public mental health in South Africa is foregrounded by a long and recent history of institutionalized White supremacist policies implemented during the *Apartheid* regime (c. 1948-1994). This period was characterized by systematic disenfranchisement of non-White communities through various modes of social, economic, and political oppression (Beinhart 2001). Despite the legislative end of *Apartheid* policies and the birth of a new democratic in 1994, South African society continues to be plagued by the persistent societal institutions of *Apartheid*, including chronic poverty (Gibbs et al. 2018), discrimination (Kuzawa & Sweet 2009), and racialized class inequality (Adjaye-Gbewonyo et al. 2016; Burns 2015), all of which are known risk factors for psychiatric disorders (Allen et al. 2014; Moomal et al. 2009; Myer et al. 2008).

In addition to these ongoing societal effects of poverty, structural violence and inequality, there is growing evidence, from South African studies and other populations, that the stressors of the past could have lingering biological effects that continue to influence socio-emotional behavior and mental health across the lifecourse. Growing evidence from the fetal origins of health and disease framework shows that past stress and trauma exposures, particularly those that occur during early development, can durably alter the development and function of various stress regulatory mechanisms in humans, including the immune system, cardiovascular system, neurobiological function, and neuroendocrine pathways (Barker et al. 1999; Gluckman & Hanson 2004; Kuzawa 2008; Taylor et al. 2010; Heim et al. 2019). Recent findings also suggest that these stress-linked alterations in stress physiology may also affect the developing offspring through various pathways of genetic and non-genetic inheritance. For instance, large population-based studies in the United Kingdom show that greater levels of maternal prenatal stress are associated with an increased future risk for externalizing disorders in their children, such as attention

deficit hyperactivity disorder (Rice et al. 2010), conduct disorders (MacKinnon et al. 2018), and internalizing disorders like anxiety and depression among adolescents (O'Donnell et al. 2014; Sharp et al. 2015). Additionally, the intergenerational signatures of maternal prenatal stress exposure have been reported in offspring during adulthood in large cohort studies and across various contexts such as Australia, the Philippines, and the United States (Betts et al. 2015; DeSantis et al. 2015; Entringer et al. 2009).

While the specific biological mechanisms that underlie the long-term psychiatric effects of prenatal stress are unknown, growing evidence from the literature on the fetal origins of psychopathology suggests that prenatal stress exposure alters the development, function, sensitivity of human stress physiological systems across the life course, which in turn may elevate one's risk for a psychopathological presentation. These stress-sensitive systems include the immune system, the catecholamine-producing sympathetic-adrenal-medullary (SAM) axis, and the hypothalamic-pituitary-adrenal (HPA) axis, which regulates physiological reactions to stress through stress-sensitive hormones like cortisol. Maternal stress-induced elevations in cortisol can enter maternal circulation, penetrate the placental wall, and reach the gestational environment of future offspring, thereby impacting the development of fetal stress physiology (D'Anna-Hernandez et al. 2012; O'Donnell et al. 2009).

Greater fetal exposure to cortisol is understood influence a wide variety of biological and health phenotypes across the lifecourse. These include birth outcomes such as restricted fetal growth rates, shorten gestations, and reduce birth size (Kim et al. 2020a; Rosa et al. 2019; Ryu 2019). Greater intrauterine cortisol exposure may also durably alter the development and sensitivity of the fetal stress physiology (e.g. HPA axis, immune system, brain development), which together with the maternal and placental systems, may have sustained impacts on the child's stress physiology across their lifecourse (Brand et al. 2010; DeSantis et al. 2015; Entringer et al. 2009; Karlén et al. 2013). Early life stress-linked dysregulation of stress physiological mechanisms, characterized by altered diurnal cortisol rhythms, glucocorticoid resistance, chronic low-grade inflammation, and modulated brain function, has consistently been reported as both a prospective risk factor and cross-sectional neuropsychiatric

phenotype of a range of psychiatric illnesses, including depression, psychosis, suicidal ideation, schizophrenia, across the lifespan (Doane et al. 2013; Heim et al. 2019; Jarcho et al. 2013; Miller et al. 2011; Taylor 2010; Vrshek-Schallhorn et al. 2013).

Additionally, recent evidence has also suggested that alterations in stress physiological systems (e.g. neuroendocrine, inflammatory, molecular, and structural pathways) due to developmental stress exposure may also alter sensitization to future stressors. The stress sensitization hypothesis (Hammen et al. 2000) proposes that the risk for adult mental illness following stressful life events is higher among individuals with a history of developmental trauma than among individuals without a history of developmental trauma. Recent findings have extended this hypothesis to suggest that greater early life trauma sensitizes or potentiates future reactions to stress and consequently, increases psychiatric disease risk (Heim et al. 2019; McLaughlin et al. 2010; Shapero et al. 2014). While an increasing number of studies are noting the long-term impacts of prenatal stress on infant, child, and early/mid adolescent psychological status (Ilg et al. 2019; Koss & Gunnar 2018; Ping et al. 2020; Ziljmans et al. 2015), few studies have examined the mental health impacts of prenatal stress into late adolescence and adulthood and also evaluated the potential stress sensitization effects of prenatal stress (Koss & Gunnar 2018; Ping et al. 2020; Ziljmans et al. 2015).

This emerging evidence suggests that some proportion of the mental health burden of current populations could reflect the lingering biological imprint of past traumatic experience during gestation (Barker 1999; Krontira et al. 2020; Kuzawa & Sweet 2009). South Africa's recent history and the country's tumultuous transition into democracy raise the question of whether the traumas of *Apartheid* may continue to have lasting effects, influencing risk for psychiatric morbidity across generations (Kim 2020). It furthermore raises the question of whether some of these effects might be reversible, opening opportunities for public health interventions to reduce the societal burden of psychiatric morbidity (Heim et al. 2019). However, little work in South Africa has explored this hypothesis and its contribution to mental health. Furthermore, as the most current research is conducted in high-income, Western contexts, it remains unclear if these effects are seen in non-Western, low- and middle-income contexts like South

Africa, where adverse psychosocial and environmental conditions are more prevalent and where the burden of mental illness is substantially greater (Patel 2007; Vigo et al. 2016).

This study examines the long-term impacts of prenatal stress and trauma from South African *Apartheid* on psychiatric morbidity during late adolescence/early adulthood and the possible reversibility of the long-term effects of prenatal stress in a longitudinal birth cohort study. Adolescence is an important period in the development of future mental illness as most psychiatric conditions emerge during this stage (Patel 2007; Breslau et al. 2017) and because adolescent psychiatric disease is a major risk factor for future adult mental illness (Pine et al. 1993; Sawyer et al. 2012). The recent history of *Apartheid*, high rates of mental illness, and low rates of healthcare access emphasize the need to elucidate possible mechanisms underlying the intergenerational mental health effects of prenatal trauma in order to improve public mental health in South Africa.

METHODS

Study setting and participants

Data come from a longitudinal birth cohort in South Africa called Birth to Twenty Plus (Bt20++), which currently spans three generations of families in the greater Johannesburg-Soweto area. The Birth to Twenty (Bt20++) study is both the largest and longest running longitudinal birth cohort study of child health and development in Africa. Bt20++ emerged as a collaboration between the University of the Witwatersrand in Johannesburg and the South African Medical Research Council with the aim to assess the impacts of rapid urbanization towards the end of *Apartheid* on the growth, health, well-being, and educational progress of children. Soweto is the largest township outside of the city of Johannesburg and a site of immense cultural and historical significance in the struggle against *Apartheid*. Racial and political violence, government divestment, and widespread protest were common in both cities, particularly at the legislative end of *Apartheid*, which is when pregnant women were first recruited into the study.

All singleton live births delivered in public sector hospitals between April 23 to June 8, 1990 and who were residents in the metropolitan Johannesburg-Soweto area six months after delivery were enrolled

in Bt20+ (Richter et al. 2007). In late 1989, BT20+ began interviewing pregnant women in public antenatal clinics to identify potential participants whose births would fall within the enrollment period. Enrolled Bt20+ neonates were cross-checked with all government birth notifications during the 7-week time period. The study area covered approximately 78 square miles at the time and included close to 3.5 million people, with about 400,000 informal housing units.

A total of 3273 singleton children were enrolled in Bt20+ and 72% of the initial cohort continued to participate in the study after 17 years (Norris et al. 2007). The cohort is roughly representative of the demographic parameters of the metropolitan Johannesburg-Soweto region. Currently, the cohort underrepresents White children due to cohort enrollment taking place in public health facilities; White families were more likely to utilize private practitioners and facilities. Although BT20+ children were all *in utero* during the *Apartheid* regime, they became among the first generation born into a democratic South Africa, and colloquially known as “Mandela’s Children” because they were born shortly after Nelson Mandela’s release from prison on February 11, 1990. Between 1990 and 2007, the scope of this analysis, BT20+ families participated in 18 waves of data collection, and follow-up studies continue. All participants provided assent and their parents provided written, informed consent. Ethical approval was obtained from the University of Witwatersrand Committee for Research on Human Subjects.

Before the full sample was achieved, 1594 women were interviewed during their third trimester about their pregnancy, social experience, and household conditions. Antenatal interviews were conducted by seven trained, multilingual, interviewers. Where translation of measures was required, consensual agreement on the phrasing of questions was reached. Continuous translation and back-translation were used in order to ensure that the meaning/s attained in the destination language mirrored those intended in the original one. The majority of interviews were conducted in antenatal services, with a quarter conducted at home. Zulu, Sotho and English were the most common languages used. Adolescents were interviewed in a research facility at the Chris Hani-Baragwanath Hospital in Soweto and at home.

Prenatal stress and social support

Prenatal stress exposure (G1) during the third trimester of pregnancy was collected using a 16-item scale ($\alpha = 0.64$), adapted from Bluen et al. (1988) capturing information on exposure to various forms of family and community stresses, including police violence, injury, and incarceration. Yes/no responses indicated the presence or absence of each stressor during the previous 6 months. After exploratory analyses, two questions were dropped: one due to a low response rate (have you experienced any problems with your other children?) and the other due to contradictory concepts being assessed at once (domestic and familial violence vs. partner separation). Of the initial 1594 women, 1051 women (65.9%) completed all of the remaining prenatal stress questions.

Social support was measured using a series of four, yes/no questions to identify the absence or presence of instrumental and emotional support, including: people available to help, a confidante, being able to speak to her partner, belonging to a community organisation/church. All questions were summed to create a total score.

Demographic, Health, and Socioeconomic Variables

During the antenatal visit in 1990, expectant women were asked about their current housing situation (e.g. number of rooms, number of inhabitants), population group (i.e. “race”), marital status, age, and obstetric history. Gravidity was calculated from information on obstetric history and operationalized into a categorical yes/no variable (0 = no past pregnancy, 1 = past history of pregnancy). They were also asked whether they used tobacco or had been drinking alcohol during their pregnancy. At delivery, a series of questions about the neonate were administered to gather information about gender, birth order, birthweight, and gestational age. Birthweight (g) and gestational age (weeks) were obtained from the child’s Road to Health Card, a patient-held child medical record provided to all new mothers in South Africa.

Household socio-economic status (SES) was assessed using an asset index which scored each participant according to the number of household physical assets that they possessed out of a possible 6

(e.g. television, refrigerator, washing machine, radio, telephone, home ownership, car). Household SES was measured again in 2006 using an updated list of assets (e.g. television, car, washing machine, refrigerator, phone, radio, microwave, cell phone, DVD, MNET, DSTV, computer, and internet). The asset index was designed based on standard measures used by the Demographic and Health Surveys (<https://dhsprogram.com/>), based on the work of Filmer and Pritchett (1999). To capture the overall SES environment during prenatal development and 2007, when the outcome measure was administered, an aggregate SES variable was constructed by summing standardized assets scores from both years of data collection.

Household Stressful Life Events

The occurrence of stressful life events in the household were assessed across two timescales, six months and twelve months, and reported by the caregiver of the index child based on yes/no responses. Thirteen life events were assessed and summed together to create a composite measure of household stress that occurred during the past six months, which included death of a sibling, parent, and other family member (three separate questions), divorce, index child changing schools, a serious illness or hospitalization experienced by the index child, caregiver, and family member (three separate questions), marital separation, increase in arguments with partner, caregiver separation from family for two weeks or more, a child leaving home, and unemployment. Additionally, eight life events that occurred in the last year were summed to create a separate composite measure, which assessed whether any member of the household experienced robbery, harassment, threats, sexual molestation, physical violence, and murder.

General Health Questionnaire (GHQ-28)

The General Health Questionnaire (GHQ-28) is a psychological screener that provides a measure of psychiatric morbidity based on four 7-item scales: somatic symptoms, anxiety and insomnia, social dysfunction, and severe depression (Goldberg & Hillier 1979). It assesses changes in mood, feelings, and behaviours during the past four weeks. The respondent evaluates their occurrence on a 4-point Likert

scale, “less than usual,” “no more than usual,” “rather more than usual,” and “much more than usual.”

Seven questions are reverse scored and transformed before all responses summed. Bt20+ index children and their caregivers completed the GHQ during a follow-up wave of data collection when the index children were 17-18 years old.

Statistical analyses

All analyses were conducted using version 15.1 of Stata (Stata Corporation, College Station, TX). All variables were examined for normal distribution and outliers. Bivariate analyses were conducted between prenatal stress, psychiatric morbidity at age 18, and covariates. Covariates were included based on a priori knowledge of social, biological, and obstetric risk factors that may potentially confound the relationship between prenatal stress and later-life psychiatric morbidity (de Mola et al. 2014; Entringer et al. 2009; O’Donnell et al. 2013). The following variables were considered for inclusion as confounding factors: maternal age, maternal education, marital status, gravidity, tobacco use, and alcohol consumption during pregnancy, household density (ratio of the number of inhabitants to the number of rooms available for sleeping), social support, and child gender. ($p > 0.10$). Fetal growth rate and gestational age were also included as key covariates to examine the role of fetal growth restriction and preterm delivery as possible pathways by which prenatal stress affects late adolescent psychiatric morbidity. A proxy measure of fetal growth was created by regressing birthweight on gestational age to create standardized residuals.

Bivariate analyses were conducted to identify potential confounding variables for inclusion in the final analytic models. With the exception of known confounding factors for the relationship between prenatal stress and later life psychiatric morbidity, only those that were statistically significant at the 0.1 level were included in the final models. These variables included infant gender, aggregate SES, gravidity, perceived social support during pregnancy, maternal education during pregnancy, household density, alcohol consumption during pregnancy, tobacco use during pregnancy, fetal growth rate, gestational age, maternal age, recent stress, past stress, and maternal GHQ in 2007. Multiple ordinary least squares (OLS)

regressions were conducted to examine the impact of prenatal stress on late adolescent psychiatric morbidity.

The final analytical sample included 304 mothers and adolescents with requisite data (Table 1). Participants included in the analytical sample were similar to those excluded ($n = 747$) with respect to psychiatric morbidity at 17-18 years, prenatal stress, child gender, gravidity, alcohol consumption, tobacco use, social support, household density, SES in 2008, fetal growth rate, recent stress, past stress, and maternal psychiatric morbidity ($p > 0.05$). Birthweight, gestational age, SES in 1990, maternal education in 1990, and the aggregate SES measure were significantly different from those excluded from the sample ($p < 0.05$). Children in the analytical sample exhibited higher birthweights and slightly longer gestations. Additionally, participants in the analytical sample reported lower household SES in 1990, lower aggregate SES levels, but more educated mothers in 1990. Thus, the analytic sample is somewhat disadvantaged compared to excluded group.

RESULTS

In our sample of 304 mothers and adolescent pairs, the average number of traumatic events experienced within the past six months of the interview was 2.4/15 total events, while the average number of social support resources was 2.8/4 (Table 1). About 24% of the sample, or 74 adolescents, surpassed the cutoff (51/112) for high psychological morbidity based on their GHQ scores. Table 2 presents the results of the OLS regression analyses of fetal, maternal, behavioral, and environmental factors that predict psychiatric morbidity during late adolescence. The unadjusted model (Model 1) predicting GHQ scores on prenatal stress was positive although not significant ($\beta = 0.53, p = 0.169$). The effect of prenatal stress did not substantively change after adjusting for gender or maternal age (Model 2). We tested for an interaction between prenatal stress and gender, but we found no evidence for gender differences. The relationship between prenatal stress and psychiatric morbidity strengthened and approached significance ($\beta = 0.69, p = 0.066$) after including gestational variables, specifically gravidity, gestational age, and fetal growth (birthweight adjusted for gestational age) (Model 3). Greater fetal growth rates significantly

predicted greater adolescent GHQ scores ($\beta = 0.035, p = 0.037$). Adjusting for maternal pregnancy exposures (Model 4), stress measures and maternal GHQ (Model 5), socioeconomic markers (Model 6), and prenatal social support (Model 7) did not appreciably modify the relationship between prenatal stress and adolescent GHQ, but the effect of prenatal stress falls out of significance in Model 6.

Testing moderators of prenatal stress-later life psychiatric morbidity pathway

We next assessed factors that could moderate the relationship between PNS and future psychiatric morbidity in the index generation: maternal age, recent/past stress, and social support. We examined the effect of an interaction between gender and prenatal stress on psychiatric morbidity, and results showed a non-significant interaction between prenatal stress and male ($\beta = -0.46, F[1, 287] = 0.35, p = 0.556$). Model 8 reports a negative and significant interaction between maternal age and prenatal stress severity ($\beta = -0.17, F[1, 287] = 7.14, p = 0.008$), showing that the adverse psychiatric effects of prenatal stress are stronger in children with younger mothers (Figure 2).

To assess the stress-sensitization hypothesis, we evaluated the interaction between prenatal stress and both measures of household stress from the past six and twelve months. Our results reported a non-significant interaction between prenatal stress and recent household stress from the past six months ($\beta = -0.027, F[1, 287] = 0.001, p = 0.951$) and a significant interaction between prenatal stress and past household stress from the past year (Model 9: $\beta = 1.19, F[1, 287] = 4.52, p = 0.0343$) (Figure 3). The effect of prenatal stress on late adolescent and early adult psychiatric morbidity was stronger in index children living in households with greater stress and trauma exposure. Finally, to explore the possible effects of positive maternal social environments during pregnancy in ameliorating the long-term mental health impacts of prenatal stress, we evaluated the effect of an interaction between social support and prenatal stress on psychiatric morbidity. Results reported a non-significant interaction between social support and prenatal stress ($\beta = -0.092, F[1, 287] = 0.04, p = 0.836$).

DISCUSSION

In this longitudinal study of the intergenerational mental health impacts of prenatal stress in South Africa, we find that pregnant women who reported greater trauma exposure during *Apartheid* had children who exhibited greater psychiatric morbidity during late adolescence, 17-18 years after the timing of their fetal stress exposure. This relationship, however, was only significant after interacting greater prenatal stress with younger maternal age ($p = 0.008$) and greater household stress and trauma in the past year ($p = 0.0343$) and remained statistically significant after controlling for key demographic, social, and biological characteristics. In sum, these data shed light on the potential fetal origins of late adolescent mental health and the intergenerational effects of trauma from *Apartheid* in our birth cohort sample of South African mothers and children in Soweto-Johannesburg. This study is among the first to prospectively assess the long-term psychiatric impacts of prenatal stress into early adulthood in a low- and middle-income country.

The finding that greater prenatal stress predicts future psychiatric morbidity is consistent with the growing literature on the fetal origins of later life psychopathology (Abbott et al. 2018; Bosch et al. 2012; Davis et al. 2020; O'Donnell et al. 2013; Ping et al. 2020). Retrospective and prospective studies of prenatal stress show that greater maternal social adversity during pregnancy predicts elevated risks for developing psychopathologies like depression, psychosis, and schizophrenia in the future (Lipner et al. 2019; McQuaid et al. 2019; Van den Bergh et al. 2008). The long-term, intergenerational effects of prenatal stress, however, were only significant when the moderating effects of maternal age and past traumatic life events on prenatal stress were independently assessed.

Our findings on the interaction between prenatal stress and younger maternal age corroborate past research highlights the numerous socioeconomic, gendered, and cultural adversities faced by younger pregnant mothers and their families in Soweto and other communities in South Africa (Makola 2011; McLeod 1999; Richter et al. 2006; Willan 2013). Young motherhood in Soweto is understood to be a highly stigmatized and morally compromised status in Soweto and a period of greater vulnerability to certain forms of violence in Soweto and other communities in South Africa (Panday et al. 2009). Past research in Bt20+, Soweto, and elsewhere shows that young mothers frequently receive disappointment

and negative attitudes from their parents, ridicule and shame from nurses, and stigma from community members (Makola 2011; McLeod 1999; Richter et al. 2006; Willan 2013) and that infants of young teenage mothers were lighter at birth relative to neonates from older mothers (Cameron et al. 1996; Lundeen et al. 2016; Rothberg et al. 1991). Our data also showed that both younger age was associated with lower birthweight. The long-term developmental effects of lower birthweight are well-documented – lower birthweight is associated with increased risk for a wide range of adolescent and adult mental illness risk across the lifecourse (Abel et al. 2010; Barker 2004; Lærum et al. 2019; Orri et al. 2019).

Results also show that fetal growth, but not gestational age, is an important predictor of later life psychiatric morbidity in this sample. Fetal growth was positively and significantly related to late adolescent/early adulthood psychiatric morbidity. Also, our finding that fetal growth weakens the coefficient on prenatal stress after inclusion into the model provides preliminary evidence that fetal growth may be involved in the relationship between prenatal stress and adolescent psychiatric morbidity. The direct effect of fetal growth rate on psychiatric morbidity at 18, however, conflicts with past literature on the fetal origins hypothesis that suggests that adverse intrauterine exposures and slower fetal growth, potentially due to stress-induced alterations in gestational neuroendocrine activity, can durably impact health, development, and disease risk in the next generation (Kuzawa 2008; O'Donnell & Meaney 2017; Monk et al. 2019). Past studies have shown sex differences in fetal growth rates and preterm birth with boys typically showing higher incidence of adverse birth outcomes (Di Renzo et al. 2007), yet there were no significant sex differences in fetal growth rates, gestational age, nor birthweights in this sample. Further research in Bt20+ should examine the potential sex differences in fetal growth rates due to prenatal stress.

Our data show that recent stress from the past 6 months was a positive and marginally significant predictor of psychiatric morbidity at 17-18, while the effect of traumatic life experiences from the past 12 months was only significant when interacted with prenatal stress. The significant interaction between prenatal stress and past year traumatic experiences, showing that the direct relationship between prenatal stress and psychiatric morbidity is stronger in children with greater household traumatic events, provide

supporting evidence for the stress sensitization hypothesis (Hammen et al. 2000; McLaughlin et al. 2010; Van den Bergh et al. 2008). While the data show that prenatal stress itself may not have sensitizing effects on later life psychiatric morbidity, it is only after interacting with elevated levels of past stress that the long-term sensitizing effects of prenatal stress on later life psychiatric morbidity become apparent. The stress sensitizing effect of prenatal stress may only significantly interact with past stress (from the past 12 months) trauma rather than with recent stress (from the past six months) because of the nature of the experiences queried by each measure. Reported by the mother during the interview, our composite measure of past stress consisted of eight questions assessing a collection of objective traumatic events that are more likely to have household-level impacts specifically on the index child compared to the events assessed in the recent stress variable. Also reported by the mother, our measure of recent stress may also query household experiences that may have marginal to no impact on the index child's psychiatric morbidity. These experiences include unemployment, divorce, or death of a relative. Nevertheless, our data show interesting stress sensitization effects due to the interaction between prenatal stress and household stress exposure from the past year.

Given these data and existing findings on the fetal origins of late adolescent/early adult psychopathology, there are two possible developmental mechanisms that may facilitate the lasting impacts of prenatal stress on future psychiatric morbidity. First, increased severity of prenatal stress may cause durable increases in psychological and physiological stress reactivity (e.g. HPA axis, the immune system, and brain function) into adulthood, which may make individuals respond worse to future stressors, and in turn increase one's risk of developing a psychopathology (Hammen et al. 2000; Heim et al. 2019; Kendler et al. 2004; McLaughlin et al. 2010). In one study, researchers found that women who experienced greater maternal anxiety during pregnancy were more likely to have children with flattened diurnal cortisol slopes, which predicted depression in female adolescents (Van den Bergh et al. 2008). The growing literature on the long-term health effects of prenatal stress is consistent with the larger body of scholarship on the effects of postnatal early life stress, which are known to have similar lasting effects

on neuroendocrine, inflammatory, and molecular mechanisms across development, extending into adulthood (Gustafsson et al. 2010; Heim & Binder 2012; Taylor et al. 2010).

Second, greater histories of prenatal stress may increase the severity of behavioral and psychiatric conditions that increase emotional and biological sensitization to future stressors and adverse events, such as depression, anxiety, and other mood disorders. Additionally, the major symptoms of depression, such as persistent feelings of victimization, learned hopelessness and helplessness, and negative appraisal (Folkman & Lazarus 1986; Peterson & Seligman 1983) may have elevated individual sensitivity to and appraisal of recent stressful events (Medrano & Hatch 2005; Peterson & Seligman 1983) and neuroendocrine sensitization (Stroud et al. 2011) to future stressors. Thus, greater sensitivity to and appraisal of past stress may have emerged as a function of the long-term depressive and psychological effects of prenatal stress. Future longitudinal research is needed to determine the underlying stress physiological mechanisms by which prenatal stress influences future stress sensitivity and psychopathological morbidity.

Additional stress physiological mechanisms and external social processes may also explain the late adolescent psychiatric impacts of prenatal stress in our sample. Growing research report that higher levels maternal prenatal stress corresponds with greater pro-inflammatory cytokines levels and other inflammatory markers across adulthood (Bilbo & Schwarz 2009; Entringer et al. 2008; Plant et al. 2016; Slopen et al. 2015). Scientists have also found increasing evidence that greater maternal prenatal stress and social experience predicts epigenetic changes at genetic loci affecting neuroendocrine systems, neurotransmission, and neurogenesis in their children (Barker et al. 2018; Glover et al. 2018; Ostlund et al. 2016; Provenzi et al. 2020). While we cannot completely rule out the potential role of certain stress physiological mechanisms in driving the prenatal stress-late adolescence relationship without directly assessing the system (e.g. salivary cortisol, cytokines, etc.), multiple stress-sensitive biological pathways likely contribute to elevations in psychiatric risk at the same time. Finally, prenatal stress may be indicative of larger and longer-term socioeconomic patterns of mothers and children in Soweto. Substantial evidence illustrates the durable impacts of poverty and material deprivation during

pregnancy. Chronically low socioeconomic status between gestation and adulthood predicted flatter diurnal cortisol rhythms in a large sample of Filipino adults (DeSantis et al. 2015).

We also report that perceived social support during pregnancy did not buffer against the later-life effects of trauma exposure during pregnancy. Prenatal social support did not appear to have an independent protective effect on adolescent psychiatric morbidity, yet the presence of a partner during pregnancy did significant predict lower adolescent GHQ scores. The insignificant effect of prenatal social support on future psychiatric morbidity, either from a direct protective effect on adolescent mental health or buffering against the psychiatric impacts of prenatal stress, may be attributed to the limited strength of our social support measure, which primarily assessed interpersonal support and a single, dichotomous measure of group membership.

Psychologists have emphasized the importance of assessing the degree, frequency, and timing, and modes of social support, similar to the impacts of stress and trauma, when examining psychosocial and physiological impacts (Dunkel Schetter 2011; Orr 2004) Past studies report protective effects of prenatal social support on postnatal outcomes, including better infant birth outcomes (Feldman et al. 2000; Orr 2004) and lower child adiposity (Katzom et al. 2019). Prenatal social support, in the form of a presence of a partner during pregnancy, has also shown to buffer the adverse effects of prenatal social adversity to predict better cognitive and psychiatric outcomes in children (Spann et al. 2020). Finally, early evidence suggest that prenatal social support may buffer against the impacts of maternal stress and contribute to healthier cortisol regulation (Field et al. 2013; Luecken et al. 2013). Further research is necessary to identify both the protective and buffering effects of prenatal stress to ameliorate the long-term disease outcomes of intergenerational stress and trauma.

In many past published studies of the lifecourse impacts of adverse prenatal experience, the source of maternal social adversity stem from violent and oppressive conditions linked with political turmoil, war, famine, and numerous forms of social inequality (Yehuda et al. 2016; Kertes et al. 2016; Roseboom et al. 2006; Kim et al. 2020). Despite the formal end to these forms of political violence, the developmental and health consequences of trauma exposures among pregnant women can durably extend

across the lifecourse of the future child and may even impact the subsequent generation (Barbarin & Richter 2013; Kuzawa & Sweet 2009). These developmental pathways and health consequences of embodied trauma may represent mechanisms that drive mental health inequalities that disproportionately impact marginalized populations with high levels of stress and trauma exposure. Inasmuch as scientists aim to reverse the past impacts of embodied trauma and social oppression, the ongoing legacies of these violent histories and historical traumas, such as *Apartheid*, must be recognized and addressed to prevent future mental health inequities from emerging.

LIMITATIONS


Unfortunately, the timeframe of the queried stress exposure in the prenatal stress measurement does not specify the exact moment of exposure as the participants were asked to report stressors that occurred in the past six months. Women completed the antenatal stress questionnaire during their third trimester, meaning that the reported stressor could have occurred anytime between the first and third trimester. Future research will benefit from knowing the exact periods of prenatal stress and cortisol exposure to understand the potential developmental and physiological effects on the child.

CONCLUSION

In this analysis of the intergenerational mental health impacts of prenatal stress in a large birth cohort in Soweto, South Africa, our data show that pregnant women who reported greater trauma exposure during *Apartheid* had children who exhibited greater psychiatric morbidity during late adolescence and early adulthood, 17-18 years after the timing of their fetal stress exposure. This relationship, however, was only significant among children born to younger mothers and children exposed to household adversity in the past year. These findings suggest that greater prenatal stress may adversely affect adolescent mental health, have stress-sensitizing effects in children, and represent possible intergenerational effects of trauma experienced under *Apartheid* in this sample of South African mothers and children in Soweto-Johannesburg.

Figure 2


Interaction effect between prenatal stress and maternal age predicting psychiatric risk.



Note. This figure demonstrates the interaction effect between prenatal stress and maternal age predicting adolescent psychiatric risk at Year 17. The effect of prenatal stress on late adolescent psychiatric risk is much stronger among younger mothers ($\beta = -0.17$, $F[1, 287] = 7.14$, $p = 0.008$). There were significant main effects for prenatal stress and maternal age.

Figure 3

Interaction effect between prenatal stress and household stress and trauma in the past year predicting psychiatric risk.



Note. This figure demonstrates the interaction effect between prenatal stress and recent stress predicting adolescent psychiatric risk at Year 17. The effect of prenatal stress on late adolescent psychiatric risk is stronger as the degree of recent stress increases ($\beta = 1.19$, $F[1, 287] = 4.52$, $p = 0.0343$). There were no significant main effects for prenatal stress and recent stress.

TABLES AND FIGURES

Table 1. Demographic characteristics, prenatal conditions, and psychiatric morbidity

Variables	N = 304	Mean (SD)	%	Range
<i>Demographics</i>				
Gender (% male)	143		47.04	
Population Group (“Race”)				
Black	270		88.8	
Coloured	31		10.2	
Indian	3		1.0	
Maternal age (at enrollment)		25.5 (6.0)		15 – 43
Maternal educational attainment (% attended)				
No school or primary school	33		10.9	
Secondary school	244		79.7	
Professional/teaching/university	27		8.9	
Household density (people/room)		3.5 (1.7)		0.67 – 16
Household assets at 1990		3.9 (1.7)		0 – 7
Household assets at 2006		6.6 (2.4)		1 – 13
Maternal alcohol use in 1990				
Never	285		93.8	
Few times a year	11		3.8	
Once a month or more	8		2.6	
Maternal tobacco use in 1990				
Not at all	268		88.2	
Occasionally	13		4.3	
Daily	23		7.5	
Marital status				
Married/Partnered	90		29.6	
Single/Widowed	214		70.4	
Birthweight (g)		3149.2 (443.8)		1150 – 4550
Low birthweight (<2500g)	20	2262.3 (292.8)	6.6	1150 – 2485
Gestational age (weeks)		38.3 (1.2)		32 – 43
Preterm birth (<37 weeks)	17	35.5 (1.3)	5.6	32 – 36
<i>Psychological status</i>				
Maternal traumatic events in 1990		2.4 (1.9)		0 – 9
Maternal Social Support in 1990		2.8 (0.9)		0 – 4
Adolescent General Health Questionnaire (score)	74	45.2 (12.7)		28 – 96
High psychological risk (≥ 51)		63.3 (11.9)		52 – 96
Social Support		2.8 (0.9)		0 – 4
Maternal General Health Questionnaire in 2007 (score)		47.5 (13.2)		28-112

1 *Table 2. Multiple regression models of prenatal stress predicting adolescent psychiatric morbidity with covariates*

	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6	Model 7
Prenatal stress (count)	0.5 ± 0.4	0.6 ± 0.4	0.7 ± 0.4 [†]	0.7 ± 0.4 [†]	0.7 ± 0.4 [†]	0.6 ± 0.4	0.6 ± 0.4
Gender (male)		-5.9 ± 1.4***	-6.0 ± 1.4***	-6.0 ± 1.4***	-5.9 ± 1.5***	-5.9 ± 1.5***	-5.8 ± 1.5***
Maternal age (year)		-0.1 ± 0.1	-0.01 ± 0.1	-0.02 ± 0.1	0.03 ± 0.2	0.02 ± 0.2	0.04 ± 0.2
Gravidity (yes/no)			-3.1 ± 1.8 [†]	-3.1 ± 1.8 [†]	-3.3 ± 1.9 [†]	-3.9 ± 1.9*	-3.7 ± 1.9 [†]
Fetal growth ^a			0.004 ± 0.002*	0.003 ± 0.002*	0.003 ± 0.002 [†]	0.003 ± 0.002 [†]	0.003 ± 0.002 [†]
Gestational age (week)			-0.1 ± 0.6	-0.1 ± 0.6	-0.1 ± 0.6	-0.2 ± 0.6	-0.2 ± 0.6
Alcohol (frequency)				0.01 ± 0.9	0.04 ± 0.9	0.002 ± 0.9	-0.09 ± 0.9
Tobacco (frequency)				0.7 ± 1.3	0.7 ± 1.3	0.3 ± 1.4	0.1 ± 1.4
Crowding (ratio)					-0.3 ± 0.4	-0.4 ± 0.4	-0.4 ± 0.4
Recent Stress (count)					1.4 ± 0.8 [†]	1.5 ± 0.8 [†]	1.5 ± 0.8 [†]
Past Stress (count)					1.0 ± 1.2	1.1 ± 1.2	1.0 ± 1.2
Maternal GHQ (score)					-0.02 ± 0.06	-0.03 ± 0.06	-0.04 ± 0.06
Assets (count)						-0.3 ± 0.5	-0.2 ± 0.5
Maternal education (count)						-0.9 ± 0.9	-0.8 ± 0.9
Social support (count)							-0.9 ± 0.9
Intercept	43.9 ± 1.2***	50.2 ± 3.3***	54.0 ± 22.4*	52.1 ± 22.8*	53.8 ± 23.3*	59.4 ± 24.2*	62.9 ± 24.4*
Model R ²	0.0063	0.0641	0.0858	0.0866	0.0986	0.1037	0.1071

[†] $p < 0.1$ * $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$

^a Birthweight adjusted for gestational age

2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

	Model 8	Model 9
Prenatal stress (count)	4.8 ± 1.6**	0.1 ± 0.5
Prenatal stress x Maternal age	-0.2 ± 0.1**	
Prenatal stress x Past Stress		1.2 ± 0.6*
Gender (male)	-5.7 ± 1.4***	-5.7 ± 1.5***
Maternal age (year)	0.4 ± 0.2*	0.7 ± 0.2
Gravidity (yes/no)	-3.6 ± 1.9 [†]	-3.8 ± 1.9 [†]
Fetal growth ^a	0.003 ± 0.002*	0.003 ± 0.002 [†]
Gestational age (week)	-0.1 ± 0.6	-0.2 ± 0.6
Alcohol (frequency)	-0.004 ± 0.9	-3.8 ± 1.9
Tobacco (frequency)	0.1 ± 1.4	-0.1 ± 1.4
Crowding ^b (ratio)	-0.3 ± 0.4	-0.3 ± 0.4
Recent Stress (count)	1.5 ± 0.8 [†]	1.5 ± 0.8 [†]
Past Stress (count)	0.9 ± 1.2	-1.6 ± 1.7
Maternal GHQ (score)	-0.03 ± 0.1	-0.04 ± 0.06
Assets (count)	-0.3 ± 0.5	-0.1 ± 0.5
Maternal education (count)	-0.7 ± 0.9	-0.7 ± 0.9
Social support (count)	-1.1 ± 0.9	-0.9 ± 0.9
Intercept	50.9 ± 24.6*	61.7 ± 24.3*
Model R ²	0.1288	0.1210

[†]p < 0.1 *p < 0.05 **p < 0.01 ***p < 0.001 ^a Birthweight adjusted for gestational age ^b Number of household inhabitants/rooms used for sleeping

40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68

BIBLIOGRAPHY

- 69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
1. Abbott, P. W., Gumusoglu, S. B., Bittle, J., Beversdorf, D. Q., & Stevens, H. E. (2018). Prenatal stress and genetic risk: How prenatal stress interacts with genetics to alter risk for psychiatric illness. *Psychoneuroendocrinology*, 90, 9-21.
 2. Abel, K. M., Wicks, S., Susser, E. S., Dalman, C., Pedersen, M. G., Mortensen, P. B., & Webb, R. T. (2010). Birth weight, schizophrenia, and adult mental disorder: is risk confined to the smallest babies?. *Archives of general psychiatry*, 67(9), 923-930.
 3. Adjaye-Gbewonyo, K., Avendano, M., Subramanian, S. V., & Kawachi, I. (2016). Income inequality and depressive symptoms in South Africa: a longitudinal analysis of the National Income Dynamics Study. *Health & place*, 42, 37-46.
 4. Adonis, C. K. (2016). Exploring the salience of intergenerational trauma among children and grandchildren of victims of apartheid-era gross human rights violations. *Indo-Pacific Journal of Phenomenology*, 16(1-2).
 5. Alexander, N., Rosenlöcher, F., Stalder, T., Linke, J., Distler, W., Morgner, J., & Kirschbaum, C. (2012). Impact of antenatal synthetic glucocorticoid exposure on endocrine stress reactivity in term-born children. *The Journal of Clinical Endocrinology & Metabolism*, 97(10), 3538-3544.
 6. Allen, J., Balfour, R., Bell, R., & Marmot, M. (2014). Social determinants of mental health. *International review of psychiatry*, 26(4), 392-407.
 7. Barbarin, O. A., & Richter, L. M. (2013). *Mandela's children: Growing up in post-apartheid South Africa*. Routledge.
 8. Barker, D. J. P. (2004). The developmental origins of adult disease. *Journal of the American College of Nutrition*, 23(sup6), 588S-595S.
 9. Betts, K. S., Williams, G. M., Najman, J. M., & Alati, R. (2015). The relationship between maternal depressive, anxious, and stress symptoms during pregnancy and adult offspring behavioural and emotional problems. *Depression and Anxiety*, 32(2), 82-90.
 10. Bluen, S. D., & Odesnik, J. (1988). Township unrest: Development of the township life events scale. *South African Journal of Psychology*, 18(2), 50-57.
 11. Bolten, M. I., Wurmser, H., Buske-Kirschbaum, A., Papoušek, M., Pirke, K. M., & Hellhammer, D. (2011). Cortisol levels in pregnancy as a psychobiological predictor for birthweight. *Archives of women's mental health*, 14(1), 33-41.
 12. Bosch, N. M., Riese, H., Reijneveld, S. A., Bakker, M. P., Verhulst, F. C., Ormel, J., & Oldehinkel, A. J. (2012). Timing matters: Long term effects of adversities from prenatal period up to adolescence on adolescents' cortisol stress response. The TRAILS study. *Psychoneuroendocrinology*, 37(9), 1439-1447.
 13. Brand, S. R., Brennan, P. A., Newport, D. J., Smith, A. K., Weiss, T., & Stowe, Z. N. (2010). The impact of maternal childhood abuse on maternal and infant HPA axis function in the postpartum period. *Psychoneuroendocrinology*, 35(5), 686-693.
 14. Breslau, J., Gilman, S. E., Stein, B. D., Ruder, T., Gmelin, T., & Miller, E. (2017). Sex differences in recent first-onset depression in an epidemiological sample of adolescents. *Translational psychiatry*, 7(5), e1139-e1139.
 15. Burns, J. K. (2015). Poverty, inequality and a political economy of mental health. *Epidemiology and psychiatric sciences*, 24(2), 107-113.
 16. Cameron, N., Richter, L., McIntyre, J., Dhlamini, N., & Garstang, L. (1996). Progress report: Teenage pregnancy and birth outcome in Soweto. Unpublished report: University of the Witwatersrand.
 17. Cherak, S. J., Giesbrecht, G. F., Metcalfe, A., Ronksley, P. E., & Malebranche, M. E. (2018). The effect of gestational period on the association between maternal prenatal salivary cortisol and birth weight: a systematic review and meta-analysis. *Psychoneuroendocrinology*, 94, 49-62.
 18. Costello, E. J., Worthman, C., Erkanli, A., & Angold, A. (2007). Prediction from low birth weight to female adolescent depression: a test of competing hypotheses. *Archives of general psychiatry*, 64(3), 338-344.

- 120 19. D'Anna-Hernandez, K. L., Hoffman, M. C., Zerbe, G. O., Coussons-Read, M., Ross, R. G., &
121 Laudenslager, M. L. (2012). Acculturation, maternal cortisol and birth outcomes in women of
122 Mexican descent. *Psychosomatic Medicine*, 74(3), 296.
- 123 20. Davis, E. P., Hankin, B. L., Glynn, L. M., Head, K., Kim, D. J., & Sandman, C. A. (2020).
124 Prenatal maternal stress, child cortical thickness, and adolescent depressive symptoms. *Child*
125 *development*, 91(2), e432-e450.
- 126 21. de Bruijn, A. T., van Bakel, H. J., Wijnen, H., Pop, V. J., & van Baar, A. L. (2009). Prenatal
127 maternal emotional complaints are associated with cortisol responses in toddler and preschool
128 aged girls. *Developmental Psychobiology: The Journal of the International Society for*
129 *Developmental Psychobiology*, 51(7), 553-563.
- 130 22. De Mola, C. L., De França, G. V. A., de Avila Quevedo, L., & Horta, B. L. (2014). Low birth
131 weight, preterm birth and small for gestational age association with adult depression: systematic
132 review and meta-analysis. *The British Journal of Psychiatry*, 205(5), 340-347.
- 133 23. Di Renzo, G. C., Rosati, A., Sarti, R. D., Cruciani, L., & Cutuli, A. M. (2007). Does fetal sex
134 affect pregnancy outcome?. *Gender medicine*, 4(1), 19-30.
- 135 24. Diego, M. A., Field, T., Hernandez-Reif, M., Schanberg, S., Kuhn, C., & Gonzalez-Quintero, V.
136 H. (2009). Prenatal depression restricts fetal growth. *Early human development*, 85(1), 65-70.
- 137 25. Doane, L. D., Mineka, S., Zinbarg, R. E., Craske, M., Griffith, J. W., & Adam, E. K. (2013). Are
138 flatter diurnal cortisol rhythms associated with major depression and anxiety disorders in late
139 adolescence? The role of life stress and daily negative emotion. *Development and*
140 *psychopathology*, 25(3), 629-642.
- 141 26. Dunkel Schetter, C. (2011). Psychological science on pregnancy: stress processes,
142 biopsychosocial models, and emerging research issues. *Annual review of psychology*, 62, 531-
143 558.
- 144 27. Dunkle, K. L., Jewkes, R. K., Brown, H. C., Yoshihama, M., Gray, G. E., McIntyre, J. A., &
145 Harlow, S. D. (2004). Prevalence and patterns of gender-based violence and revictimization
146 among women attending antenatal clinics in Soweto, South Africa. *American journal of*
147 *epidemiology*, 160(3), 230-239.
- 148 28. Eleftheriades, M., Creatsas, G., & Nicolaides, K. (2006). Fetal growth restriction and postnatal
149 development. *Annals of the New York Academy of Sciences*, 1092(1), 319-330.
- 150 29. Entringer, S., Khumsta, R., Hellhammer, H. D., Wadhwa, D. P., Wust, S. (2009). Prenatal exposure
151 to maternal psychosocial stress and HPA axis regulation in young adults. *Hormones and*
152 *Behaviour*, 55(2), 292-298. <https://doi.org/10.1016/j.yhbeh.2008.11.006>
- 153 30. Entringer, S., Buss, C., Rasmussen, J. M., Lindsay, K., Gillen, D. L., Cooper, D. M., & Wadhwa,
154 P. D. (2017). Maternal cortisol during pregnancy and infant adiposity: a prospective investigation.
155 *The Journal of Clinical Endocrinology & Metabolism*, 102(4), 1366-1374.
- 156 31. Feldman, P. J., Dunkel-Schetter, C., Sandman, C. A., & Wadhwa, P. D. (2000). Maternal social
157 support predicts birth weight and fetal growth in human pregnancy. *Psychosomatic medicine*,
158 62(5), 715-725.
- 159 32. Field, T., Diego, M., Delgado, J., & Medina, L. (2013). Yoga and social support reduce prenatal
160 depression, anxiety and cortisol. *Journal of bodywork and movement therapies*, 17(4), 397-403.
- 161 33. Filmer, D., & Pritchett, L. (1999). The effect of household wealth on educational attainment:
162 evidence from 35 countries. *Population and development review*, 25(1), 85-120.
- 163 34. Gibbs, A., Jewkes, R., Willan, S., & Washington, L. (2018). Associations between poverty,
164 mental health and substance use, gender power, and intimate partner violence amongst young
165 (18-30) women and men in urban informal settlements in South Africa: A cross-sectional study
166 and structural equation model. *PLoS one*, 13(10), e0204956.
- 167 35. Global Burden of Disease Collaborative Network. (2017). Global Burden of Disease Study 2017.
168 Institute for Health Metrics and Evaluation (IHME). Retrieved from:
169 <https://vizhub.healthdata.org/gbd-compare/>.

- 170 36. Gluckman, P. D., & Hanson, M. A. (2004, October). Maternal constraint of fetal growth and its
171 consequences. In *Seminars in fetal and neonatal medicine* (Vol. 9, No. 5, pp. 419-425). WB
172 Saunders.
- 173 37. Goel, N., Workman, J. L., Lee, T. T., Innala, L., & Viau, V. (2011). Sex differences in the HPA
174 axis. *Comprehensive Physiology*, 4(3), 1121-1155.
- 175 38. Goldberg, D. P., & Hillier, V. F. (1979). A scaled version of the General Health Questionnaire.
176 *Psychological medicine*, 9(1), 139-145.
- 177 39. Hammen, C., Henry, R., & Daley, S. E. (2000). Depression and sensitization to stressors among
178 young women as a function of childhood adversity. *Journal of consulting and clinical psychology*,
179 68(5), 782.
- 180 40. Hankin, B. L. (2009). Development of sex differences in depressive and co-occurring anxious
181 symptoms during adolescence: Descriptive trajectories and potential explanations in a multiwave
182 prospective study. *Journal of Clinical Child & Adolescent Psychology*, 38(4), 460-472.
- 183 41. Heim, C. M., Entringer, S., & Buss, C. (2019). Translating basic research knowledge on the
184 biological embedding of early-life stress into novel approaches for the developmental
185 programming of lifelong health. *Psychoneuroendocrinology*, 105, 123-137.
- 186 42. Heim, C., Newport, D. J., Mletzko, T., Miller, A. H., & Nemeroff, C. B. (2008). The link between
187 childhood trauma and depression: insights from HPA axis studies in humans.
188 *Psychoneuroendocrinology*, 33(6), 693-710.
- 189 43. Jarcho, M. R., Slavich, G. M., Tylova-Stein, H., Wolkowitz, O. M., & Burke, H. M. (2013).
190 Dysregulated diurnal cortisol pattern is associated with glucocorticoid resistance in women with
191 major depressive disorder. *Biological psychology*, 93(1), 150-158.
- 192 44. Karlén, J., Frostell, A., Theodorsson, E., Faresjö, T., & Ludvigsson, J. (2013). Maternal influence
193 on child HPA axis: a prospective study of cortisol levels in hair. *Pediatrics*, 132(5), e1333-e1340.
- 194 45. Katzow, M., Messito, M. J., Mendelsohn, A. L., Scott, M. A., & Gross, R. S. (2019). The
195 Protective Effect of Prenatal Social Support on Infant Adiposity in the First 18 Months of Life.
196 *The Journal of pediatrics*, 209, 77-84.
- 197 46. Kertes, D. A., Kamin, H. S., Hughes, D. A., Rodney, N. C., Bhatt, S., & Mulligan, C. J. (2016).
198 Prenatal maternal stress predicts methylation of genes regulating the hypothalamic–pituitary–
199 adrenocortical system in mothers and newborns in the Democratic Republic of Congo. *Child
200 development*, 87(1), 61-72.
- 201 47. Kim, A. W. How should we study intergenerational trauma? Reflections on a 30-year birth cohort
202 study in Soweto, South Africa. *Somatosphere*.
- 203 48. Kim, A.W., Mohamed, R.S., Kuzawa, C.K., Norris, S.A. (2020). Maternal prenatal stress during
204 the first trimester and infant birthweight in Soweto, South Africa. In review.
- 205 49. Krontira, A. C., Cruceanu, C., & Binder, E. B. (2020). Glucocorticoids as Mediators of Adverse
206 Outcomes of Prenatal Stress. *Trends in Neurosciences*.
- 207 50. Kudielka, B. M., & Kirschbaum, C. (2005). Sex differences in HPA axis responses to stress: a
208 review. *Biological psychology*, 69(1), 113-132.
- 209 51. Kuzawa, C. W. (2008). The developmental origins of adult health: intergenerational inertia in
210 adaptation and disease. In *Evolution and health* (pp. 325-349). Oxford University Press.
- 211 52. Ladwig, K. H., Marten-Mittag, B., Erazo, N., & Gündel, H. (2001). Identifying somatization
212 disorder in a population-based health examination survey: psychosocial burden and gender
213 differences. *Psychosomatics*, 42(6), 511-518.
- 214 53. Lærum, A. M., Reitan, S. K., Evensen, K. A. I., Lydersen, S., Brubakk, A. M., Skranes, J., &
215 Indredavik, M. S. (2019). Psychiatric symptoms and risk factors in adults born preterm with very
216 low birthweight or born small for gestational age at term. *BMC psychiatry*, 19(1), 223.
- 217 54. Lipner, E., Murphy, S. K., & Ellman, L. M. (2019). Prenatal maternal stress and the cascade of
218 risk to schizophrenia spectrum disorders in offspring. *Current psychiatry reports*, 21(10), 99.

- 219 55. Luecken, L. J., Lin, B., Coburn, S. S., MacKinnon, D. P., Gonzales, N. A., & Crnic, K. A. (2013).
220 Prenatal stress, partner support, and infant cortisol reactivity in low-income Mexican American
221 families. *Psychoneuroendocrinology*, 38(12), 3092-3101.
- 222 56. MacKinnon, N., Kingsbury, M., Mahedy, L., Evans, J., & Colman, I. (2018). The association
223 between prenatal stress and externalizing symptoms in childhood: Evidence from the Avon
224 Longitudinal Study of Parents and Children. *Biological psychiatry*, 83(2), 100-108.
- 225 57. Macleod, C. (1999). Teenage pregnancy and its 'negative' consequences: Review of South
226 African research—Part 1. *South African journal of psychology*, 29(1), 1-7.
- 227 58. Maiello, S. (2001). On the transgenerational transmission of trauma and violence. *Psycho-*
228 *analytic Psychotherapy in South Africa*, 9(2), 13-31.
- 229 59. Makola, M. P. (2011). Teenage pregnancy: views of parents/caregivers, teenagers and teachers at
230 two high schools in Soweto, Gauteng (Doctoral dissertation).
- 231 60. Martel, M. M. (2013). Sexual selection and sex differences in the prevalence of childhood
232 externalizing and adolescent internalizing disorders. *Psychological bulletin*, 139(6), 1221.
- 233 61. Maxwell, S. D., Fineberg, A. M., Drabick, D. A., Murphy, S. K., & Ellman, L. M. (2018).
234 Maternal prenatal stress and other developmental risk factors for adolescent depression: Spotlight
235 on sex differences. *Journal of abnormal child psychology*, 46(2), 381-397.
- 236 62. McQuaid, G. A., Darcey, V. L., Avalos, M. F., Fishbein, D. H., & VanMeter, J. W. (2019).
237 Altered cortical structure and psychiatric symptom risk in adolescents exposed to maternal stress
238 in utero: A retrospective investigation. *Behavioural brain research*, 375, 112145.
- 239 63. Miller, A. H., Maletic, V., & Raison, C. L. (2009). Inflammation and its discontents: the role of
240 cytokines in the pathophysiology of major depression. *Biological psychiatry*, 65(9), 732-741.
- 241 64. Miller, G. E., Chen, E., & Parker, K. J. (2011). Psychological stress in childhood and
242 susceptibility to the chronic diseases of aging: moving toward a model of behavioural and
243 biological mechanisms. *Psychological bulletin*, 137(6), 959.
- 244 65. Moomal, H., Jackson, P. B., Stein, D. J., Herman, A., Myer, L., Seedat, S., & Madela-Mntla, E.
245 (2009). Perceived discrimination and mental health disorders: the South African Stress and
246 Health study. *South African Medical Journal*, 99(5).
- 247 66. Myer, L., Smit, J., Roux, L. L., Parker, S., Stein, D. J., & Seedat, S. (2008). Common mental
248 disorders among HIV-infected individuals in South Africa: prevalence, predictors, and validation
249 of brief psychiatric rating scales. *AIDS patient care and STDs*, 22(2), 147-158.
- 250 67. Naninck, E. F. G., Lucassen, P. J., & Bakker, J. (2011). Sex differences in adolescent depression:
251 do sex hormones determine vulnerability?. *Journal of neuroendocrinology*, 23(5), 383-392.
- 252 68. Nolen-Hoeksema, S., & Girgus, J. S. (1994). The emergence of gender differences in depression
253 during adolescence. *Psychological bulletin*, 115(3), 424.
- 254 69. Norris, S. A., Richter, L. M., & Fleetwood, S. A. (2007). Panel studies in developing countries:
255 Case analysis of sample attrition over the past 16 years within the Birth to Twenty cohort in
256 Johannesburg, South Africa. *Journal of International Development*, 19, 1-8.
- 257 70. O'Donnell, K. J., Glover, V., Holbrook, J. D., & O'Connor, T. G. (2014). Maternal prenatal
258 anxiety and child brain-derived neurotrophic factor (BDNF) genotype: effects on internalizing
259 symptoms from 4 to 15 years of age. *Development and psychopathology*, 26(4pt2), 1255-1266.
- 260 71. O'Donnell, K. J., Glover, V., Jenkins, J., Browne, D., Ben-Shlomo, Y., Golding, J., & O'Connor,
261 T. G. (2013). Prenatal maternal mood is associated with altered diurnal cortisol in adolescence.
262 *Psychoneuroendocrinology*, 38(9), 1630-1638.
- 263 72. O'Donnell, K., O'Connor, T. G., & Glover, V. (2009). Prenatal stress and neurodevelopment of
264 the child: focus on the HPA axis and role of the placenta. *Developmental neuroscience*, 31(4),
265 285-292.
- 266 73. Orr, S. T. (2004). Social support and pregnancy outcome: a review of the literature. *Clinical*
267 *obstetrics and gynecology*, 47(4), 842-855.

- 268 74. Orri, M., Gunnell, D., Richard-Devantoy, S., Bolanis, D., Boruff, J., Turecki, G., & Geoffroy, M.
269 C. (2019). In-utero and perinatal influences on suicide risk: a systematic review and meta-
270 analysis. *The Lancet Psychiatry*, 6(6), 477-492.
- 271 75. Ostlund, B. D., Conradt, E., Crowell, S. E., Tyrka, A. R., Marsit, C. J., & Lester, B. M. (2016).
272 Prenatal stress, fearfulness, and the epigenome: exploratory analysis of sex differences in DNA
273 methylation of the glucocorticoid receptor gene. *Frontiers in behavioural neuroscience*, 10, 147.
- 274 76. Panday, S., Makiwane, M., Ranchod, C., & Letsoalo, T. (2009). Teenage pregnancy in South
275 Africa - with a specific focus on school-going learners. *Child, Youth, Family and Social*
276 *Development, Human Sciences Research Council*. Pretoria: Department of Basic Education
- 277 77. Pariante, C. M., & Lightman, S. L. (2008). The HPA axis in major depression: classical theories
278 and new developments. *Trends in neurosciences*, 31(9), 464-468.
- 279 78. Patel, V., Flisher, A. J., Hetrick, S., & McGorry, P. (2007). Mental health of young people: a
280 global public-health challenge. *The Lancet*, 369(9569), 1302-1313.
- 281 79. Peltzer, K., & Pengpid, S. (2008). Sexual abuse, violence and HIV risk among adolescents in
282 South Africa. *Gender and Behaviour*, 6(1), 1462-1478.
- 283 80. Perrin, M. A., Chen, H., Sandberg, D. E., Malaspina, D., & Brown, A. S. (2007). Growth
284 trajectory during early life and risk of adult schizophrenia. *The British Journal of Psychiatry*,
285 191(6), 512-520.
- 286 81. Pine, D. S., Cohen, P., Gurley, D., Brook, J., & Ma, Y. (1998). The risk for early-adulthood
287 anxiety and depressive disorders in adolescents with anxiety and depressive disorders. *Archives*
288 *of general psychiatry*, 55(1), 56-64.
- 289 82. Ping, E. Y., Laplante, D. P., Elgbeili, G., Hiller, K. M., Brunet, A., O'Hara, M. W., & King, S.
290 (2015). Prenatal maternal stress predicts stress reactivity at 2½ years of age: The Iowa Flood
291 Study. *Psychoneuroendocrinology*, 56, 62-78.
- 292 83. Ping, E. Y., Laplante, D. P., Elgbeili, G., Jones, S. L., Brunet, A., & King, S. (2020). Disaster-
293 related prenatal maternal stress predicts HPA reactivity and psychopathology in adolescent
294 offspring: Project Ice Storm. *Psychoneuroendocrinology*, 104697.
- 295 84. Plant, D. T., Pawlby, S., Sharp, D., Zunszain, P. A., & Pariante, C. M. (2016). Prenatal maternal
296 depression is associated with offspring inflammation at 25 years: a prospective longitudinal
297 cohort study. *Translational psychiatry*, 6(11), e936-e936.
- 298 85. Quesada, A. A., Tristao, R. M., Pratesi, R., & Wolf, O. T. (2014). Hyper-responsiveness to acute
299 stress, emotional problems and poorer memory in former preterm children. *Stress*, 17(5), 389-399.
- 300 86. Rice, F., Harold, G. T., Boivin, J., Van den Bree, M., Hay, D. F., & Thapar, A. (2010). The links
301 between prenatal stress and offspring development and psychopathology: disentangling
302 environmental and inherited influences. *Psychological Medicine*, 40(2), 335-345.
- 303 87. Richter, L. M., Norris, S. A., & Ginsburg, C. (2006). The silent truth of teenage pregnancies–
304 Birth to Twenty cohort's next generation. *South African Medical Journal*, 96(2), 122.
- 305 88. Riis, J. L., Granger, D. A., Woo, H., Voegtline, K., DiPietro, J. A., & Johnson, S. B. (2019).
306 Long-term associations between prenatal maternal cortisol and child neuroendocrine-immune
307 regulation. *International Journal of Behavioral Medicine*, 1-15.
- 308 89. Rosa, M. J., Nentin, F., Bosquet Enlow, M., Hacker, M. R., Pollas, N., Coull, B., & Wright, R. J.
309 (2019). Sex-specific associations between prenatal negative life events and birth outcomes. *Stress*,
310 22(6), 647-653.
- 311 90. Rothberg, A. D., Shuenyane, E., Lits, B., & Strebel, P. M. (1991). Effect of stress on birth weight
312 in two Johannesburg populations. *South African Medical Journal*, 79(1), 35-38.
- 313 91. Ryu, H. (2019). Maternal prenatal stress and birth weight (Doctoral dissertation, KDI School).
- 314 92. Sawyer, S. M., Afifi, R. A., Bearinger, L. H., Blakemore, S. J., Dick, B., Ezech, A. C., & Patton, G.
315 C. (2012). Adolescence: a foundation for future health. *The Lancet*, 379(9826), 1630-1640.
- 316 93. Sharp, H., Hill, J., Hellier, J., & Pickles, A. (2015). Maternal antenatal anxiety, postnatal stroking
317 and emotional problems in children: outcomes predicted from pre-and postnatal programming
318 hypotheses. *Psychological medicine*, 45(2), 269-283.

- 319 94. Spann, M. N., Bansal, R., Hao, X., Rosen, T. S., & Peterson, B. S. (2020). Prenatal
320 socioeconomic status and social support are associated with neonatal brain morphology, toddler
321 language and psychiatric symptoms. *Child Neuropsychology*, 26(2), 170-188.
- 322 95. Stout SA , Espel EV , Sandman CA , Glynn LM , Davis EP . Fetalprogramming of children’s
323 obesity risk. *Psychoneuroendocrinology*. 2015;53:29–39.
- 324 96. Taylor, S. E. (2010). Mechanisms linking early life stress to adult health outcomes. *Proceedings*
325 *of the National Academy of Sciences*, 107(19), 8507-8512.
- 326 97. Van den Bergh, B. R., Van Calster, B., Smits, T., Van Huffel, S., & Lagae, L. (2008). Antenatal
327 maternal anxiety is related to HPA-axis dysregulation and self-reported depressive symptoms in
328 adolescence: a prospective study on the fetalorigins of depressed mood.
329 *Neuropsychopharmacology*, 33(3), 536-545.
- 330 98. Van den Bergh, B. R., van den Heuvel, M. I., Lahti, M., Braeken, M., de Rooij, S. R., Entringer,
331 S., Hoyer, D., Roseboom, T., Räikkönen, K., King, S. & Schwab, M. (2017). Prenatal
332 developmental origins of behaviour and mental health: The influence of maternal stress in
333 pregnancy. *Neuroscience & Biobehavioural Reviews*.
- 334 99. Vigo, D., Thornicroft, G., & Atun, R. (2016). Estimating the true global burden of mental illness.
335 *The Lancet Psychiatry*, 3(2), 171-178.
- 336 100. Vrshek-Schallhorn, S., Doane, L. D., Mineka, S., Zinbarg, R. E., Craske, M. G., & Adam,
337 E. K. (2013). The cortisol awakening response predicts major depression: predictive stability over
338 a 4-year follow-up and effect of depression history. *Psychological medicine*, 43(3), 483-493.
- 339 101. Willan, S. (2013). A review of teenage pregnancy in South Africa—experiences of
340 schooling, and knowledge and access to sexual & reproductive health services. *Partners in Sexual*
341 *Health*, 1-63.
- 342 102. Yehuda, R., Daskalakis, N. P., Bierer, L. M., Bader, H. N., Klengel, T., Holsboer, F., &
343 Binder, E. B. (2016). Holocaust exposure induced intergenerational effects on FKBP5
344 methylation. *Biological psychiatry*, 80(5), 372-380.
- 345 103. Zhang, B., & Wing, Y. K. (2006). Sex differences in insomnia: a meta-analysis. *Sleep*,
346 29(1), 85-93.
- 347 104. Zijlmans, M. A., Riksen-Walraven, J. M., & de Weerth, C. (2015). Associations between
348 maternal prenatal cortisol concentrations and child outcomes: A systematic review. *Neuroscience*
349 *& Biobehavioral Reviews*, 53, 1-24.
- 350