

Supplement Table 3. Additional Effects from Vascular Comorbidities on All-Cause Mortality Using Cox Proportional Hazards Regression Models.

	Model 1 ^a	Model 2 ^b
Retinopathy and DM status		
Both DM and retinopathy v.s. Retinopathy only	1.28 (0.86-1.88)	1.10 (0.68-1.76)
Both DM and retinopathy v.s. DM only	1.43 (0.92-2.21)	1.28 (0.82-2.01)
Retinopathy and HBP status		
Both HBP and retinopathy v.s. Retinopathy only	0.97 (0.63-1.49)	0.71 (0.44-1.14)
Both HBP and retinopathy v.s. HBP only	1.54 (1.17-2.03)	1.23 (0.92-1.63)
Retinopathy and CKD status		
Both CKD and retinopathy v.s. Retinopathy only	1.50 (0.99-2.25)	1.00 (0.66-1.54)
Both CKD and retinopathy v.s. CKD only	1.51 (1.15-1.96)	1.20 (0.92-1.56)
Retinopathy and CVD status		
Both CVD and retinopathy v.s. Retinopathy only	1.57 (1.05-2.35)	1.11 (0.69-1.78)
Both CVD and retinopathy v.s. CVD only	1.38 (0.96-1.99)	1.07 (0.70-1.64)

Abbreviations: DM, diabetes mellitus; HBP, high blood pressure; CKD, chronic kidney disease; CVD, cardiovascular disease.

Boldface indicates statistical significance. Values are hazard ratios (95% confidence interval).

All-cause mortality was assessed through December 31, 2015.

^a Model 1: Adjusted for age, gender, race, education level, marital status, income status.

^b Model 2: Model 1 plus additional adjustments for BMI, smoking status, drinking status, hypertension, diabetes mellitus, diabetes duration, hypercholesterolaemia, C-reactive protein, self-rated health status, walking disability, self-reported history of cardiovascular disease and chronic kidney disease.