

Table S1 Summary of mass spectrometry results for the three types of samples

	Peptides	PSM	Proteins	MeOx	NQ
CP	770	2827	65	126 (4.5%)	87 (3.1%)
aWF	7809	47228	373	10243 (21.7%)	3009 (6.4%)
Dressing	10789	52091	418	5059 (9.7%)	1774 (3.4%)

Total numbers of identified unique peptides, peptide spectrum matches (PSM), and unique proteins found in three citrate plasma samples, five acute wound fluids and six dressing extracts. Furthermore, total numbers and percentages of peptides with post translational modifications (MeOx: methionine oxidation; NQ: deamidation) are indicated.

Table S2 Identified unique peptides, proteins and average peptide length from five acute wound fluids

	WF1	WF2	WF3	WF4	WF5
Number of peptides	3876	4271	2932	2649	3435
Number of proteins	188	222	150	143	196
Average length (Da)	1414.70	1367.43	1463.87	1516.11	1373.36
Average number of AA	12.69	12.28	13.26	13.68	12.28

AA: amino acids

Table S3 Identified unique peptides, proteins and average peptide length from six dressing extracts

	Low inflammation			High inflammation		
	1	2	3	4	5	6
Number of peptides	3146	4863	4277	5103	1412	4929
Number of proteins	179	228	200	170	88	175
Average length (Da)	1548.67	1591.19	1546.77	1589.84	1351.38	1668.49
Average number of AA	14.33	14.52	14.13	14.44	12.59	15.29

AA: amino acids

Supplementary figure 1: Venn diagrams comparing the numbers of identified proteins and their peptides of each sample in the low (A) and high (B) inflammation group.

