

Appendix A

Results of the logistic regression model to clarify the factors related to increase in care needs of long-term care insureds in Japan in 3 years.




Fig. A.1. Flow chart of data selection; the study sample for analysis comprised 76,102 subjects.

Table A.1.

Demographic Characteristics and Chi-Square Test Result of Numbers of Stable or Improved and Deteriorated Groups.

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Sex						<0.0001
Male	10,879	13,835	24,714	56.0%	32.5%	
Female	24,410	26,978	51,388	52.5%	67.5%	
Former care needs certification result_1st judgment						<0.0001
Unknown	6,100	8,298	14,398	57.6%	18.9%	
Unqualified	718	1,095	1,813	60.4%	2.4%	
Temporary long-term care level	303	512	815	62.8%	1.1%	
Support required level 1	5,799	5,974	11,773	50.7%	15.5%	
Support required level 2	5,606	6,332	11,938	53.0%	15.7%	
Care needs level 1	6,660	8,497	15,157	56.1%	19.9%	
Care needs level 2	4,788	5,129	9,917	51.7%	13.0%	
Care needs level 3	2,759	2,249	5,008	44.9%	6.6%	
Care needs level 4	2,086	1,866	3,952	47.2%	5.2%	
Care needs level 5	470	861	1,331	64.7%	1.7%	
Former care needs certification result_2nds judgment						<0.0001
Unknown	5,781	7,926	13,707	57.8%	18.0%	
Unqualified	224	343	567	60.5%	0.7%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Temporary long-term care level	55	80	135	59.3%	0.2%	
Support required level 1	5,896	5,918	11,814	50.1%	15.5%	
Support required level 2	5,385	5,753	11,138	51.7%	14.6%	
Care needs level 1	6,741	8,257	14,998	55.1%	19.7%	
Care needs level 2	4,997	5,803	10,800	53.7%	14.2%	
Care needs level 3	3,379	3,570	6,949	51.4%	9.1%	
care needs level 4	2,389	2,043	4,432	46.1%	5.8%	
Care needs level 5	442	1,120	1,562	71.7%	2.1%	
Living arrangement						0.076
Unknown	2	0	2	0.0%	0.0%	
Home	34,332	39,788	74,120	53.7%	97.4%	
Facility	955	1,025	1,980	51.8%	2.6%	
Care Needs Level_1st judgment						<0.0001
Unqualified	16	1,799	1,815	99.1%	2.4%	
Support required level 1	5,572	8,290	13,862	59.8%	18.2%	
Support required level 2	6,518	7,467	13,985	53.4%	18.4%	
Care needs level 1	6,827	9,914	16,741	59.2%	22.0%	
Care needs level 2	6,593	6,638	13,231	50.2%	17.4%	
Care needs level 3	4,210	3,532	7,742	45.6%	10.2%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Care needs level 4	5,553	3,173	8,726	36.4%	11.5%	
Care Needs Level_2nd judgment						<0.0001
Unqualified	41	366	407	89.9%	0.5%	
Support required level 1	5,596	7,595	13,191	57.6%	17.3%	
Support required level 2	6,199	5,876	12,075	48.7%	15.9%	
Care needs level 1	7,035	10,049	17,084	58.8%	22.4%	
Care needs level 2	6,531	7,079	13,610	52.0%	17.9%	
Care needs level 3	4,389	5,433	9,822	55.3%	12.9%	
Care needs level 4	5,439	3,287	8,726	37.7%	11.5%	
Care needs level 5	59	1,128	1,187	95.0%	1.6%	
Degree of independent living for elderly with disability						<0.0001
Dependent	164	274	438	62.6%	0.6%	
J 1 (able to go out using public transportation)	3,514	3,344	6,858	48.8%	9.0%	
J 2 (able to go out but limited to the neighborhood)	6,999	8,186	15,185	53.9%	20.0%	
A 1 (able to go out with assistance and usually not bedridden)	7,195	9,927	17,122	58.0%	22.5%	
A 2 (rarely able to go out and naps several times during daytime)	8,363	11,249	19,612	57.4%	25.8%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
B 1 (able to eat and use a wheelchair to access the toilet away from bed)	3,491	3,442	6,933	49.6%	9.1%	
B 2 (requires care to get on a wheelchair)	4,222	2,729	6,951	39.3%	9.1%	
C 1 (able to roll over by him/herself)	794	612	1,406	43.5%	1.8%	
C 2 (fully assisted in rolling over by him/herself)	547	1,050	1,597	65.7%	2.1%	
Degree of independent living for elderly people with dementia						<0.0001
Dependent	9,478	9,365	18,843	49.7%	24.8%	
I	9,917	11,536	21,453	53.8%	28.2%	
II a	3,902	4,279	8,181	52.3%	10.8%	
II b	6,495	7,221	13,716	52.6%	18.0%	
III a	3,775	5,450	9,225	59.1%	12.1%	
III b	869	1,251	2,120	59.0%	2.8%	
IV	759	1,442	2,201	65.5%	2.9%	
M	94	269	363	74.1%	0.5%	
Short-term memory						<0.0001
Loss	12,432	12,864	25,296	50.9%	33.2%	
Unknown	11	3	14	21.4%	0.0%	
Independent	11,066	11,962	23,028	51.9%	30.3%	
With difficulty	11,780	15,984	27,764	57.6%	36.5%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Cognitive ability (daily decision-making)						<0.0001
Loss	12,432	12,864	25,296	50.9%	33.2%	
Unknown	13	5	18	27.8%	0.0%	
Independent	11,208	12,289	23,497	52.3%	30.9%	
Some difficulty	6,482	8,183	14,665	55.8%	19.3%	
Partially assisted	3,903	5,374	9,277	57.9%	12.2%	
Fully assisted	1,251	2,098	3,349	62.6%	4.4%	
Communication ability						<0.0001
Unknown	14	7	21	33.3%	0.0%	
Capable	13,528	15,312	28,840	53.1%	37.9%	
Somewhat challenging	5,876	7,572	13,448	56.3%	17.7%	
Limited to specific requirements	2,953	4,050	7,003	57.8%	9.2%	
Incapable	486	1,008	1,494	67.5%	2.0%	
Eating						<0.0001
Unknown	17	25	42	59.5%	0.1%	
Independent	22,326	26,937	49,263	54.7%	64.7%	
Needs to be fed	514	987	1,501	65.8%	2.0%	
BPSD						<0.0001
None	5,225	5,686	10,911	52.1%	14.3%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Yes	3,272	5,204	8,476	61.4%	11.1%	
No	14,360	17,059	31,419	54.3%	41.3%	
Turn in bed						<0.0001
Independent	11,742	17,824	29,566	60.3%	38.9%	
Can do holding railings	20,575	20,069	40,644	49.4%	53.4%	
Fully assisted	2,972	2,920	5,892	49.6%	7.7%	
Get up						<0.0001
Independent	2,738	4,963	7,701	64.4%	10.1%	
Can do holding railings	28,633	32,345	60,978	53.0%	80.1%	
Fully assisted	3,918	3,505	7,423	47.2%	9.8%	
Maintain sitting position						<0.0001
Independent	12,519	17,827	30,346	58.7%	39.9%	
Can do holding railings	13,864	15,032	28,896	52.0%	38.0%	
Partially assisted	8,626	7,312	15,938	45.9%	20.9%	
Fully assisted	280	642	922	69.6%	1.2%	
Stand on both feet						<0.0001
Independent	15,376	21,239	36,615	58.0%	48.1%	
Partially assisted	16,690	16,636	33,326	49.9%	43.8%	
Fully assisted	3,223	2,938	6,161	47.7%	8.1%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Walk						<0.0001
Independent	8,718	12,402	21,120	58.7%	27.8%	
Can do with assistive technology	19,301	22,299	41,600	53.6%	54.7%	
Fully assisted	7,270	6,112	13,382	45.7%	17.6%	
Transfer						<0.0001
Independent	20,631	27,936	48,567	57.5%	63.8%	
Needs supervision	6,700	7,025	13,725	51.2%	18.0%	
Partially assisted	5,514	3,523	9,037	39.0%	11.9%	
Fully assisted	2,444	2,329	4,773	48.8%	6.3%	
Stand up						<0.0001
Independent	1,027	2,955	3,982	74.2%	5.2%	
Can do holding railings	30,411	34,545	64,956	53.2%	85.4%	
Fully assisted	3,851	3,313	7,164	46.2%	9.4%	
Stand on one foot						<0.0001
Independent	1,909	3,369	5,278	63.8%	6.9%	
Partially assisted	21,312	26,671	47,983	55.6%	63.1%	
Fully assisted	12,068	10,773	22,841	47.2%	30.0%	
Bathing						<0.0001
Independent	13,659	16,599	30,258	54.9%	39.8%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Partially assisted	11,511	13,982	25,493	54.8%	33.5%	
Fully assisted	7,984	7,793	15,777	49.4%	20.7%	
Does not take bath	2,135	2,439	4,574	53.3%	6.0%	
Swallowing						<0.0001
Independent	26,431	32,093	58,524	54.8%	76.9%	
Needs supervision	8,635	7,888	16,523	47.7%	21.7%	
Fully assisted	223	832	1,055	78.9%	1.4%	
Eating						<0.0001
Independent	28,818	33,982	62,800	54.1%	82.5%	
Needs supervision	4,146	4,439	8,585	51.7%	11.3%	
Partially assisted	1,969	1,394	3,363	41.5%	4.4%	
Fully assisted	356	998	1,354	73.7%	1.8%	
Oral hygiene						<0.0001
Independent	23,642	28,750	52,392	54.9%	68.8%	
Partially assisted	9,545	9,580	19,125	50.1%	25.1%	
Fully assisted	2,102	2,483	4,585	54.2%	6.0%	
Washing face						<0.0001
Independent	23,568	29,054	52,622	55.2%	69.1%	
Partially assisted	9,623	9,292	18,915	49.1%	24.9%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Fully assisted	2,098	2,467	4,565	54.0%	6.0%	
Hair care						<0.0001
Independent	25,800	31,066	56,866	54.6%	74.7%	
Partially assisted	6,004	6,057	12,061	50.2%	15.8%	
Fully assisted	3,485	3,690	7,175	51.4%	9.4%	
Nail-cutting						<0.0001
Independent	13,044	16,303	29,347	55.6%	38.6%	
Partially assisted	8,910	10,085	18,995	53.1%	25.0%	
Fully assisted	13,335	14,425	27,760	52.0%	36.5%	
Putting on/removing top						<0.0001
Independent	19,826	24,869	44,695	55.6%	58.7%	
Needs supervision	2,795	4,673	7,468	62.6%	9.8%	
Partially assisted	9,783	8,444	18,227	46.3%	24.0%	
Fully assisted	2,885	2,827	5,712	49.5%	7.5%	
Putting on/removing pants						<0.0001
Independent	18,836	24,133	42,969	56.2%	56.5%	
Needs supervision	2,739	4,487	7,226	62.1%	9.5%	
Partially assisted	8,805	8,109	16,914	47.9%	22.2%	
Fully assisted	4,909	4,084	8,993	45.4%	11.8%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Taking medication						0.048
Independent	14,734	16,962	31,696	53.5%	41.6%	
Partially assisted	15,489	18,206	33,695	54.0%	44.3%	
Fully assisted	5,066	5,645	10,711	52.7%	14.1%	
Managing money						<0.0001
Independent	12,791	14,261	27,052	52.7%	35.5%	
Partially assisted	7,320	8,957	16,277	55.0%	21.4%	
Fully assisted	15,178	17,595	32,773	53.7%	43.1%	
Vision						<0.0001
Normal	23,788	28,364	52,152	54.4%	68.5%	
Able to see over 1m away	9,180	9,669	18,849	51.3%	24.8%	
Nearsightedness	1,726	1,794	3,520	51.0%	4.6%	
Poor sight	434	472	906	52.1%	1.2%	
Cannot be judged	161	514	675	76.1%	0.9%	
Hearing ability						<0.0001
Normal	17,935	20,977	38,912	53.9%	51.1%	
Manages to hear	10,771	11,823	22,594	52.3%	29.7%	
Can hear loud sounds	6,039	7,132	13,171	54.1%	17.3%	
Poor hearing	459	555	1,014	54.7%	1.3%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Cannot be judged	85	326	411	79.3%	0.5%	
Expressing intentions						<0.0001
Capable	27,708	30,171	57,879	52.1%	76.1%	
Sometimes capable	5,722	7,852	13,574	57.8%	17.8%	
Almost incapable	1,616	2,116	3,732	56.7%	4.9%	
Incapable	243	674	917	73.5%	1.2%	
Understanding the daily routine						<0.0001
Capable	26,935	29,239	56,174	52.1%	73.8%	
Incapable	8,354	11,574	19,928	58.1%	26.2%	
Remembering birth date						<0.0001
Capable	32,890	36,972	69,862	52.9%	91.8%	
Incapable	2,399	3,841	6,240	61.6%	8.2%	
Short-term memory						<0.0001
Capable	23,907	26,807	50,714	52.9%	66.6%	
Incapable	11,382	14,006	25,388	55.2%	33.4%	
Remembering one's name						<0.0001
Capable	34,665	39,516	74,181	53.3%	97.5%	
Incapable	624	1,297	1,921	67.5%	2.5%	
current season						<0.0001

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Capable	29,374	32,331	61,705	52.4%	81.1%	
Incapable	5,915	8,482	14,397	58.9%	18.9%	
Recognize places						<0.0001
Capable	32,065	35,762	67,827	52.7%	89.1%	
Incapable	3,224	5,051	8,275	61.0%	10.9%	
Conviction of being threatened						<0.0001
Absent	31,412	35,271	66,683	52.9%	87.6%	
Sometimes present	1,835	2,366	4,201	56.3%	5.5%	
Present	2,042	3,176	5,218	60.9%	6.9%	
Delusional thoughts and imaginary conversation						<0.0001
Absent	29,704	33,281	62,985	52.8%	82.8%	
Sometimes present	2,308	2,887	5,195	55.6%	6.8%	
Present	3,277	4,645	7,922	58.6%	10.4%	
Emotional instability						<0.0001
Absent	26,978	30,829	57,807	53.3%	76.0%	
Sometimes present	3,948	4,568	8,516	53.6%	11.2%	
Present	4,363	5,416	9,779	55.4%	12.8%	
Day–night reversal						<0.0001
Absent	30,455	34,828	65,283	53.3%	85.8%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Sometimes present	2,440	2,936	5,376	54.6%	7.1%	
Present	2,394	3,049	5,443	56.0%	7.2%	
Speech repetition						<0.0001
Absent	23,960	26,976	50,936	53.0%	66.9%	
Sometimes present	3,653	4,102	7,755	52.9%	10.2%	
Present	7,676	9,735	17,411	55.9%	22.9%	
Screaming/Calling out						<0.0001
Absent	31,584	35,951	67,535	53.2%	88.7%	
Sometimes present	1,894	2,383	4,277	55.7%	5.6%	
Present	1,811	2,479	4,290	57.8%	5.6%	
Resistance to care						<0.0001
Absent	30,242	33,641	63,883	52.7%	83.9%	
Sometimes present	2,469	3,177	5,646	56.3%	7.4%	
Present	2,578	3,995	6,573	60.8%	8.6%	
Frequent wandering						<0.0001
Absent	34,046	38,298	72,344	52.9%	95.1%	
Sometimes present	479	887	1,366	64.9%	1.8%	
Present	764	1,628	2,392	68.1%	3.1%	
Restlessness						<0.0001

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Absent	33,514	37,996	71,510	53.1%	94.0%	
Sometimes present	818	1,160	1,978	58.6%	2.6%	
Present	957	1,657	2,614	63.4%	3.4%	
Not returning after going out						<0.0001
Absent	34,244	38,805	73,049	53.1%	96.0%	
Sometimes present	446	944	1,390	67.9%	1.8%	
Present	599	1,064	1,663	64.0%	2.2%	
Insistence on going out alone						<0.0001
Absent	33,970	38,439	72,409	53.1%	95.1%	
Sometimes present	625	1,035	1,660	62.3%	2.2%	
Present	694	1,339	2,033	65.9%	2.7%	
Hoarding behavior						<0.0001
Absent	34,130	38,892	73,022	53.3%	96.0%	
Sometimes present	350	587	937	62.6%	1.2%	
Present	809	1,334	2,143	62.2%	2.8%	
Destroying objects and clothing						<0.0001
Absent	34,839	40,015	74,854	53.5%	98.4%	
Sometimes present	234	412	646	63.8%	0.8%	
Present	216	386	602	64.1%	0.8%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Movement						<0.0001
Independent	17,281	23,284	40,565	57.4%	53.3%	
Needs supervision	7,892	9,782	17,674	55.3%	23.2%	
Partial assistance	5,667	4,286	9,953	43.1%	13.1%	
Fully assisted	4,449	3,461	7,910	43.8%	10.4%	
Passing urine/Urination						<0.0001
Independent	20,617	25,710	46,327	55.5%	60.9%	
Needs supervision	2,192	3,212	5,404	59.4%	7.1%	
Partial assistance	7,428	7,679	15,107	50.8%	19.9%	
Fully assisted	5,052	4,212	9,264	45.5%	12.2%	
Passing motion/Defecation						<0.0001
Independent	21,658	27,468	49,126	55.9%	64.6%	
Needs supervision	1,975	3,029	5,004	60.5%	6.6%	
Partial assistance	6,465	6,017	12,482	48.2%	16.4%	
Fully assisted	5,191	4,299	9,490	45.3%	12.5%	
Everyday decision-making						<0.0001
Capable	12,221	13,789	26,010	53.0%	34.2%	
Capable except during special occasions	16,855	18,017	34,872	51.7%	45.8%	
Capable but very challenging	5,314	7,423	12,737	58.3%	16.7%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Incapable	899	1,584	2,483	63.8%	3.3%	
Forgetfulness						<0.0001
Absent	18,446	19,587	38,033	51.5%	50.0%	
Sometimes present	6,553	7,699	14,252	54.0%	18.7%	
Present	10,290	13,527	23,817	56.8%	31.3%	
Frequency of going out						<0.0001
More than once a week	19,802	22,323	42,125	53.0%	55.4%	
More than once a month	6,039	7,625	13,664	55.8%	18.0%	
Less than once a month	9,448	10,865	20,313	53.5%	26.7%	
Self-talk/Laughing to oneself						<0.0001
Absent	32,938	37,637	70,575	53.3%	92.7%	
Sometimes present	880	1,102	1,982	55.6%	2.6%	
Present	1,471	2,074	3,545	58.5%	4.7%	
Selfish behaviors						<0.0001
Absent	31,122	35,095	66,217	53.0%	87.0%	
Sometimes present	1,918	2,379	4,297	55.4%	5.6%	
Present	2,249	3,339	5,588	59.8%	7.3%	
Disorganized talking						<0.0001
Absent	27,869	31,039	58,908	52.7%	77.4%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Sometimes present	3,322	4,069	7,391	55.1%	9.7%	
Present	4,098	5,705	9,803	58.2%	12.9%	
Not fitting in a group						<0.0001
Absent	32,789	37,536	70,325	53.4%	92.4%	
Sometimes present	1,160	1,501	2,661	56.4%	3.5%	
Present	1,340	1,776	3,116	57.0%	4.1%	
Shopping						<0.0001
Independent	5,586	5,549	11,135	49.8%	14.6%	
Needs supervision	369	385	754	51.1%	1.0%	
Partial assistance	7,777	9,245	17,022	54.3%	22.4%	
Fully assisted	21,557	25,634	47,191	54.3%	62.0%	
Cooking						<0.0001
Independent	9,788	10,502	20,290	51.8%	26.7%	
Needs supervision	527	679	1,206	56.3%	1.6%	
Partial assistance	3,278	4,241	7,519	56.4%	9.9%	
Fully assisted	21,696	25,391	47,087	53.9%	61.9%	
BPSD_visual/auditory hallucination						<0.0001
No	34,530	39,642	74,172	53.4%	97.5%	
Yes	759	1,171	1,930	60.7%	2.5%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
BPSD_ Delusions						<0.0001
No	33,942	38,683	72,625	53.3%	95.4%	
Yes	1,347	2,130	3,477	61.3%	4.6%	
BPSD_ Day-night reversal						<0.0001
No	34,586	39,671	74,257	53.4%	97.6%	
Yes	703	1,142	1,845	61.9%	2.4%	
BPSD_ Violent language						<0.0001
No	34,743	39,828	74,571	53.4%	98.0%	
Yes	546	985	1,531	64.3%	2.0%	
BPSD_ Assaulting						<0.0001
No	35,114	40,480	75,594	53.5%	99.3%	
Yes	175	333	508	65.6%	0.7%	
BPSD_ Refusal to care						<0.0001
No	34,496	39,444	73,940	53.3%	97.2%	
Yes	793	1,369	2,162	63.3%	2.8%	
BPSD_ Wandering						<0.0001
No	34,823	39,891	74,714	53.4%	98.2%	
Yes	466	922	1,388	66.4%	1.8%	
BPSD_ Mismanagement of fire						<0.0001

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
No	34,915	39,961	74,876	53.4%	98.4%	
Yes	374	852	1,226	69.5%	1.6%	
BPSD_ Filthy behaviors						<0.0001
No	35,028	40,359	75,387	53.5%	99.1%	
Yes	261	454	715	63.5%	0.9%	
BPSD_ Pica behaviors						0.084
No	35,227	40,718	75,945	53.6%	99.8%	
Yes	62	95	157	60.5%	0.2%	
BPSD_ Sexual disinhibition						0.792
No	35,274	40,794	76,068	53.6%	100.0%	
Yes	15	19	34	55.9%	0.0%	
BPSD_ Others						<0.0001
No	34,576	39,813	74,389	53.5%	97.7%	
Yes	713	1,000	1,713	58.4%	2.3%	
No paralysis in limbs						<0.0001
No	24,377	26,676	51,053	52.3%	67.1%	
Yes	10,912	14,137	25,049	56.4%	32.9%	
Paralysis in left upper limb						<0.0001
No	30,832	36,628	67,460	54.3%	88.6%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Yes	4,457	4,185	8,642	48.4%	11.4%	
Paralysis in right upper limb						<0.0001
No	30,363	36,113	66,476	54.3%	87.4%	
Yes	4,926	4,700	9,626	48.8%	12.6%	
Paralysis in left lower limb						<0.0001
No	14,736	17,590	32,326	54.4%	42.5%	
Yes	20,553	23,223	43,776	53.0%	57.5%	
Paralysis in right lower limb						0.014
No	14,950	17,652	32,602	54.1%	42.8%	
Yes	20,339	23,161	43,500	53.2%	57.2%	
Paralysis outside limbs						<0.0001
No	31,478	37,246	68,724	54.2%	90.3%	
Yes	3,811	3,567	7,378	48.3%	9.7%	
Joint movement without restriction						<0.0001
No	19,301	19,050	38,351	49.7%	50.4%	
Yes	15,988	21,763	37,751	57.6%	49.6%	
Shoulder joint movement restriction						<0.0001
No	29,180	35,623	64,803	55.0%	85.2%	
Yes	6,109	5,190	11,299	45.9%	14.8%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Hip joint movement restriction						<0.0001
No	31,488	37,388	68,876	54.3%	90.5%	
Yes	3,801	3,425	7,226	47.4%	9.5%	
Knee joint movement restriction						<0.0001
No	24,019	29,199	53,218	54.9%	69.9%	
Yes	11,270	11,614	22,884	50.8%	30.1%	
Other joint movement restriction						<0.0001
No	26,891	32,369	59,260	54.6%	77.9%	
Yes	8,398	8,444	16,842	50.1%	22.1%	
Medical treatment in the last 2 weeks_Intravenous therapy						<0.0001
No	34,348	40,002	74,350	53.8%	97.7%	
Yes	941	811	1,752	46.3%	2.3%	
Medical treatment in the last 2 weeks_Total parenteral nutrition						0.001
No	35,252	40,731	75,983	53.6%	99.8%	
Yes	37	82	119	68.9%	0.2%	
Medical treatment in the last 2 weeks_Dialysis						<0.0001
No	34,840	40,410	75,250	53.7%	98.9%	
Yes	449	403	852	47.3%	1.1%	
Medical treatment in the last 2 weeks_Stoma						

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
No	35,070	40,631	75,701	53.7%	99.5%	
Yes	219	182	401	45.4%	0.5%	
Medical treatment in the last 2 weeks_Oxygen therapy						0.001
No	34,683	39,982	74,665	53.5%	98.1%	
Yes	606	831	1,437	57.8%	1.9%	
Medical treatment in the last 2 weeks_Artificial respirator						0.345
No	35,273	40,788	76,061	53.6%	99.9%	
Yes	16	25	41	61.0%	0.1%	
Medical treatment in the last 2 weeks_Tracheostomy						<0.0001
No	35,262	40,745	76,007	53.6%	99.9%	
Yes	27	68	95	71.6%	0.1%	
Medical treatment in the last 2 weeks_Pain care						0.009
No	35,065	40,488	75,553	53.6%	99.3%	
Yes	224	325	549	59.2%	0.7%	
Medical treatment in the last 2 weeks_Tube feeding						<0.0001
No	35,108	40,151	75,259	53.4%	98.9%	
Yes	181	662	843	78.5%	1.1%	
Medical treatment in the last 2 weeks_Vital sign monitor						<0.0001
No	35,020	40,619	75,639	53.7%	99.4%	

Variables	Stable or Improved (care needs remained the same or decreased) n = 35,289	Deteriorated (care needs increased) n = 40,813	Total n = 76,102	Deterioration Proportion (%)	% in each category	P value*
Yes	269	194	463	41.9%	0.6%	
Medical treatment in the last 2 weeks_Bedsore treatment						0.001
No	34,960	40,521	75,481	53.7%	99.2%	
Yes	329	292	621	47.0%	0.8%	
Medical treatment in the last 2 weeks_Catheter						<0.0001
No	34,600	40,301	74,901	53.8%	98.4%	
Yes	689	512	1,201	42.6%	1.6%	

* p values are two-tailed and are considered to be statistically significant when < 0.005 .

Results are after conducting a descriptive analysis of the presence or absence of care needs increases of the total study sample and the sub subgroup. We also conducted a single regression model to identify variables that were statistically significant with p value < 0.05 and area under ROC curve ≥ 0.53 . None of the variables extracted from the long-term care insurance claim data were statistically significant. We used the selected variables to establish our multivariable binary logistic regression model.

Table A.2. Results of Multivariable Logistic Regression Model of All Care Needs Increases (AUC:0.7143).

Variables	Odds Ratio (95% Confidence Interval)	P Value
Sex		
Female	Ref	
Male	1.22 (1.17–1.26)	<0.05
Care needs level of 1st judgment		
Care needs level 1	Ref	
Unqualified	187 (114–308)	<0.05
Support required Level 1	2.78 (2.62–2.95)	<0.05
Support required Level 2	1.66 (1.58–1.76)	<0.05
Care needs level 2	0.3 (0.28–0.32)	<0.05
Care needs level 3	0.11 (0.1–0.13)	<0.05
Care needs level 4	0.03 (0.02–0.03)	<0.05
Degree of independent living for elderly with disability		
A 2 (rarely goes out and naps several times during daytime)	Ref	
Dependent	0.38 (0.30–0.48)	<0.05
J 1 (able to go out using public transportation)	0.41 (0.38–0.44)	<0.05
J 2 (able to go out but only within the neighborhood)	0.57 (0.54–0.60)	<0.05
A 1 (goes out with assistance and usually not bedridden)	0.84 (0.81–0.88)	<0.05
B 1 (able to eat and use toilet away from bed using wheel chair)	1.14 (1.06–1.21)	<0.05
B 2 (requires assistance to get onto a wheelchair)	1.17 (1.07–1.29)	<0.05
C 1 (able to roll over by him/herself)	1.66 (1.43–1.93)	<0.05
C 2 (fully assisted in rolling over by him/herself)	1.75 (1.45–2.12)	<0.05
Cognitive ability (daily decision-making)		
Loss	Ref	
Unknown	0.30 (0.04–2.50)	0.268
Independent	1.07 (1.03–1.11)	<0.05
Some difficulty	1.36 (1.28–1.45)	<0.05
Partially assisted	1.49 (1.36–1.63)	<0.05
Fully assisted	1.48 (1.29–1.70)	<0.05
Communication ability		
Capable	Ref	
Unknown	1.16 (0.18–7.65)	0.878

Variables	Odds Ratio (95% Confidence Interval)	P Value
Somewhat challenging	0.99 (0.93–1.06)	0.743
Limited to specific requirements	0.99 (0.89–1.09)	0.792
Incapable	1.03 (0.86–1.23)	0.747
Turn in bed		
Can do holding railings	Ref	
Independent	1.11 (1.07–1.16)	<0.05
Fully assisted	1.10 (1.02–1.18)	<0.05
Get up		
Can do holding railings	Ref	
Independent	1.06 (1.00–1.13)	0.052
Fully assisted	1.36 (1.25–1.47)	<0.05
Stand on both feet		
Independent	Ref	
Partially assisted	1.11 (1.06–1.16)	<0.05
Fully assisted	1.11 (1.00–1.25)	0.057
Walk		
Can do with assistive technology	Ref	
Independent	0.87 (0.83–0.92)	<0.05
Fully assisted	1.08 (1.01–1.16)	<0.05
Transfer		
Independent	Ref	
Needs supervision	0.97 (0.92–1.03)	0.328
Partially assisted	1.04 (0.96–1.13)	0.367
Fully assisted	0.98 (0.85–1.14)	0.835
Stand up		
Can do holding railings	Ref	
Independent	1.39 (1.26–1.53)	<0.05
Fully assisted	1.14 (1.03–1.26)	<0.05
Stand on one foot		
Partially assisted	Ref	
Independent	1.03 (0.95–1.11)	0.505
Fully assisted	1.06 (1.01–1.11)	<0.05
Swallowing		

Variables	Odds Ratio (95% Confidence Interval)	P Value
Independent	Ref	
Needs supervision	0.96 (0.92–1.00)	<0.05
Fully assisted	3.10 (2.48–3.88)	<0.05
Washing face		
Independent	Ref	
Partially assisted	1.11 (1.06–1.18)	<0.05
Fully assisted	1.32 (1.19–1.46)	<0.05
Putting on/removing top		
Independent	Ref	
Needs supervision	1.30 (1.19–1.43)	<0.05
Partially assisted	1.44 (1.33–1.55)	<0.05
Fully assisted	1.65 (1.46–1.87)	<0.05
Putting on/removing pants		
Independent	Ref	
Needs supervision	0.97 (0.88–1.06)	0.487
Partially assisted	0.99 (0.91–1.07)	0.812
Fully assisted	1.08 (0.96–1.22)	0.21
Expressing intentions		
Capable	Ref	
Sometimes capable	1.26 (1.19–1.32)	<0.05
Almost incapable	1.40 (1.27–1.54)	<0.05
Incapable	1.78 (1.45–2.18)	<0.05
Understanding the daily routine		
Capable	Ref	
Incapable	1.29 (1.22–1.36)	<0.05
Remembering birth date		
Capable	Ref	
Incapable	1.18 (1.11–1.25)	<0.05
Recognize places		
Capable	Ref	
Incapable	1.21 (1.12–1.30)	<0.05
Conviction of being threatened		
Absent	Ref	

Variables	Odds Ratio (95% Confidence Interval)	P Value
Sometimes absent	1.27 (1.19–1.35)	<0.05
Present	1.45 (1.36–1.54)	<0.05
Movement		
Independent	Ref	
Needs supervision	1.13 (1.07–1.19)	<0.05
Partial assistance	1.27 (1.18–1.37)	<0.05
Fully assisted	1.15 (1.03–1.28)	<0.05
Pass motion/Defecation		
Independent	Ref	
Needs supervision	1.32 (1.23–1.42)	<0.05
Partial assistance	1.64 (1.53–1.76)	<0.05
Fully assisted	1.78 (1.60–1.97)	<0.05
Disorganized talking		
Absent	Ref	
Sometimes absent	1.25 (1.19–1.32)	<0.05
Present	1.25 (1.18–1.32)	<0.05
No paralysis in limbs		
No	Ref	
Yes	0.85 (0.81–0.88)	<0.05
Joint movement without restriction		
No	Ref	
Yes	1.19 (1.15–1.23)	<0.05
Shoulder joint movement restriction		
No	Ref	
Yes	0.98 (0.94–1.03)	0.541
Care needs certification reference time_Dementia addition (minute)	1.01 (1.01–1.01)	<0.05