

## Appendix B

Outcomes of the machine learning models with four care needs level (CNL) categories (declined needs and benefit loss as one group, support required levels 1 and 2 into one group, CNL 1–3 as one group, and CNL 4–5 to another group) and nine CNL categories (the original CNL, benefit loss, and declined needs).

Table B.1. Labels of Classification Models.

# of labels	Label description
4	0: Declined needs; Loss of benefits 1: Support required level 1–2 2: Care needs level 1–3 3: Care needs level 4–5
9	00: Declined needs 10: Loss of benefits 12: Support required level 1 13: Support required level 2 21: Care needs level 1 22: Care needs level 2 23: Care needs level 3 24: Care needs level 4 25: Care needs level 5

Table B.2. Demographic Characteristics and Chi-square or Fisher's Exact Test Result of Numbers of Stable or Improved and Deteriorated Groups.

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
Living arrangement	Unknown	0	1	0	1	0.00%	0.00%	<0.001
	Home		17,785	11,370	29,155	39.00%	95.90%	
	facility		873	358	1,231	29.10%	4.10%	
Signs of elderly people with dementia	None	0	18,659	11,728	30,387	38.60%	100.00%	1
Applicant type	1st insured	0	18,658	11,728	30,386	38.60%	100.00%	1
	2nd insured		1		1	0.00%	0.00%	
Sex	Male	0	5,993	2,517	8,510	29.60%	28.00%	<0.001
	Female		12,666	9,211	21,877	42.10%	72.00%	
Care needs level_1st judgment	Unqualified	0	223	207	430	48.10%	1.40%	<0.001
	Support required level 1		2,315	2,106	4,421	47.60%	14.50%	
	Support required level 2		2,464	2,487	4,951	50.20%	16.30%	
	Care needs level 1		2,973	2,459	5,432	45.30%	17.90%	
	Care needs level 2		3,137	2,036	5,173	39.40%	17.00%	
	Care needs level 3		2,025	1,071	3,096	34.60%	10.20%	
	Care needs level 4		2,722	890	3,612	24.60%	11.90%	
	Care needs level 5		2,800	472	3,272	14.40%	10.80%	

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
Care needs level_2nd judgment	Unqualified	0	26	35	61	57.40%	0.20%	<0.001
	Support required level 1		2046	1,892	3,938	48.00%	13.00%	
	Support required level 2		2,115	2,131	4,246	50.20%	14.00%	
	Care needs level 1		2,921	2,628	5,549	47.40%	18.30%	
	Care needs level 2		3,068	2,116	5,184	40.80%	17.10%	
	Care needs level 3		2,552	1,454	4,006	36.30%	13.20%	
	Care needs level 4		2,722	977	3,699	26.40%	12.20%	
	Care needs level 5		3,209	495	3,704	13.40%	12.20%	
Year of certification criteria adopted	2009	0	8,172	3,565	11,737	30.40%	38.60%	<0.001
	2012		10,487	8,163	18,650	43.80%	61.40%	
Turn in bed	Independent	0	5,158	4,071	9,229	44.10%	30.40%	<0.001
	Can do holding railings		9,604	6,609	16,213	40.80%	53.40%	
	Fully assisted		3,897	1,048	4,945	21.20%	16.30%	
Get up	Independent	0	1,233	947	2,180	43.40%	7.20%	<0.001
	Can do holding railings		12,765	9,659	22,424	43.10%	73.80%	
	Fully assisted		4,661	1,122	5,783	19.40%	19.00%	
Maintain sitting position	Independent	0	5,270	4,387	9,657	45.40%	31.80%	<0.001
	Can do holding railings		6,457	4,756	11,213	42.40%	36.90%	

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
	Partially assisted		6,024	2,470	8,494	29.10%	28.00%	
	Fully assisted		908	115	1,023	11.20%	3.40%	
Sitting, with feet dangling	Independent	0	18,659	11,728	30,387	38.60%	100.00%	1
Stand on both feet	Independent	0	6,378	5,288	11,666	45.30%	38.40%	<0.001
	Partially assisted		8,025	5,464	13,489	40.50%	44.40%	
	Fully assisted		4,256	976	5,232	18.70%	17.20%	
Walk	Independent	0	3,409	2,614	6,023	43.40%	19.80%	<0.001
	Can do with assistive technology		8,997	7,233	16,230	44.60%	53.40%	
	Fully assisted		6,253	1,881	8,134	23.10%	26.80%	
Transfer	Independent	0	8,722	7,480	16,202	46.20%	53.30%	<0.001
	Needs supervision		3,177	2,099	5,276	39.80%	17.40%	
	Partially assisted		2,702	1,278	3,980	32.10%	13.10%	
	Fully assisted		4,058	871	4,929	17.70%	16.20%	
Stand up	Independent	0	485	400	885	45.20%	2.90%	<0.001
	Can do holding railings		13,465	10,181	23,646	43.10%	77.80%	
	Fully assisted		4,709	1,147	5,856	19.60%	19.30%	
Stand on one foot	Independent	0	613	459	1,072	42.80%	3.50%	<0.001
	Can do holding railings		9,228	7,502	16,730	44.80%	55.10%	

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
	Fully assisted		8,818	3,767	12,585	29.90%	41.40%	
Stepping into a bathtub	Independent	0	18,659	11,728	30,387	38.60%	100.00%	1
Bathing	Independent	0	5,112	4,728	9,840	48.00%	32.40%	<0.001
	Partially assisted		5,393	4,119	9,512	43.30%	31.30%	
	Fully assisted		6,973	2,499	9,472	26.40%	31.20%	
	Does not take bath		1,181	382	1,563	24.40%	5.10%	
Bedsores	Present	0	18,659	11,728	30,387	38.60%	100.00%	1
Skin disease	Present	0	18,659	11,728	30,387	38.60%	100.00%	1
Raising an arm till chest level	Capable	0	18,659	11,728	30,387	38.60%	100.00%	1
Swallowing	Independent	0	11,863	8,887	20,750	42.80%	68.30%	<0.001
	Needs supervision		5,982	2,760	8,742	31.60%	28.80%	
	Fully assisted		814	81	895	9.10%	2.90%	
Urination urge	Present	0	18,659	11,728	30,387	38.60%	100.00%	1
Bowel movement urge	Present	0	18,659	11,728	30,387	38.60%	100.00%	1
Cleaning after urination	Independent	0	18,659	11,728	30,387	38.60%	100.00%	1
Cleaning after bowel movement	Independent	0	18,659	11,728	30,387	38.60%	100.00%	1
Eating	Independent	0	12,166	9,628	21,794	44.20%	71.70%	<0.001
	Needs supervision		2,395	1,195	3,590	33.30%	11.80%	

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
	Partially assisted		1,791	584	2,375	24.60%	7.80%	
	Fully assisted		2,307	321	2,628	12.20%	8.60%	
Oral hygiene	Independent	0	9,663	8,356	18,019	46.40%	59.30%	<0.001
	Partially assisted		4,925	2,468	7,393	33.40%	24.30%	
	Fully assisted		4,071	904	4,975	18.20%	16.40%	
Washing face	Independent	0	9,895	8,481	18,376	46.20%	60.50%	<0.001
	Partially assisted		4,733	2,371	7,104	33.40%	23.40%	
	Fully assisted		4,031	876	4,907	17.90%	16.10%	
Hair care	Independent	0	10,727	8,885	19,612	45.30%	64.50%	<0.001
	Partially assisted		3,122	1,606	4,728	34.00%	15.60%	
	Fully assisted		4,810	1,237	6,047	20.50%	19.90%	
Cutting nails	Independent	0	5,116	4,543	9,659	47.00%	31.80%	<0.001
	Partially assisted		3,683	2,912	6,595	44.20%	21.70%	
	Fully assisted		9,860	4,273	14,133	30.20%	46.50%	
Buttoning/unbuttoning clothes	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Putting on/ removing top	Independent	0	8,022	7,168	15,190	47.20%	50.00%	<0.001
	Needs supervision		1,670	1,239	2,909	42.60%	9.60%	
	Partially assisted		4,532	2,368	6,900	34.30%	22.70%	

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
	Fully assisted		4,435	953	5,388	17.70%	17.70%	
Putting on/removing pants	Independent	0	7,672	6,916	14,588	47.40%	48.00%	<0.001
	Needs supervision		1,575	1,203	2,778	43.30%	9.10%	
	Partially assisted		3,851	2,228	6,079	36.70%	20.00%	
	Fully assisted		5,561	1,381	6,942	19.90%	22.80%	
Putting on/removing socks	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Cleaning up rooms	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Taking medication	Independent	0	5,798	5,138	10,936	47.00%		<0.001
	Partially assisted		7,106	4,847	11,953	40.60%	39.30%	
	Fully assisted		5,755	1,743	7,498	23.20%	24.70%	
Managing money	Independent	0	4,699	4,264	8,963	47.60%	29.50%	<0.001
	Partially assisted		3,088	2,489	5,577	44.60%	18.40%	
	Fully assisted		10,872	4,975	15,847	31.40%	52.20%	
Severe forgetfulness	Absent	0	18,659	11,728	30,387	38.60%	100.00%	1
Indifference to surroundings	Absent	0	18,659	11,728	30,387	38.60%	100.00%	1
Vision	Normal	0	11,021	7,397	18,418	40.20%	60.60%	<0.001
	Able to see over 1m away		5,294	3,403	8,697	39.10%	28.60%	
	Nearsightedness		1,270	692	1,962	35.30%	6.50%	

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
	Poor sight		368	144	512	28.10%	1.70%	
	Cannot be judged		706	92	798	11.50%	2.60%	
Hearing ability	Normal	0	7,600	5,064	12,664	40.00%	41.70%	<0.001
	Manages to hear		5,849	3,811	9,660	39.50%	31.80%	
	Able to hear loud sounds		4,359	2,614	6,973	37.50%	22.90%	
	Poor hearing		436	195	631	30.90%	2.10%	
	Cannot be judged		415	44	459	9.60%	1.50%	
Expressing intentions	Capable	0	11,419	8,855	20,274	43.70%	66.70%	<0.001
	Sometimes capable		4,066	2033	6,099	33.30%	20.10%	
	Almost incapable		2,078	684	2,762	24.80%	9.10%	
	Incapable		1,096	156	1,252	12.50%	4.10%	
Responding to request	Capable	0	18,659	11,728	30,387	38.60%	100.00%	1
Understanding the daily routine	Capable	0	10,814	8,507	19,321	44.00%	63.60%	<0.001
	Incapable		7,845	3,221	11,066	29.10%	36.40%	
Remembering birth date	Capable	0	15,018	10,641	25,659	41.50%	84.40%	<0.001
	Incapable		3,641	1,087	4,728	23.00%	15.60%	
Short-term memory	Capable	0	9,929	7,622	17,551	43.40%	57.80%	<0.001
	Incapable		8,730	4,106	12,836	32.00%	42.20%	



Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
Remembering one's name	Capable	0	16,886	11,380	28,266	40.30%	93.00%	<0.001
	Incapable		1,773	348	2,121	16.40%	7.00%	
Aware of current season	Capable	0	12,475	9,337	21,812	42.80%	71.80%	<0.001
	Incapable		6,184	2,391	8,575	27.90%	28.20%	
Recognizing places	Capable	0	14,254	10,310	24,564	42.00%	80.80%	<0.001
	Incapable		4,405	1,418	5,823	24.40%	19.20%	
Conviction to be threatened	Absent	0	16,313	10,135	26,448	38.30%	87.00%	0.005
	Sometimes present		1,026	646	1,672	38.60%	5.50%	
	Present		1,320	947	2,267	41.80%	7.50%	
Delusional thoughts and imaginary conversation	Absent	0	15,327	9,578	24,905	38.50%	82.00%	0.137
	Sometimes present		1,286	879	2,165	40.60%	7.10%	
	Present		2046	1,271	3,317	38.30%	10.90%	
Illusion	Absent	0	18,659	11,728	30,387	38.60%	100.00%	1
Emotional instability	Absent	0	14,075	8,873	22,948	38.70%	75.50%	0.004
	Sometimes present		2017	1,367	3,384	40.40%	11.10%	
	Present		2,567	1,488	4,055	36.70%	13.30%	
Day–night reversal	Absent	0	15,084	9,970	25,054	39.80%	82.40%	<0.001
	Sometimes present		1,680	904	2,584	35.00%	8.50%	

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
	Present		1,895	854	2,749	31.10%	9.00%	
Violent behaviors	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Speech repetition	Absent	0	12,311	7,297	19,608	37.20%	64.50%	<0.001
	Sometimes present		1922	1,401	3,323	42.20%	10.90%	
	Present		4,426	3,030	7,456	40.60%	24.50%	
Screaming/Calling out	Absent	0	15,957	10,412	26,369	39.50%	86.80%	<0.001
	Sometimes present		1,274	642	1916	33.50%	6.30%	
	Present		1,428	674	2,102	32.10%	6.90%	
Resistance to care	Absent	0	14,698	9,702	24,400	39.80%	80.30%	<0.001
	Sometimes present		1,675	957	2,632	36.40%	8.70%	
	Present		2,286	1,069	3,355	31.90%	11.00%	
Frequent wandering	Absent	0	17,567	11,161	28,728	38.90%	94.50%	<0.001
	Sometimes present		353	223	576	38.70%	1.90%	
	Present		739	344	1,083	31.80%	3.60%	
Restlessness	Absent	0	17,224	10,936	28,160	38.80%	92.70%	0.007
	Sometimes present		620	327	947	34.50%	3.10%	
	Present		815	465	1,280	36.30%	4.20%	
Not returning after going out	Absent	0	17,889	11,234	29,123	38.60%	95.80%	0.238

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
	Sometimes absent		293	211	504	41.90%	1.70%	
	Present		477	283	760	37.20%	2.50%	
Insistence on going out alone	Absent	0	17,642	11,154	28,796	38.70%	94.80%	0.011
	Sometimes present		429	274	703	39.00%	2.30%	
	Present		588	300	888	33.80%	2.90%	
Hoarding behavior	Absent	0	17,903	11,211	29,114	38.50%	95.80%	0.108
	Sometimes present		244	147	391	37.60%	1.30%	
	Present		512	370	882	42.00%	2.90%	
Mismanagement of fire	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Destroying objects and clothing	Absent	0	18,305	11,528	29,833	38.60%	98.20%	0.3
	Sometimes present		160	99	259	38.20%	0.90%	
	Present		194	101	295	34.20%	1.00%	
Unhygienic behaviors	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Pica behaviors	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Inappropriate sexual behavior	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Movement	Independent	0	7,097	6,243	13,340	46.80%	43.90%	<0.001
	Needs supervision		3,832	2,805	6,637	42.30%	21.80%	
	Partially assisted		2,814	1,456	4,270	34.10%	14.10%	

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
	Fully assisted		4,916	1,224	6,140	19.90%	20.20%	
Cleaning after urination	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Passing urine/ Urination	Independent	0	8,030	7,216	15,246	47.30%	50.20%	<0.001
	Needs supervision		1,225	880	2,105	41.80%	6.90%	
	Partially assisted		3,766	2,167	5,933	36.50%	19.50%	
	Fully assisted		5,638	1,465	7,103	20.60%	23.40%	
Passing motion/ Defecation	Independent	0	8,606	7,582	16,188	46.80%	53.30%	<0.001
	Needs supervision		1,187	807	1,994	40.50%	6.60%	
	Partially assisted		3,190	1,858	5,048	36.80%	16.60%	
	Fully assisted		5,676	1,481	7,157	20.70%	23.60%	
Managing money	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Everyday decision-making	Capable	0	4,761	3,976	8,737	45.50%	28.80%	<0.001
	Capable except during special occasions		7,322	5,246	12,568	41.70%	41.40%	
	Capable but very challenging		4,396	2,045	6,441	31.70%	21.20%	
	Incapable		2,180	461	2,641	17.50%	8.70%	
Forgetfulness	Absent	0	8,570	5,198	13,768	37.80%	45.30%	<0.001
	Sometimes present		3,234	2,464	5,698	43.20%	18.80%	
	Present		6,855	4,066	10,921	37.20%	35.90%	

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
Activity level during daytime	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Frequency of going out	More than once a week	0	8,683	7,055	15,738	44.80%	51.80%	<0.001
	More than once a month		2,811	1,972	4,783	41.20%	15.70%	
	Less than once a month		7,165	2,701	9,866	27.40%	32.50%	
Change in environment	Unknown	0	18,659	11,728	30,387	38.60%	100.00%	1
Self-talk / laughing to oneself	Absent	0	16,720	10,746	27,466	39.10%	90.40%	<0.001
	Sometimes present		643	346	989	35.00%	3.30%	
	Present		1,296	636	1,932	32.90%	6.40%	
Selfish behaviors	Absent	0	16,186	10,195	26,381	38.60%	86.80%	0.135
	Sometimes present		1,021	681	1,702	40.00%	5.60%	
	Present		1,452	852	2,304	37.00%	7.60%	
Disorganized talking	Absent	0	13,812	8,911	22,723	39.20%	74.80%	<0.001
	Sometimes present		1,866	1,244	3,110	40.00%	10.20%	
	Present		2,981	1,573	4,554	34.50%	15.00%	
Not fitting in a group	Absent	0	16,986	10,695	27,681	38.60%	91.10%	0.194
	Sometimes present		731	487	1,218	40.00%	4.00%	
	Present		942	546	1,488	36.70%	4.90%	
Shopping	Independent	0	1,942	1,790	3,732	48.00%	12.30%	<0.001

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
	Needs supervision		134	149	283	52.70%	0.90%	
	Partially assisted		2,793	2,665	5,458	48.80%	18.00%	
	Fully assisted		13,790	7,124	20,914	34.10%	68.80%	
Light cooking	Independent	0	3,624	3,446	7,070	48.70%	23.30%	<0.001
	Needs supervision		199	208	407	51.10%	1.30%	
	Partially assisted		1,369	1,347	2,716	49.60%	8.90%	
	Fully assisted		13,467	6,727	20,194	33.30%	66.50%	
Bedridden (levels)	Independent	0	53	48	101	47.50%	0.30%	<0.001
	J1		1,251	1,130	2,381	47.50%	7.80%	
	J2		2,641	2,368	5,009	47.30%	16.50%	
	A1		3,268	3,049	6,317	48.30%	20.80%	
	A2		4,283	2,850	7,133	40.00%	23.50%	
	B1		1,595	815	2,410	33.80%	7.90%	
	B2		2,693	986	3,679	26.80%	12.10%	
	C1		782	189	971	19.50%	3.20%	
	C2		2,093	293	2,386	12.30%	7.90%	
Dementia (levels)	Independent	0	2,779	2,134	4,913	43.40%	16.20%	<0.001
	I		4,539	3,812	8,351	45.60%	27.50%	

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
	IIa		1,849	1,398	3,247	43.10%	10.70%	
	IIb		3,284	2,052	5,336	38.50%	17.60%	
	IIIa		3,331	1,550	4,881	31.80%	16.10%	
	IIIb		937	354	1,291	27.40%	4.20%	
	IV		1,680	406	2,086	19.50%	6.90%	
	M		260	22	282	7.80%	0.90%	
Short-term memory	Unknown	0	5	4	9	44.40%	0.00%	<0.001
	Capable		6,792	5,119	11,911	43.00%	39.20%	
	Problematic		11,862	6,605	18,467	35.80%	60.80%	
Cognitive ability	Unknown	0	6	5	11	45.50%	0.00%	<0.001
	Independent		6,819	5,389	12,208	44.10%	40.20%	
	Somewhat challenging		4,798	3,364	8,162	41.20%	26.90%	
	Needs assistance		4,117	2,129	6,246	34.10%	20.60%	
	Cannot be judged		2,919	841	3,760	22.40%	12.40%	
Communication ability	Unknown	0	5	8	13	61.50%	0.00%	<0.001
	Capable		8,460	6,689	15,149	44.20%	49.90%	
	Somewhat challenging		4,715	3,004	7,719	38.90%	25.40%	
	Limited to specific requests		3,727	1,645	5,372	30.60%	17.70%	

Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
	Incapable		1,752	382	2,134	17.90%	7.00%	
Eating	Unknown	0	33	7	40	17.50%	0.10%	<0.001
	Independent		16,267	11,336	27,603	41.10%	90.80%	
	Fully dependent		2,359	385	2,744	14.00%	9.00%	
BPSD	Unknown	0	3,551	2,141	5,692	37.60%	18.70%	<0.001
	Present		4,063	2,067	6,130	33.70%	20.20%	
	Absent		11,045	7,520	18,565	40.50%	61.10%	
BPSD	Others	0	392	223	615	36.30%	2.00%	<0.001
	Filthy behaviors		99	31	130	23.80%	0.40%	
	Refusal to care		375	140	515	27.20%	1.70%	
	Delusion		980	573	1,553	36.90%	5.10%	
	Visual/auditory hallucination		15,563	10,185	25,748	39.60%	84.70%	
	Wandering		169	88	257	34.20%	0.80%	
	Sexual disinhibition		1	5	6	83.30%	0.00%	
	Day–night reversal		526	185	711	26.00%	2.30%	
	Assaulting		30	11	41	26.80%	0.10%	
	Violent language		378	149	527	28.30%	1.70%	
	Mismanagement of fire		125	130	255	51.00%	0.80%	



Features	Level	Missing	Stable or Improved (care needs remained the same or decreased) n = 18,659	Deteriorated (care needs increased) n = 11,728	Total N = 30,387	Deterioration Proportion (%)	% in each category	P value
Paralysis in limbs	Pica behaviors		21	8	29	27.60%	0.10%	<0.001
	Others	0	275	260	535	48.60%	1.80%	
Joint movement without restriction	Right_upper_limb		1,282	838	2,120	39.50%	7.00%	<0.001
	Right_lower_limb		349	321	670	47.90%	2.20%	
	Left_upper_limb		3,431	1,423	4,854	29.30%	16.00%	
	Left_lower_limb		8,533	5,360	13,893	38.60%	45.70%	
	No_paralysis		4,789	3,526	8,315	42.40%	27.40%	
	Others	0	1,507	1,109	2,616	42.40%	8.60%	
Medical treatment in the last 2 weeks_Intravenous therapy	Hips		1,442	787	2,229	35.30%	7.30%	<0.001
	Shoulders		3,912	1,941	5,853	33.20%	19.30%	
	Knees		3,655	2,629	6,284	41.80%	20.70%	
	No restrictions		8,143	5,262	13,405	39.30%	44.10%	
	Skin disease treatment	0	341	73	414	17.60%	1.40%	
Medical treatment in the last 2 weeks_Intravenous therapy	Catheter		371	83	454	18.30%	1.50%	<0.001
	Stoma treatment		115	23	138	16.70%	0.50%	
	Monitoring		36	18	54	33.30%	0.20%	
	Respirator		8	3	11	27.30%	0.00%	

<b>Features</b>	<b>Level</b>	<b>Missing</b>	<b>Stable or Improved (care needs remained the same or decreased) n = 18,659</b>	<b>Deteriorated (care needs increased) n = 11,728</b>	<b>Total N = 30,387</b>	<b>Deterioration Proportion (%)</b>	<b>% in each category</b>	<b>P value</b>
	Central venous nutrition		87	4	91	4.40%	0.30%	
	Tracheostomy		26		26	0.00%	0.10%	
	Infusion management		16,559	11,229	27,788	40.40%	91.40%	
	Care for pain		81	52	133	39.10%	0.40%	
	Tube feeding		484	59	543	10.90%	1.80%	
	Dialysis		180	72	252	28.60%	0.80%	
	Oxygen treatment		371	112	483	23.20%	1.60%	

Table B.3. Accuracy and Model Fitness Results of Care Needs Level Predictive Model in Three Years.

Model	Classes	Accuracy (Mean $\pm$ SD), %
GBDT	4	74.64 $\pm$ 0.04
GBDT	9	53.87 $\pm$ 0.55
Logistic Regression	9	41.52 $\pm$ 7.91

# Confusion Matrix

3 year prediction





Fig. B.1. Confusion Matrix of the Four-Category Predictive Model.

# Confusion Matrix

3 year prediction



**Prediction of care level needed in 3 years**

Fig. B.2. Confusion Matrix of the Nine-Category Predictive Model.




Fig. B.3. ROC Curve of the Binary Gradient Boosting Decision Tree Model.




Fig. B.4. Probability distribution of an insureds' predictions.