Infection control report summary

Supplementary File 1

Likely sources of infection in HCP: definitions

Outside community. Among the sequences available for comparison, the likely source of infection was <u>not</u> a patient and was <u>not</u> a co-worker/employee.

Patient source. The most likely source of infection in the HCP was a patient source.

Employee source. The most likely source of infection in the HCP was a co-worker/employee.

Combined patient and employee cluster. A patient to HCP transmission event likely started this cluster and was followed by HCP-to-HCP transmission. However, we are unable to pinpoint the first HCP to become infected and/or are unable to distinguish ongoing sources of transmission as patient-to-HCP and HCP-to-HCP are both possible.

Inconclusive. No consensus sequence available, and/or there were no appropriate comparator sequences available, and/or epidemiological information were insufficient to interpret sequence data

Summary table

Likely source of infection in HCP	Number of cases
Outside community	58 (60.4%)
Patient source (via employee-patient interactions)	4 (4.2%)
Employee source (via employee-employee interactions)	10 (10.4%)
Combined patient and employee cluster	12 (12.5)
Inconclusive	12 (12.5)
Total	96

Total number of patient comparator samples = 140 (96 consensus sequences).

Each case included in the above table is summarized below. For each case, we include the likely source of infection for all involved healthcare personnel (HCP). Next, we provide essential information for each associated sample in the form of a table, including sample collection date, GISAID identifier, Nextstrain clade, and Pangolin lineage. We report clades using the updated Nextstrain clade naming strategy as outlined by Bedford, Hodcroft, and Neher in Virological.org¹. We report lineages according to the Pangolin nomenclature as outlined by Rambaut and colleagues³. A description of each Pangolin lineage can be found at cov-lineages². Next, we provide a very brief overview of any known epidemiological interactions among the involved individuals. The level of epidemiological detail associated with each case is variable, but we have included all known information here. We include a simple alignment showing the consensus sequences mapped against

the Wuhan-Hu-1 reference sequence. Consensus-level differences amongst the reference and the sample sequences are denoted with a vertical black line. Particular variant identities for each sample can be found on the <u>GitHub accompanying this manuscript</u>. Finally, we include a time-resolved phylogenetic tree, built using Nextstrain algorithms, for each case. These trees include all sequences which are publicly available in the GISAID database from the state of Wisconsin. We highlight the samples involved in each case using bolded branches and nodes. An interactive view of this tree can be found <u>here</u>.

¹ https://virological.org/t/updated-nextstain-sars-cov-2-clade-naming-strategy/581

² https://cov-lineages.org/lineages.html

³ Rambaut A, Holmes EC, O'Toole Á, Hill V, McCrone JT, Ruis C, du Plessis L, Pybus OG. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. Nat Microbiol. 2020 Nov;5(11):1403-1407. doi: 10.1038/s41564-020-0770-5. Epub 2020 Jul 15. PMID: 32669681.

Report #1. 2020-04-15.

Likely source of HCP infection

HCP 1. Outside community.


Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	April 2020	hCoV-19/USA/IA-UW-121/2020	20A	B.1
patient A	April 2020	hCoV-19/USA/WI-UW-120/2020	20A	B.1.19
patient B	April 2020	hCoV-19/USA/WI-UW-122/2020	20A	B.1.19

Epidemiological information

In the 14 days before symptom onset, HCP 1 interacted with patients A and B per comprehensive caregiver trace (see Methods, "Sample approvals and sample selection criteria" for further information). HCP 1 reported wearing appropriate personal protective equipment (PPE) while providing care to these patients.

MN908947.3 (Se	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
HCP_1 Patient_A Patient_B	1						1	1 1 1 1 1		1			1 1 1 1 1	ŀ	1



Report #2. 2020-04-16.

Likely source of HCP infection

HCP 1. Outside community (household contact).

HCP 2. Outside community.

HCP 3. Outside community.

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	April 2020	hCoV-19/USA/WI-UW-119/2020	20A	B.1.19
HCP 2	April 2020	hCoV-19/USA/WI-UW-259/2020	20A	B.1.19
HCP 3	April 2020	hCoV-19/USA/WI-UW-260/2020	20C	B.1
Household contact (HCP 1)	April 2020	hCoV-19/USA/WI-UW-120/2020	20A	B.1.19
patient A	March 2020	hCoV-19/USA/WI-UW-118/2020	20C	B.1


patient B	March 2020	hCoV-19/USA/WI-UW-110/2020	20A	B.1.139

HCP 1 tested positive for SARS-CoV-2 after providing care for two SARS-CoV-2 positive patients, patients A and B. HCP 1 also had a household contact who tested positive for SARS-CoV-2 16 days before HCP 1. In the week following HCP 1's positive test, two additional HCP, HCP 2 and HCP 3 tested positive for SARS-CoV-2. HCPs 1, 2, and 3 all work in the same department, but we do not know whether these individuals had any high-risk contact with HCP 1 before their positive test results.

Alignment

	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,90
MN908947.3															500 (2 - 405 00 C 40
HCP_1	1						1	1.1					1.1		- L
HCP_1 HCP_2			1				1	11	1				1 11		
HCP_3	1	1	I	1				1					1		
HCP_1_household-contact	1		1				1	1.1					1 1		1
Patient_A Patient_B	1	1	1					1					1	1	
Patient_B	1					1		1					1		

Phylogeny



Notes

This case was published as an independent case report in Emerging Infectious Disease {32758345}.

Report #3. 2020-06-04.

Likely source of HCP infection

HCP 1. Outside community.


Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	May 2020	hCoV-19/USA/WI-UW-388/2020	20A	B.1.139
HCP-C	Sample was not avai	lable		
patient A	May 2020	hCoV-19/USA/WI-UW-386/2020	20A	B.1.139
patient B	May 2020	hCoV-19/USA/WI-UW-390/2020	20A	B.1.139
patient C	May 2020	hCoV-19/USA/WI-UW-389/2020	20A	B.1.139
patient D	May 2020	hCoV-19/USA/WI-UW-387/2020	20A	B.1.139

Epidemiological information

HCP 1 did not have direct contact with any of the patients included here. In this case, a household contact (HCP-C) of HCP 1 also works in healthcare, but in a different healthcare facility than HCP 1. The healthcare facility employing HCP-C was experiencing a COVID-19 outbreak at the time that HCP 1 tested positive. Patients A-D were patient samples collected from the HCP-C outbreak. A sample from the HCP-C was not available for comparison. Given the similarity in viral sequences between HCP 1 and all four patients from the outside healthcare facility, it is likely HCP 1 was exposed/infected via their household contact, who was likely exposed through patient contact.

MN 0000 47 0	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
MN908947.3															
HCP 1	1 1		1 11 1	1	1			1					T.		1
Patient_A	1 1		1 11 1		1			1.1					I.		1
Patient_B	1 1			1	1			1					1		1
Patient_C	1 1		1 11 1	1	1			1				1	1		1
Patient_D	1.1		1 11 1	1	1			1					1		1



Report #4. 2020-06-04.

Likely source of HCP infection

HCP 1. Patient source (patient D).

Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	May 2020	hCoV-19/USA/WI-UW-391/2020	20A	B.1.139
patient A	May 2020	hCoV-19/USA/WI-UW-392/2020	20A	B.1.276
patient B	May 2020	hCoV-19/USA/WI-UW-393/2020	20A	B.1.139
patient C	May 2020	N/A - no consensus sequence		
patient D	May 2020	hCoV-19/USA/WI-UW-389/2020	20A	B.1.139


Epidemiological information

In the two weeks before symptom onset, HCP 1 provided direct care to patients A-D. HCP 1 wore appropriate PPE while providing care and was also present during patient D's treatment with the Aerobika nebulizer.

Alignment

MN908947.3	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
HCP_1	1		1 11 1	L	1			1				I	I.		1
Patient_A Patient_B		1	1				1			1		1		1 1	1
Patient_D	1 1		1 11 1	1	1			1				1	I.		1

Phylogeny



Notes

The Aerobika nebulizer was added to the list of aerosol-generating procedures requiring enhanced PPE based on this case.

Report #5. 2020-07-08.

Likely source of HCP infection

HCP 1. Outside community.

HCP 2. Outside community (likely source was HCP 1, but infection took place outside of the workplace).


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	June 2020	hCoV-19/USA/WI-UW-588/2020	20C	B.1

HCP 2	June 2020	hCoV-19/USA/WI-UW-610/2020	20C	B.1
patient A	June 2020	hCoV-19/USA/FL-UW-473/2020	20A	B.1.162
patient B	June 2020	hCoV-19/USA/WI-UW-469/2020	20A	B.1.162
patient C	June 2020	hCoV-19/USA/WI-UW-497/2020	20A	B.1.139
patient D	June 2020	hCoV-19/USA/WI-UW-493/2020	20A	B.1.139
patient E	June 2020	hCoV-19/USA/WI-UW-479/2020	20A	B.1.139
patient F	June 2020	N/A - no consensus sequence		

HCP 1 had direct contact while wearing appropriate PPE with patients A-E. HCP 1 and HCP 2 had unmasked interactions (>15 mins) with each other outside of the workplace. HCP 2 did not have contact with any SARS-CoV-2 positive patients in the 14 days before symptom onset.

Alignment

MN908947.3	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000		29,903
HCP_1			1					-						-	1	1 1
HCP_2	1 1		1					1					1	1	1	1 1
Patient_A	1	1	1	1		1	1-	1	1	1	1			1		1
Patient_B	1.1	1	1		1			1	1	1			1	1		
Patient_C			1 11 1	1	1			1			1		1			1
Patient_D	1 1		1 11 1	1	1		1	1				1	1			1
Patient_E	1	I I	1 1				1				1		Ш			



Report #6. 2020-08-13.

Likely source of HCP infection

HCP 1. Outside community (based on epidemiological risk factors). HCP 2. Employee source (likely source was HCP 1).

Samples


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	June 2020	hCoV-19/USA/WI-UW-520/2020	20A	B.1.139
HCP 2	June 2020	hCoV-19/USA/WI-UW-726/2020	20A	B.1.139

Epidemiological information

HCP 1 had a high-risk exposure event in the community before testing positive. This event was indoors, unmasked, and lasted longer than 15 minutes. HCP 1 works in the same department as HCP 2. Neither HCP 1 nor HCP 2 provided direct care to patients diagnosed with COVID-19 in the 14 days before their symptom onset. HCP 2 reported wearing a mask around all coworkers except while eating in the breakroom. HCP 2 reports removing their mask while eating, but maintaining a 6-foot physical distance from others during this time.

Alignment





Report #7. 2020-08-04.

Likely source of HCP infection

HCP 1. Outside community.


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	July 2020	hCoV-19/USA/WI-UW-973/2020	20C	B.1.2
patient A	July 2020	N/A - no consensus sequence		
patient B	July 2020	N/A - no consensus sequence		
patient C	July 2020	N/A - no consensus sequence		
patient D	July 2020	N/A - no consensus sequence		
patient E	July 2020	N/A - no consensus sequence		
patient F	July 2020	hCoV-19/USA/WI-UW-852/2020	20A	B.1.139
patient G	July 2020	N/A - no consensus sequence		
patient H	July 2020	hCoV-19/USA/WI-UW-870/2020	20A	B.1.139
patient I	July 2020	hCoV-19/USA/WI-UW-847/2020	20A	B.1.139
patient J	July 2020	N/A - no consensus sequence		
patient K	July 2020	N/A - no consensus sequence		
patient L	July 2020	hCoV-19/USA/WI-UW-927/2020	20A	B.1.139
patient M	July 2020	hCoV-19/USA/WI-UW-878/2020	20C	B.1.369
patient N	July 2020	N/A - no consensus sequence		
patient O	July 2020	hCoV-19/USA/WI-UW-886/2020	20A	B.1.162
patient P	July 2020	hCoV-19/USA/WI-UW-880/2020	20A	B.1.139
patient Q	July 2020	hCoV-19/USA/WI-UW-894/2020	20B	B1.1
patient R	July 2020	N/A - no consensus sequence	<u>-</u>	
patient S	July 2020	hCoV-19/USA/WI-UW-895/2020	20C	B.1.1369
patient T	July 2020	hCoV-19/USA/WI-UW-876/2020	20C	B.1.369
patient U	July 2020	hCoV-19/USA/WI-UW-931/2020	20C	B.1.369
patient V	July 2020	N/A - no consensus sequence		

patient W	July 2020	N/A - no consensus sequence		
patient X	July 2020	hCoV-19/USA/WI-UW-898/2020	20A	B.1.369
patient Y	July 2020	hCoV-19/USA/WI-UW-883/2020	20C	B.1
patient Z	July 2020	N/A - no consensus sequence	<u>-</u>	
patient AA	July 2020	N/A - no consensus sequence		

HCP 1 collected nasopharyngeal specimens from patients with suspected COVID-19. HCP 1 wore appropriate PPE while collecting these specimens and reported no breach in PPE. Patients A-AA were collected in the 14 days before symptom onset in HCP 1.

Alignment

	1	2,500	5,000	7,500	10,000	12,500	15,000	17,500	20,000	22,500	25,000		29,90
MN908947.3													
HCP_1	1 1	1 1			1		1		1	1	1	1 1	1
Patient F		1	1 1	1			1		1				1
Patient_H	1.1	1 1		1	1		1		1				1
Patient_I	1.1	1		1			1		1	1 1			1
Patient_L	1.1	1	11 1	1			1		1	1			1
Patient_M	1 1	1	1.1	I			1	1 1		1 1	п		1
Patient O	1.1	1		1			1 1		1	1	1		1
Patient_P Patient_Q	1.1	1		1			1		1				1
Patient Q	1	1.1	l.		1 1		н		1	1		1 1	
Patient_S	1 1	1	1				11	1		1	1 1		1 1
Patient_T	1 1	1	1				11	1		1	1 1		1 1
Patient_U	1 1	1	1				-11 - 1	1		1	1 1 1		1 1
Patient_Y	1 1	1					1			1	1 1	1	1.1
Patient_X	1.1	1 1		1									1



Report #8. 2020-08-18.

Likely source of HCP infection

HCP 1. Outside community.


Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	July 2020	hCoV-19/USA/WI-UW-1022/2020	20B	B.1.1.73
patient A	July 2020	N/A - no consensus sequence		
patient B	Unknown	N/A - no consensus sequence		
patient C	July 2020	N/A - no consensus sequence		
patient D	July 2020	hCoV-19/USA/WI-UW-923/2020	20A	B.1.240
patient E	June 2020	hCoV-19/USA/WI-UW-614/2020	20C	B.1.330
patient F	July 2020	hCoV-19/USA/WI-UW-875/2020	20C	B.1.369
patient G	July 2020	hCoV-19/USA/WI-UW-889/2020	20A	B.1.139
patient H	July 2020	hCoV-19/USA/WI-UW-1100/2020	20A	B.1.139

Epidemiological information

HCP 1 likely did not have direct interactions with any of the patients listed here. HCP 1 did, however, perform cleaning duties in the rooms of each of these patients in the 14 days before symptom onset.

MN908947.3	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
HCP 1	1	1	1.1			1	Ш				1		11		1 1
Patient_D			1	1						1	1	1-1	1		1
Patient_E		11 1	1		1 1			1		1			1	1	
Patient_F		1 1	1					1 1	Ш			1	11 1	11 1	1
Patient_G			1 11	1	1			1			1		1		1
Patient H	1		1 11	1	1			1	1			1	1 1		11



Report #9. 2020-08-21.

Likely source of HCP infection


HCP 1. Outside community.

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	July 2020	hCoV-19/USA/WI-UW-965/2020	20B	B.1.1.73
patient A	July 2020	hCoV-19/USA/WI-UW-765/2020	20A	B.1.139
patient B	July 2020	hCoV-19/USA/WI-UW-792/2020	20A	B.1.139
patient C	July 2020	hCoV-19/USA/WI-UW-761/2020	20C	B.1.3
patient D	July 2020	hCoV-19/USA/WI-UW-781/2020	20A	B.1.139
patient E	July 2020	hCoV-19/USA/WI-UW-794/2020	20A	B.1.139
patient F	July 2020	hCoV-19/USA/WI-UW-796/2020	20C	B.1.294
patient G	July 2020	hCoV-19/USA/WI-UW-790/2020	20A	B.1.139

patient H	July 2020	hCoV-19/USA/WI-UW-768/2020	20A	B.1.139
patient I	July 2020	hCoV-19/USA/WI-UW-803/2020	20A	B.1.162
patient J	July 2020	N/A - no consensus sequence		
patient K	July 2020	hCoV-19/USA/MO-UW-771/2020	20C	B.1.370
patient L	July 2020	N/A - no consensus sequence		
patient M	July 2020	N/A - no consensus sequence		
patient N	July 2020	hCoV-19/USA/WI-UW-784/2020	20A	B.1
patient O	July 2020	hCoV-19/USA/WI-UW-787/2020	20A	B.1.139
patient P	July 2020	N/A - no consensus sequence		
patient Q	July 2020	hCoV-19/USA/WI-UW-786/2020	20A	B.1.112
patient R	July 2020	hCoV-19/USA/WI-UW-766/2020	20C	B.1.370
patient S	July 2020	hCoV-19/USA/WI-UW-798/2020	20A	B.1

HCP 1 collected nasopharyngeal specimens from patients with suspected COVID-19. HCP 1 wore appropriate PPE while collecting these specimens and reported no breach in PPE. Patients A-S were collected in the 14 days before symptom onset in HCP 1.

	1 2,0	000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000		29,903
MN908947.3																
HCP_1	1	1			1		1	1			11		11 1			1 1
Patient_A	1 1	1	11 1	1	1			1			1		1			1
Patient_B	1 1	1		1	1			1	1			1	1			1
Patient_C	1 1 1	1 1					1	1	1	1			11	1	1	
Patient D		1	11 1	1	1			1				1	1			1
Patient_E			11	1				1			1		1			1
Patient_F	1 1	1	1			1		1			1	1	1	11		
Patient_G			1	3	1		1	1				1	1			
Patient H	1	ា	11 1	1	1			F				1	1			1
Patient_I	1	1					1	I I	1	1		11	1	1		1
Patient_K	1 11 1		6	1				1	1			1	1		1	1
Patient_N	4	1						1			1	1	1		1	1
Patient_O	1 1	1		1				1			1		1			1
Patient_Q	1	1	1	1				1			I.		1	1		
Patient_R	1 1	1				1 1		1		1			1	1 1	П	
Patient_S	4	ш		1				1			1	1	1		1	1



Report #10. 2020-08-26.

Likely source of HCP infection

HCP 1. Patient source (patient G).


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	August 2020	hCoV-19/USA/WI-UW-1125/2020	20A	B.1.139
patient A	August 2020	hCoV-19/USA/WI-UW-867/2020	20A	B.1.255
patient B	July 2020	hCoV-19/USA/WI-UW-889/2020	20A	B.1.139
patient C	July 2020	N/A - no consensus sequence		
patient D	July 2020	N/A - no consensus sequence		
patient E	August 2020	N/A - no consensus sequence		
patient F	July 2020	hCoV-19/USA/WI-UW-923/2020	20A	B.1.240
patient G	July 2020	hCoV-19/USA/WI-UW-1100/2020	20A	B.1.139
patient H	July 2020	hCoV-19/USA/WI-UW-998/2020	20B	B.1.1.73

patient I	August 2020	hCoV-19/USA/WI-UW-1035/2020	20A	B.1.162
patient J	July 2020	N/A - no consensus sequence		
patient K	July 2020	N/A - no consensus sequence		
patient L	July 2020	N/A - no consensus sequence		
patient M	July 2020	hCoV-19/USA/WI-UW-1054/2020	20B	B.1.1.73
patient N	July 2020	hCoV-19/USA/WI-UW-805/2020	20C	B.1.330
patient O	July 3030	hCoV-19/USA/WI-UW-1090/2020	20B	B.1.1.73

HCP 1 had direct contact while wearing appropriate PPE with patients A-O in the 14 days before symptom onset.

Alignment

MN908947.3	1 2,0	00	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000		29,903
HCP_1	1.1	1	11.1	1	1			T	1			1	1 1			11
Patient_A	1	Т						I		1			1	1	_	
Patient_B	1 1	1	11 1	1	I			1			I		1		_	1
Patient_F	1	1		1				1		1	016	1 - 1	1			1
Patient_G	1 1	1	11 1	1	1			1	1			1	1 1		-	1.1
Patient_H								1			1		11			- F
Patient_I	1	1	1				1	1	1	1			1 1	1	-	
Patient_M	11				1		1 1			11			11 1			-
Patient_N	1 1 1	1 1			1 1			1		1		246	1	1	_	
Patient_O	1	1					1.1	П			1	1	11 1	1	-	1 1



Report #11. 2020-09-11.

Likely source of HCP infection

HCP 1. Outside community.


Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	August 2020	hCoV-19/USA/WI-UW-1213/2020	20C	B.1.2
patient A	August 2020	N/A - no consensus sequence		
patient B	August 2020	hCoV-19/USA/WI-UW-1163/2020	20B	B.1.1.130
patient C	August 2020	hCoV-19/USA/WI-UW-1240/2020	20C	B.1.2
patient D	August 2020	hCoV-19/USA/WI-UW-1170/2020	20A	B.1.5

Epidemiological information

HCP 1 had direct contact while wearing appropriate PPE with patients A-D in the 14 days before symptom onset. HCP 1 denied any lapses in PPE and did not interact with coworkers without a surgical mask on.

MN908947.3	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
HCP_1	1	1	ı			1		1.1		1	1	1	1	1.1	11.1.1
Patient_B	-	1 1 1	I				1	1	1				1 1		1
Patient_C	1	l.	1 1		1	1 1		1.1		1		1	1	1.1	1 1 1 1
Patient_D	1		l.					1					1		



Report #12. 2020-09-11.

Likely source of HCP infection

HCP 1. Inconclusive.

HCP 2. Outside community.


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	August 2020	N/A - no consensus sequence		
HCP 2	August 2020	hCoV-19/USA/WI-UW-1278/2020	20A	B.1.139
patient A	July 2020	hCoV-19/USA/WI-UW-1100/2020	20A	B.1.139
patient B	July 2020	N/A - no consensus sequence		
patient C	July 2020	N/A - no consensus sequence		
patient D	August 2020	N/A - no consensus sequence		

patient E	July 2020	hCoV-19/USA/WI-UW-850/2020	20C	B.1.2
patient F	August 2020	hCoV-19/USA/WI-UW-1248/2020	20B	B.1.1.244
patient G	August 2020	hCoV-19/USA/WI-UW-1204/2020	20A	B.1.139
patient H	August 2020	N/A - no consensus sequence		

HCP 1 and HCP 2 are household contacts and one or both of these HCP provided direct patient care to patients A-H while wearing appropriate PPE in the two weeks before their symptom onset.

Alignment

MN908947.3 (Se	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
10111908947.5 (3e															
HCP_2		1		1	1		1	1	1			1	1	11	1
Patient_A	1 1			1	1		~	1	1			1	1 1	266 - 296	1.1
Patient_E	1 1	1				1 1	1	l.		1			1	1	1
Patient F		1			1.1	1 1	1	п т		940	1	1	1	1	1 11
Patient_G					12		1			1			1		



Report #13. 2020-09-14.

Likely source of HCP infection

HCP 1. Outside community.


Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	August 2020	hCoV-19/USA/WI-UW-1305/2020	20A	B.1.139
patient A	July 2020	hCoV-19/USA/WI-UW-1100/2020	20A	B.1.139
patient B	August 2020	hCoV-19/USA/WI-UW-1231/2020	20C	B.1.337
patient C	August 2020	N/A - no consensus sequence		
patient D	August 2020	N/A - no consensus sequence		

Epidemiological information

HCP 1 provided direct care while wearing appropriate PPE to patients A-D in the 14 days before symptom onset.

Alignment



2019-Dec 2021-Feb 200-Apr 200-Jun 2020-Aug 202-Oct 2020-Dec 2021-Feb

Report #14. 2020-09-22.

Likely source of HCP infection

HCP 1. Outside community.

HCP 2. Combined patient and employee cluster.

HCP 3. Inconclusive.

HCP 4. Combined patient and employee cluster.

HCP 5. Outside community.

HCP 6. Employee source (HCP 9 is a likely source of infection, although the HCP 6 \rightarrow HCP 9 is also possible).

HCP 7. Outside community.

HCP 8. Inconclusive.

HCP 9. Outside community.

HCP 10. Inconclusive.

HCP 11. Combined patient and employee cluster.

HCP 12. Combined patient and employee cluster.

HCP 13. Combined patient and employee cluster.

HCP 14. Inconclusive.

HCP 15. Inconclusive.

HCP 16. Combined patient and employee cluster.

HCP 17. Combined patient and employee cluster.

HCP 18. Inconclusive.

HCP 19. Inconclusive.

HCP 20. Combined patient and employee cluster.

HCP 21. Inconclusive.


Patients A and C are the patients involved in this combined patient and employee cluster.

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	September 2020	hCoV-19/USA/WI-UW-1348/2020	20G	B.1.2
HCP 2	September 2020	hCoV-19/USA/WI-UW-1344/2020	20G	B.1.2
HCP 3	September 2020	N/A - no consensus sequence		
HCP 4	September 2020	hCoV-19/USA/WI-UW-1412/2020	20G	B.1.2
HCP 5	September 2020	hCoV-19/USA/WI-UW-1325/2020	20G	B.1.5
HCP 6	September 2020	hCoV-19/USA/WI-UW-1480/2020	20B	B.1.1.251
HCP 7	September 2020	hCoV-19/USA/WI-UW-1477/2020	20G	B.1.369
HCP 8	September 2020	N/A - no consensus sequence		
HCP 9	September 2020	hCoV-19/USA/WI-UW-1475/2020	20B	B.1.1.251
HCP 10	September 2020	N/A - no consensus sequence		

HCP 11	September 2020	hCoV-19/USA/WI-UW-1441/2020	20G	B.1.2
HCP 12	September 2020	hCoV-19/USA/WI-UW-1445/2020	20G	B.1.2
HCP 13	September 2020	hCoV-19/USA/WI-UW-1452/2020	20G	B.1.2
HCP 14	September 2020	N/A - no consensus sequence		
HCP 15	September 2020	N/A - no consensus sequence		
HCP 16	September 2020	hCoV-19/USA/WI-UW-1459/2020	20G	B.1.2
HCP 17	September 2020	hCoV-19/USA/WI-UW-1436/2020	20G	B.1.2
HCP 18	September 2020	N/A - no consensus sequence		
HCP 19	September 2020	N/A - no consensus sequence		
HCP 20	September 2020	hCoV-19/USA/WI-UW-1461/2020	20G	B.1.2
HCP 21	September 2020	N/A - no consensus sequence		
patient A	September 2020	hCoV-19/USA/WI-UW-1406/2020	20G	B.1.2
patient B	September 2020	hCoV-19/USA/WI-UW-1499/2020	20G	B.1.2
patient C	September 2020	hCoV-19/USA/WI-UW-1900/2020	20G	B.1.2
patient D	September 2020	hCoV-19/USA/WI-UW-1941/2020	20G	B.1.2

These HCP work in the same department. HCP 2, 4, 11, 12, 13, 16, and 20 provided direct care to patient A and/or patient C. HCP 17 and HCP 1 did not provide direct care to patient A or C.

MN908947.3	1	2,500	5,000	7,500	10,000	12,500	15,000	17,500	20,000		22,500	25,000	29,903
WIN900947.5	D				orf1ab g	ene					S gene		N.
			orf1a	ab CDS							S CDS		N.)
HCP_1	1 1	11 1			1.1		1	l	1	11	Ш	1.1	1 111
HCP_2 HCP_4 HCP_5 HCP_6 HCP_7 HCP_9		11 1			11		1	1	1	11			1 1 1
HCP_4	1 1	11 1			1.1		1	1	1	11	1		1 1 1
HCP ₅		1					1		1		1		
HCP_6	1 1	Ĩ			1	1	1.1			"	1 1	1	1
HCP ⁷	1 1	1 1		1				1 1			1.1		1 1
HCP_9	1 1	I			1	1	1.1			11	1 1	1	
HCP ¹¹	1 1	11 1			1.1		1	1	1	11	1	11	1 1 1
HCP_11 HCP_12 HCP_13 HCP_16 HCP_17	1 1	11 1			1.1		1	1	1	11	1	11	1 1 1
HCP ¹³	1 1	11 1			11			1	1	11	1	11	1 1 1
HCP ¹⁶	1 1	11 1			1.1		1	1	1	11	1	1.1	1 1 1
HCP ¹⁷	1 1	11 1			11		1	1	1	11	1	11	1 1 1
HCP_20	1 1	11 1			1.1		1	1	1	11	1	11	1 1 1
Patient_A	1 1	11 1			11		1	I.	1	11	1	11	1 1 1
Patient_B	1 1	I		1	1		11	1		1	11	1 11	1 1 1 1
Patient_C	1				11		1	1	1	11	1	11	1 1 1
Patient D	1 1	1		1	1		11	1	1		1	11	111



Report #15. 2020-09-18.

Likely source of HCP infection

HCP 1. Outside community.


Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	September 2020	hCoV-19/USA/WI-UW-1347/2020	20G	B.1.2
patient A	August 2020	hCoV-19/USA/WI-UW-1220/2020	20G	B.1.2

Epidemiological information

HCP 1 provided direct care to patient A while wearing appropriate PPE and with no reported lapses in PPE use.

MN908947.3	1 2	,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
HCP_1	1 1	16			п	1		1.1		1		1	1	1.1	11.11.1
Patient_A	1 1	0			1	1		1.1		1	1	1	1	1.1	11 1 1



Report #16. 2020-10-29.

Likely source of HCP infection

HCP 1. Employee source (HCP 2). HCP 2. Inconclusive.

HCP 3. Outside community.

Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	September 2020	hCoV-19/USA/WI-UW-1901/2020	20G	B.1.2
HCP 2	September 2020	hCoV-19/USA/WI-UW-1350/2020	20G	B.1.2
HCP 3	September 2020	hCoV-19/USA/WI-UW-1898/2020	20A	B.1


Epidemiological information

Contact tracing revealed HCP 2 worked for two days prior to symptom onset and may have had unmasked contact with HCP 1 during overlapping meal breaks. Contact tracing additionally revealed HCP 3 had a high-risk exposure even lasting >15 minutes in the outside community prior to testing positive.

Alignment

MN908947.3	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
		1 1	1			1	r.	1 1		1		1.1	11	1.1	1 1 1
HCP_1 HCP_2 HCP_3		1 1	1			1	1	1 1		1		1.1	11	1.1	1 1 1
HCP_3	1	1	1		1	1	1	1			1		1	I	1 11

Phylogeny



Report #17. 2020-10-29.

Likely source of HCP infection

HCP 1. Outside community.


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)			
HCP 1	September 2020	hCoV-19/USA/WI-UW-1895/2020	20C	B.1.369			
patient A	September 2020	N/A - no consensus sequence					
patient B	July 2020	hCoV-19/USA/WI-UW-774/2020	20A	B.1.139			
patient C	August 2020	hCoV-19/USA/WI-UW-1198/2020	20A	B.1.139			

patient D	August 2020	hCoV-19/USA/WI-UW-1166/2020	20A	B.1.139			
patient E	August 2020	N/A - no consensus sequence					
patient F	September 2020	N/A - no consensus sequence					
patient G	August 2020	N/A - no consensus sequence					
patient H	August 2020	hCoV-19/USA/WI-UW-1301/2020	20C	B.1			
patient I	August 2020	N/A - no consensus sequence					

HCP 1 provided direct care to patients A-I while wearing appropriate PPE and with no reported lapses in PPE use. HCP 1 also had a household contact with confirmed SARS-CoV-2 infection who had symptoms onset prior to HCP 1's symptom onset.

Alignment

MN908947.3	1 2,0	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
HCP_1	1 1	1 1		I.			11	1 11				1 1 1	I	1 1
Patient_B Patient_C		1 11	1 1	1		1				I		1		1
Patient_D Patient_H		1 11 1	1 1		1.1	1		1	'	l.		1		П



Report #18. 2020-10-29.

Likely source of HCP infection

HCP 1. Outside community.

HCP 2. Outside community.

HCP 3. Outside community.

Samples


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	September 2020	hCoV-19/USA/WI-UW-1475/2020	20B	B.1.1.251
HCP 2	September 2020	hCoV-19/USA/WI-UW-1896/2020	20B	B.1.1.251
HCP 3	September 2020	hCoV-19/USA/WI-UW-1894/2020	20B	B.1.1.251

Epidemiological information

HCP 1 worked with HCP 3. HCP 3 was a household contact of HCP 2. None of these HCP had direct interactions with patients with known SARS-CoV-2 infection.

Alignment

MN908947.3 (Se	. 1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
HCP_1 HCP_2 HCP_3	1	1	1				1						1 1		1
HCP_3		1	1 1				1	1 1 1				11 1	1 1		11



Report #19. 2020-11-05.

Likely source of HCP infection

HCP 1. Outside community.

Samples


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	October 2020	hCoV-19/USA/WI-UW-1928/2020	19A	B.1
patient A	September 2020	hCoV-19/USA/WI-UW-1930/2020	20G	B.1.2

Epidemiological information

In the two weeks before symptom onset, HCP 1 provided direct care to patient A. HCP 1 wore appropriate PPE while providing care and reported no lapses in PPE use.

Alignment

MN908947.3	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000		29,903
HCP_1 patient_A	1	1	1		1	1		1.1		1		1	1	1.1	1.1	



Report #20. 2020-11-05.

Likely source of HCP infection

HCP 1. Outside community.

Samples


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	October 2020	hCoV-19/USA/WI-UW-1931/2020	20G	B.1.2
patient A	September 2020	hCoV-19/USA/WI-UW-1935/2020	20A	B.1

Epidemiological information

In the two weeks before symptom onset, HCP 1 provided direct care to patient A. HCP 1 wore appropriate PPE while providing care and reported no lapses in PPE use.

Alignment

	-44														
	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
MN908947.3					÷		-								
HCP_1	-	E	1			1 1		1		1		1	1	1.1	
patient_A		1	1 1			1		1 1 1				11 1	1	1	1



Report #21. 2020-11-05.

Likely source of HCP infection

HCP 1. Outside community.

HCP 2. Outside community.


Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	October 2020	hCoV-19/USA/WI-UW-1929/2020	20G	B.1.2
HCP 2	October 2020	hCoV-19/USA/WI-UW-1938/2020	20B	B.1.1.73
patient A	September 2020	N/A - no consensus sequence		
patient B	September 2020	N/A - no consensus sequence		
patient C	September 2020	hCoV-19/USA/WI-UW-1934/2020	20G	B.1.2
patient D	September 2020	hCoV-19/USA/IL-UW-1937/2020	20G	B.1.2
patient E	October 2020	N/A - no consensus sequence		

Epidemiological information

HCP 1 and 2 worked in the same department and both provided direct patient care to patients A-E in the two weeks before their onset of symptoms. HCP 1 and HCP 2 reported no lapses in PPE with each other or with their patients.

MN908947.3	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
HCP_1	1	1	1		1	1		1.1		1		1	1.1	1.1	1 1 1 1
HCP_2	-1	1 1					1 11 1	1					1.1	1	1 1
patient_C	- 1	1 1	1		1	1 1	1	1.1		1		1	11	1.1	1 1 1 1
patient_D	-1	1	1	1		1	1	1				1	1	1.1	1 1 1



Report #22. 2020-11-05.

Likely source of HCP infection

HCP 1. Outside community.

Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	October 2020	hCoV-19/USA/WI-UW-1936/2020	20G	B.1.2
patient A	July 2020	hCoV-19/USA/WI-UW-1100/2020	20A	B.1.139


Epidemiological information

HCP 1 performed a postmortem examination on patient A. At the time the patient expired, they were known to have an active COVID-19 infection.

Alignment

MN908947.3	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
HCP	-	I.	1		1	1	L	1.1		1		1	I.	1.1	1 1 1 1
patient_A	1 1		1 11 1	1	1			1	1			1	1 1		11

Phylogeny



Report #23. 2020-11-05.

Likely source of HCP infection

- HCP 1. Outside community (household contact).
- HCP 2. Employee source.
- HCP 3. Employee source.
- HCP 4. Employee source.
- HCP 5. Outside community.

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	October 2020	hCoV-19/USA/WI-UW-1933/2020	20G	B.1.2
HCP 2	October 2020	hCoV-19/USA/WI-UW-1926/2020	20G	B.1.2


HCP 3	October 2020	hCoV-19/USA/WI-UW-1939/2020	20G	B.1.2
HCP 4	October 2020	hCoV-19/USA/WI-UW-1927/2020	20G	B.1.2
HCP 5	October 2020	hCoV-19/USA/WI-UW-1932/2020	20A	B.1.139

All of these HCP work in the same department, with the exception of HCP 1 who is a household contact of HCP 2. HCP 2-5 reported sharing an unmasked meal together prior to testing positive.

Alignment

MN908947.3 (Se.	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
10110300347.3 (36.															
HCP_1	-	1	1		1	1 11	1	1.1	1	1		1	1	1.1	1 1 1
HCP_1 HCP_2	-	1	1		1	1 11	1	1.1	1	1		1	1	1.1	
HCP_3			1		1	1 11	L.	1.1	1			1	1	1.1	1 1 1 1
HCP_4	-1	1	1		1	1 11	1	1.1	1	1		1	1	1.1	1 1 1 1
HCP_5			1 11 1	1 1	12						1	3	1		1 1

Phylogeny



Report #25. 2020-12.08.

Likely source of HCP infection HCP 1. Patient source (patient A).

Samples


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	October 2020	hCoV-19/USA/WI-UW-2326/2020	20C	B.1.2
patient A	October 2020	hCoV-19/USA/WI-UW-2362/2020	20C	B.1.2
patient B	October 2020	hCoV-19/USA/WI-UW-2389/2020	20C	B.1.2

Epidemiological information

In the two weeks before symptom onset, HCP 1 provided direct care to patients A and B.

MN908947.3 (Se	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
HCP_1				1	1	1		F 1		1		1	1	1.1	1 1 1 1
Patient_A	1			1	1	1		1 1		1		1	1	1.1	1 1 1 1
Patient_B	-	1 1			1	1 1		1.1		1		1	- 11	1.1	1 1 1 1

Alignment



Report #27. 2020-12.09.

Likely source of HCP infection

HCP 1. Outside community.

HCP 2. Outside community.

HCP 3. Outside community.

HCP 4. Outside community.


Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	October 2020	hCoV-19/USA/WI-UW-2226/2020	20A	B.1
HCP 2	October 2020	N/A	20G	B.1.2
HCP 3	October 2020	hCoV-19/USA/WI-UW-2227/2020	20G	B.1.2
HCP 4	October 2020	hCoV-19/USA/WI-UW-2228/2020	20G	B.1.2
patient A	September 2020	hCov-19/USA/IL-UW-2327/2020	20A	B.1
patient B	September 2020	hCoV-19/USA/WI-UW-1972/2020	20G	B.1

Epidemiological information

HCP 1-4 work in the same department. HCP 1-4 all had direct interactions with patients A and/or B during the 14 days prior to symptom onset.

/ //N908947.3 (Se	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000		29,903
HCP_1	1				1		1		1			11	-		
HCP_2 HCP_3							1 1 1		1		1	1	- 1	1	
HCP_4	1	1		1	1	1	11			Ш	1	11	1	1	1 1



Report #28. 2020-12.09.

Likely source of HCP infection

HCP 1. Outside community.

HCP 2. Outside community.

HCP 3. Outside community.

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
Patient A	November 2020	hCoV-19/USA/WI-UW-2265/2020	20G	B.1.2
HCP 1	October 2020	hCoV-19/USA/WI-UW-2266/2020	20G	B.1.2
HCP 2	October 2020	hCoV-19/USA/WI-UW-2267/2020	20G	B.1.2
HCP 3	October 2020	hCoV-19/USA/WI-UW-2268/2020	20G	B.1.2

HCP 1-3 provided direct patient care to patient A in the 14 days before symptom onset. All HCP reported appropriate use of PPE with no lapses. HCP 1 and 2 had no known interactions prior to their infections. HCP 2 and HCP 3 both attended a high-risk community event for greater than 15 minutes together.

Alignment

P_1	1 2,000	4,000	6,000	8,000 I I	10,000 I I I	12,000	14,000	1		20,000	22,000	24,000	26,000	
23	1 1	I	Ĺ	-	1		1.1		I		I	1	I I	1 1 1
nylogen	У													
20	G									U :	SA/WI-U			
											USA/W	/WI-UW- /I-UW-22	2266/2020 267/2020	
											o US/	∿	/-2265/2020	
												_		
		C			-									
										1				

Report #29. 2020-12.09.

Likely source of HCP infection

HCP 1. Employee source (HCP 3).

HCP 2. Outside community.

HCP 3. Outside community.

Samples


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	November 2020	hCoV-19/USA/WI-UW-2269/2020	20G	B.1.2
HCP 2	November 2020	hCoV-19/USA/WI-UW-2270/2020	20G	B.1.2
HCP 3	November 2020	hCoV-19/USA/WI-UW-2271/2020	20G	B.1.2

Epidemiological information

HCP 1-3 work in the same department. HCP 1 and 3 attended an in person meeting together and reported sitting 6-feet apart while wearing masks. HCP 2 did not attend this meeting. HCP 3 reported symptoms before HCP 1.

Alignment

MN908947.3	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000			29,903
HCP_1 HCP_2 HCP_3	11	1	1 1		1	1		1.1		1		1	1	1.1		1 1	1 1
HCP_2		1	п		1	1	1	1 11		1		1	1	11	1	1 1	1
HCP_3	1	1	1 1		1	1		1.1		1		1	1	1.1		1 1	1 1



Report #30. 2020-12.09.

Likely source of HCP infection

HCP 1. Patient source (patient C).


Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	November 2020	hCoV-19/USA/WI-UW-2272/2020	20G	B.1.2
patient A	November 2020	hCov-19/USA/WI-UW-2324/2020	20G	B.1.2
patient B	October 2020	hCoV-19/USA/WI-UW-2273/2020	20G	B.1.2
patient C	November 2020	hCoV-19/USA/WI-UW-2274/2020	20G	B.1.2
patient D	November 2020	hCoV-19/USA/WI-UW-2275/2020	20A	B.1.139
patient E	November 2020	hCoV-19/USA/WI-UW-2276/2020	20G	B.1.2

Epidemiological information

HCP 1 provided direct patient care to patients A-E and reported no lapses in PPE use.

	1		2,000	4,0	00	6,000	8,0	000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	28,000	29,903
MN908947.3 (Se																			
HCP 1		1	1	1			1	1	1 1		1 1	ļ.	1		1.1	11	1 1 1	111	1 1
Patient A	-	1.1		1			1	1	11		1		1 1		1	1.1	1.1	1 1	1 1
Patient_B	- 11	1				1		1	1		1		1		1	1	1.1	1 - 1	1 1
Patient_C	-	1	1	1			1	1	1 1		1 1		1		1	11	1 1 1	1.1.1	1 1
Patient_D		- 1		11 11		1 1			1 1		11		1	1 1		1 1			
Patient_E	-1	- 1		1		1		1	1		1		1		1	1	1111	11-1	1 1



Report #31. 2020-12.09.

Likely source of HCP infection

- HCP 1. Outside community.
- HCP 2. Outside community.
- HCP 3. Outside community.
- HCP 4. Employee source (HCP 2).
- HCP 5. Employee source (HCP 2).
- HCP 6. Outside community.
- HCP 7. Outside community.

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	November 2020	hCoV-19/USA/WI-UW-2312/2020	20C	B.1.363
HCP 2	November 2020	hCoV-19/USA/WI-UW-2277/2020	20G	B.1.2
HCP 3	November 2020	hCoV-19/USA/WI-UW-2278/2020	20G	B.1.2
HCP 4	November 2020	hCoV-19/USA/WI-UW-2279/2020	20G	B.1.2


HCP 5	November 2020	hCoV-19/USA/WI-UW-2280/2020	20G	B.1.2
HCP 6	November 2020	hCoV-19/USA/WI-UW-2281/2020	20G	B.1.2
HCP 7	November 2020	hCoV-19/USA/WI-UW-2282/2020	20G	B.1.2

These HCP are all in the same department. Their level of interaction with each other is unclear.

Alignment

MN908947.3	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
CP_1	1911		1				11	I	1	1	1	1	1	1 1	1
ICP_2	121	1	1		1	1		1.1		1		1 1	1	1.1	1 1 1 1
CP_3		1	1 1	11 1		1		1		1		1	1	1 11	II 1 1
P_4		1	1	1	1	1		1.1		1		1 1	1	1.1	1 1 1 1
CP_5	-1	1	1	1	1	1		1.1		1		1 1	1	1.1	1 1 1 1
CP_6	1	1	1			1		1				1	1	1	1 1 1
CP_7	1	1	1	1	1	1		1.1		1		1	1	11 1	

Phylogeny



Report #32. 2020-12.09.

Likely source of HCP infection HCP 1. Outside community. HCP 2. Outside community.

- HCP 3. Outside community.
- HCP 4. Outside community.
- HCP 5. Outside community.
- HCP 6. Outside community.
- HCP 7. Outside community.
- HCP 8. Outside community.
- HCP 9. Outside community.
- HCP 10. Outside community.
- HCP 11. Outside community.
- HCP 12. Outside community.
- HCP 13. Inconclusive (could be HCP 8, but these samples were collected >14 apart).
- HCP 14. Employee source (HCP 8 or HCP 13).

Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	October 2020	hCoV-19/USA/WI-UW-2283/2020	20G	B.1.370
HCP 2	October 2020	hCoV-19/USA/WI-UW-2284/2020	20G	B.1.2
HCP 3	October 2020	hCoV-19/USA/WI-UW-2285/2020	20A	B.1.216
HCP 4	October 2020	hCoV-19/USA/WI-UW-2286/2020	20G	B.1.2
HCP 5	October 2020	hCoV-19/USA/WI-UW-2390/2020	20G	B.1.2
HCP 6	October 2020	hCoV-19/USA/WI-UW-2391/2020	20G	B.1
HCP 7	November 2020	hCoV-19/USA/WI-UW-2287/2020	20G	B.1.2
HCP 8	November 2020	hCoV-19/USA/WI-UW-2288/2020	20G	B.1.2
HCP 9	November 2020	hCoV-19/USA/WI-UW-2289/2020	20G	B.1.2
HCP 10	November 2020	hCoV-19/USA/WI-UW-2313/2020	20G	B.1.2
HCP 11	November 2020	hCoV-19/USA/WI-UW-2290/2020	20G	B.1.2
HCP 12	November 2020	hCoV-19/USA/WI-UW-2291/2020	20G	B.1.2
HCP 13	November 2020	hCoV-19/USA/WI-UW-2292/2020	20G	B.1.2
HCP 14	November 2020	hCoV-19/USA/WI-UW-2293/2020	20G	B.1.2
patient A	November 2020	hCoV-19/USA/WI-UW-2325/2020	20B	B.1.1.73


Epidemiological information

HCP 1-14 work in the same department together. One or more of these HCP provided direct patient care to patient A.

Alignment

MN908947.3 (Se	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	2	29,903
HCP_1								1.1								
HCP_2						1 1					. '					
HCP_3								1 1			1				1 1111	
HCP_4	1			1	1			1.1				11	1	1.1	1 1 1	1
HCP 5		1						1.1		316			1			1
HCP_6		1	1 1			1	1	1					1.1	1	1 1	11
HCP_6 HCP_7 HCP_8					1			1.1					1	1.1	1 1 1	
HCP 8	1		1		1	1		1.1		1		1	1	1.1	1 1 1	
HCP_9	1	1	1	1		1		1 1		1		1	1		1 1 1	1
HCP_10	-		1	1	1	1		1 1		1		1	1	1.1	1 1 1	1
HCP_11						1	1			1						1
HCP_12	1	1	1	1 1	1	1		1.1		1	r	T	1	- F F	1 1 1	_
HCP_13		1	1					1.1		1		1	1	1.1	1 1 1	
HCP_14		1			1	1		1.1				1	1	1.1	1.1.1	
patient_A	-	1.1	11.1			08.51	11.0	1		1	1	0.92	11 1	25-23	1 1	1

Phylogeny



Report #33. 2021-01-03.

Likely source of HCP infection

- HCP 1. Combined patient and employee cluster.
- HCP 2. Combined patient and employee cluster.
- HCP 3. Combined patient and employee cluster.
- HCP 4. Outside community.
- HCP 5. Combined patient and employee cluster.

Patients A, B and C are the patients involved in this combined patient and employee cluster.


Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	December 2020	hCoV-19/USA/WI-UW-2394/2020	20G	B.1.2
HCP 2	December 2020	hCoV-19/USA/WI-UW-2396/2020	20G	B.1.2
HCP 3	December 2020	hCoV-19/USA/WI-UW-2451/2020	20G	B.1.2
HCP 4	December 2020	hCoV-19/USA/WI-UW-2452/2020	20G	B.1.2
HCP 5	December 2020	hCoV-19/USA/WI-UW-2450/2020	20G	B.1.2
patient A	December 2020	hCoV-19/USA/WI-UW-2392/2020	20G	B.1.2
patient B	December 2020	hCoV-19/USA/WI-UW-2393/2020	20G	B.1.2
patient C	December 2020	hCoV-19/USA/IL-UW-2449/2020	20G	B.1.2

Samples

Epidemiological information

HCP 1-5 provided direct care to one or more of these patients, A-C. HCP 1-5 may have also interacted with each other. No lapses in PPE were reported. Patient A had the earliest reported symptom onset.

MN908947.3 (Se	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,	000	26,000			29,903
HCP 1			1		1	1	1	1.1				1	1		11 1	-	1.1	
HCP_2	-	1	1		1	1	1	1.1		1.1		1	1	1	II I	1	1.1	н
HCP_3		1	1		1	1	T.	1.1		1.1			1	1	11 1	1	1.1	
HCP_4	_	1	1		1	1	1	1.1		1.1		1		1	- U - I	1	1	п
HCP_5		1					1	1.1		11		8	1	2	11 1	1	1.1	
Patient A	-	1	1		1	1	1	1.1		1.1		1	1	- 1	11 1	1	1.1	
Patient_B	-	1	1		1	1	1	1.1		11		1	1	1	11 1	1	1.1	н
Patient_C	-	1	1		1	1	I.	1.1		1.1		1	1	1	11 1	1	1.1	н



Report #34. 2021-01-03.

Likely source of HCP infection

HCP 1. Outside community.

Samples

Sample type	Sample collection date	GISAID identifier	Clade (Nextstrain)	Lineage (Pangolin)
HCP 1	December 2020	hCoV-19/USA/WI-UW-2453/2020	20G	B.1.2
patient A	December 2020	hCoV-19/USA/WI-UW-2395/2020	20G	B.1.2

Epidemiological information

HCP 1 provided direct care to patient A during the 14 days prior to HCP 1's symptom onset. HCP 1 reported no lapse in PPE when providing care to patient A.

MN908947.3 (Se	1	2,000	4,000	6,000	8,000	10,000	12,000	14,000	16,000	18,000	20,000	22,000	24,000	26,000	29,903
HCP_1			I		1	1 1		1.1		1.1	1 1	1	П	1.1	
Patient_A	1	1	1	1 1	1	1		1.1		1		1	1	1.1	1 1 1 1

