

**Supplementary Table 1:** Multivariate linear regression coefficients and exponentiated coefficients for all the variables analyzed.

	Coeff	5%	95%	Exponentiated Coefficients	5%	95%	
Constant	2.98	2.78	3.25	19.65	16.05	25.67	*
IgG levels pre VAX	0.87	0.59	1.13	2.38	1.80	3.10	*
Fever	0.16	-0.28	0.60	1.18	0.76	1.82	
Low-grade Fever	-0.06	-0.45	0.22	0.94	0.64	1.24	
Headache	-0.03	-0.35	0.28	0.97	0.70	1.32	
Cough	0.04	-0.25	0.34	1.04	0.78	1.41	
Sore throat	-0.35	-0.53	-0.11	0.70	0.59	0.89	*
Muscle pain	0.26	-0.08	0.49	1.30	0.92	1.63	
Asthenia	0.18	-0.18	0.48	1.20	0.83	1.61	
Anosmia/dysgeusia	0.50	0.14	0.87	1.65	1.16	2.38	*
Gastrointestinal disorders	0.59	0.16	0.97	1.81	1.17	2.63	*
Conjunctivitis	-0.18	-0.48	0.17	0.84	0.62	1.19	
Dyspnea	0.17	-0.10	0.48	1.19	0.90	1.61	
Chest pain	0.11	-0.14	0.38	1.12	0.87	1.46	
Tachycardia	0.26	0.02	0.60	1.30	1.02	1.83	*
Pneumonia	-0.11	-0.48	0.23	0.90	0.62	1.26	
Age	-0.13	-0.24	-0.001	0.88	0.79	1.00	*
Sex	-0.01	-0.16	0.19	0.99	0.85	1.21	
Serology days from VAX	0.87	0.65	1.03	2.39	1.91	2.80	*
Serology days before VAX	-0.003	-0.01	0.01	1.00	0.99	1.01	
COVID-19	1.48	1.07	1.93	4.39	2.92	6.92	*

\*  $p<0.05$