

Variable	Index cases (n=50)	Household contacts (n=137)
Age median [IQR]	46 [37-54]	21 [13-46]
<b>Age groups n(%)</b>		
0-12 years	-	33 (24)
13-18 years	-	23 (17)
19-49 years	31 (62)	50 (36)
50+ years	19 (28)	31 (23)
<b>Sex Female n(%)</b>	38 (76)	54 (39)
<b>Diagnosis n(%)</b>		
PCR+	46 (92)	48 (35)
PCR- Sero+	3 (6)	22 (15)
PCR- Sero- MLF+	1 (2)	5 (4)

**Table S1 | General characteristics of study participants.**

	Index cases (n=50)	Household cases (n=75)
<b>PCR +</b>	<b>46 (92)</b>	<b>48 (64)</b>
PCR+ sero+ MLF+	43 (86)	42 (56)
PCR+ sero+ MLF-	2 (4)	1 (1)
PCR+ sero- MLF+	0	1 (1)
PCR+ sero- MLF-	1 (2)	4 (5)
<b>Sero +</b>	<b>48 (96)</b>	<b>65 (87)</b>
PCR- sero+ MLF+	3 (6)	9 (12)
PCR- sero+ MLF-	0	13 (17)
<b>MLF +</b>	<b>46 (92)</b>	<b>57 (76)</b>
PCR- sero- MLF+	1 (2)	5 (7)

**Table S2 | PCR, serum and mucosal positivity against SARS-CoV-2 in index and household cases.** The PCR positive threshold was set at a Ct value <36, corresponding to a viral load of  $\geq 10^3$  copies/ml of extracted sample. The seroconversion threshold was based on the mean + 2\*SD of the log<sub>2</sub>-transformed pre-SARS-CoV-2 control samples. To identify cases based on mucosal antibodies against S, a naive Bayes model was used, trained by the mucosal antibody measurements of the PCR positive study participants (cases) and the pre-SARS-CoV-2 control MLF samples (controls).