

Table S1. Participant characteristics based on observed data

	Total (n=10,619)	Boys (n=5340)	Girls (n=5279)
Age (y)	14.3 ± 0.34	14.3 ± 0.34	14.2 ± 0.34
Ethnicity			
White	8439 (79.5)	4239 (79.4)	4200 (79.5)
South Asian	1027 (9.7)	501 (9.4)	526 (10.0)
Other / Mixed	1153 (10.8)	600 (11.2)	553 (10.5)
Net disposable household income (£/week)	416 ± 178	420 ± 177	413 ± 178
Birth weight (kg)	3.4 ± 0.6	3.4 ± 0.6	3.3 ± 0.6
Breast fed ≥3 months	4253 (40.1)	2122 (39.7)	2131 (40.4)
MVPA (60 min/d)			
Not at all	446 (4.2)	190 (3.6)	256 (4.9)
1-2 days per week	2523 (23.8)	956 (17.9)	1567 (29.7)
3-4 days per week	3606 (34.0)	1694 (31.7)	1912 (36.2)
5-6 days per week	2096 (19.7)	1198 (22.4)	898 (17.0)
Every day	1948 (18.3)	1302 (24.4)	646 (12.2)
TV viewing (hours/d)			
≤1	1375 (13.0)	764 (14.3)	611 (11.6)
1-1.99	2133 (20.1)	1161 (21.7)	972 (18.4)
2-2.99	2579 (24.3)	1265 (23.7)	1314 (24.9)
3-4.99	2508 (23.6)	1216 (22.8)	1292 (24.5)
5-6.99	1151 (10.8)	538 (10.1)	613 (11.6)
≥7	873 (8.2)	396 (7.4)	477 (9.0)
Social media (hours/d)			
None	826 (7.8)	567 (10.6)	259 (4.9)
<0.5	1309 (12.3)	886 (16.6)	423 (8.0)
0.5-0.99	1532 (14.4)	953 (17.9)	579 (11.0)
1-1.99	1811 (17.1)	981 (18.4)	830 (15.7)
2-2.99	1639 (15.4)	760 (14.2)	879 (16.6)
3-4.99	1493 (14.1)	558 (10.4)	935 (17.7)
5-6.99	1013 (9.5)	323 (6.1)	690 (13.1)
≥7	996 (9.4)	312 (5.8)	684 (13.0)
Electronic gaming (hours/d)			
None	1986 (18.7)	304 (5.7)	1682 (31.9)
<0.5	1446 (13.6)	312 (5.8)	1134 (21.5)
0.5-0.99	1246 (11.7)	478 (8.9)	768 (14.6)
1-1.99	1617 (15.2)	941 (17.6)	676 (12.8)
2-2.99	1449 (13.7)	1031 (19.3)	418 (7.9)
3-4.99	1379 (13.0)	1039 (19.5)	340 (6.4)
5-6.99	717 (6.8)	590 (11.1)	127 (2.4)
≥7	779 (7.3)	645 (12.1)	134 (2.5)
Diet index	5.5 ± 2.1	5.4 ± 2.0	5.5 ± 2.2
BMI z-score	0.57 ± 1.21	0.53 ± 1.24	0.61 ± 1.17
Body fat (%)	24.8 ± 8.1	21.3 ± 7.6	28.4 ± 6.9
Weight status			
Normal	6942 (65.4)	3543 (66.4)	3399 (64.4)
Overweight	1588 (14.9)	733 (13.7)	855 (16.2)
Obese	2089 (19.7)	1064 (19.9)	1025 (19.4)

Unweighted sample. Values are *n* (proportions) or mean ± standard deviation. BMI, Body mass index. MVPA, Moderate to vigorous physical activity.

Table S2. Sample characteristics based on survey weights

	Total	Boys	Girls
Sex (boys)	52.3	-	-
Age (y)	14.3 (14.3 to 14.3)	14.3 (14.3 to 14.3)	14.3 (14.3 to 14.3)
Ethnicity			
White	80.8	80.1	81.6
South Asian	6.7	6.6	6.8
Other / Mixed	12.5	13.3	11.6
Net disposable household income (£/week)	399 (388 to 409)	401 (389 to 412)	397 (385 to 408)
Birth weight (kg)	3.3 (3.3 to 3.4)	3.4 (3.4 to 3.4)	3.3 (3.3 to 3.3)
Breast fed ≥3 months	35.4	34.9	35.9
MVPA (60 min/d)			
Not at all	4.9	4.5	5.3
1-2 days per week	23.8	18.3	29.9
3-4 days per week	33.3	31.1	35.7
5-6 days per week	19.1	21.4	16.5
Every day	18.9	24.7	12.6
TV viewing (hours/d)			
≤1	12.5	13.7	11.2
1-1.99	19.6	21.1	17.9
2-2.99	23.7	22.6	24.9
3-4.99	23.5	22.6	24.5
5-6.99	11.5	11.2	11.9
≥7	9.2	8.8	9.6
Social media (hours/d)			
None	7.7	10.5	4.7
<0.5	12.1	16.2	7.6
0.5-0.99	13.7	16.9	10.1
1-1.99	16.7	18.2	15.0
2-2.99	15.2	14.4	16.2
3-4.99	14.1	10.4	18.3
5-6.99	10.0	6.6	13.5
≥7	10.5	6.8	14.6
Electronic gaming (hours/d)			
None	17.4	5.4	30.7
<0.5	13.0	5.3	21.3
0.5-0.99	11.2	8.0	14.7
1-1.99	15.4	17.3	13.2
2-2.99	13.7	18.6	8.3
3-4.99	13.5	20.2	6.2
5-6.99	7.3	11.4	2.8
≥7	8.5	13.8	2.8
Diet index	5.3 (5.2 to 5.4)	5.3 (5.2 to 5.4)	5.3 (5.2 to 5.5)
BMI z-score	0.59 (0.56 to 0.61)	0.55 (0.51 to 0.59)	0.63 (0.58 to 0.67)
Body fat (%)	24.7 (24.4 to 24.9)	21.3 (21.0 to 21.6)	28.4 (28.1 to 28.7)
Weight status			
Normal	64.8	65.8	63.8
Overweight	14.8	13.5	16.2
Obese	20.4	20.7	20.0

Values are proportions or mean (95% confidence interval) based on survey weights. BMI, Body mass index. MVPA, Moderate to vigorous physical activity.

Table S3. Distributions of sleep onset times stratified by sex

	Before 10pm	10–10:59pm	11– 11:59pm	After midnight
Boys	1202 (22.5); 23.1	2194 (41.1); 40.8	1623 (30.4); 29.8	321 (6.0); 6.3
Girls	1236 (23.4); 24.0	2176 (41.2); 40.8	1576 (29.9); 29.4	291 (5.5); 5.8

Values are unweighted sample sizes (proportions); *weighted proportions based on survey weights*. Chi² test on unweighted data: $p=0.50$; Chi² test on weighted data: $p=0.67$

Table S4. Distributions of sleep duration stratified by sex

	≤8 hours	8-9 hours	9-10 hours	>10 hours
Boys	521 (9.8); 10.8	1768 (33.1); 31.6	2244 (42.0); 42.6	807 (15.1); 15.0
Girls	580 (11.0); 11.9	1774 (33.6); 33.8	2228 (42.2); 41.6	697 (13.2); 12.7

Values are unweighted sample sizes (proportions); *weighted proportions based on survey weights*. Chi² test on unweighted data: $p=0.012$; Chi² test on weighted data: $p=0.02$

Table S5. Distributions of sleep latency stratified by sex

	0-15 mins	16-30 mins	31-45 mins	46-60 mins	>60 mins
Boys	1989 (37.3); 37.1	1743 (32.6); 31.6	770 (14.4); 14.8	361 (6.8); 6.4	477 (8.9); 10.1
Girls	1585 (30.0); 29.7	1789 (33.9); 33.4	909 (17.2); 17.6	451 (8.6); 8.5	545 (10.3); 10.8

Values are unweighted sample sizes (proportions); *weighted proportions based on survey weights*. Chi² test on unweighted data: $p<0.001$; Chi² test on weighted data: $p<0.001$

Table S6. Distributions of night waking frequency stratified by sex

	Never	A little	Sometimes	Often	Habitually
Boys	1919 (35.9); 35.5	1868 (35.0); 33.8	769 (14.4); 14.3	345 (6.5); 6.7	439 (8.2); 9.7
Girls	1313 (24.9); 24.1	1814 (34.4); 32.9	914 (17.3); 17.8	526 (9.9); 10.4	712 (13.5); 14.8

Values are unweighted sample sizes (proportions); *weighted proportions based on survey weights*. Chi² test on unweighted data: $p<0.001$; Chi² test on weighted data: $p<0.001$

Table S7. Cross-tabulation of sleep onset time and sleep duration

Boys	≤8 hours	>8 to 9 hours	>9 to 10 hours	>10 hours
Before 10pm	0 (0.0); 0.0	110 (9.1); 8.8	560 (46.6); 47.9	532 (44.3); 43.3
10 – 10:59pm	59 (2.7); 3.0	707 (32.2); 31.7	1170 (53.3); 54.1	258 (11.8); 11.2
11 – 11:59pm	273 (16.8); 19.0	839 (51.7); 48.4	494 (30.4); 30.2	17 (1.1); 1.4
After midnight	189 (58.9); 58.3	112 (34.9); 34.5	20 (6.2); 7.2	0 (0.0); 0.0
Girls	≤8 hours	>8 to 9 hours	>9 to 10 hours	>10 hours
Before 10pm	0 (0.0); 0.0	80 (6.5); 6.7	644 (52.1); 54.2	512 (41.4); 39.1
10 – 10:59pm	64 (2.9); 3.2	763 (35.1); 38.7	1180 (54.2); 51.0	169 (7.8); 7.1
11 – 11:59pm	330 (20.9); 22.4	836 (53.1); 50.6	394 (25.0); 25.8	16 (1.0); 1.2
After midnight	186 (63.9); 70.2	95 (32.7); 26.4	10 (3.4); 3.4	0 (0.0); 0.0

Values are unweighted sample sizes (proportions); *weighted proportions based on survey weights.*

Table S8. Cross-tabulation of sleep onset time and sleep latency

Boys	0-15 mins	16-30 mins	31-45 mins	46-60 mins	>60 mins
Before 10pm	487 (40.5); 40.4	426 (35.4); 35.0	157 (13.1); 12.6	65 (5.4); 5.9	67 (5.6); 6.1
10 – 10:59pm	851 (38.8); 38.4	752 (34.3); 33.6	331 (15.1); 16.2	155 (7.0); 5.9	105 (4.8); 5.9
11 – 11:59pm	568 (35.0); 35.2	508 (31.3); 29.6	247 (15.2); 15.3	110 (6.8); 7.4	190 (11.7); 12.5
After midnight	83 (25.8); 26.0	57 (17.8); 15.5	35 (10.9); 10.5	31 (9.7); 7.3	115 (35.8); 40.7
Girls	0-15 mins	16-30 mins	31-45 mins	46-60 mins	>60 mins
Before 10pm	468 (37.9); 37.6	444 (35.9); 36.0	188 (15.2); 15.5	64 (5.2); 5.2	72 (5.8); 5.7
10 – 10:59pm	683 (31.4); 31.1	813 (37.4); 36.4	391 (18.0); 18.5	171 (7.8); 7.8	118 (5.4); 6.2
11 – 11:59pm	394 (25.0); 24.6	479 (30.4); 30.6	284 (18.0); 17.8	179 (11.4); 11.3	240 (15.2); 15.7
After midnight	40 (13.8); 12.2	53 (18.2); 15.9	46 (15.8); 19.5	37 (12.7); 12.9	115 (39.5); 39.5

Values are unweighted sample sizes (proportions); *weighted proportions based on survey weights.*

Table S9. Cross-tabulation of sleep onset time and night waking frequency

Boys	Never	A little	Sometimes	Often	Habitually
Before 10pm	411 (34.2); 35.3	450 (37.5); 36.5	177 (14.7); 14.6	64 (5.3); 4.7	100 (8.3); 8.9
10 – 10:59pm	816 (37.2); 37.0	784 (35.7); 33.9	317 (14.5); 14.4	138 (6.3); 7.1	139 (6.3); 7.6
11 – 11:59pm	579 (35.6); 34.1	561 (34.6); 34.0	230 (14.2); 14.2	110 (6.8); 6.2	143 (8.8); 11.5
After midnight	113 (35.2); 33.8	73 (22.7); 22.0	45 (14.0); 12.6	33 (10.3); 13.5	57 (17.8); 18.1
Girls	Never	A little	Sometimes	Often	Habitually
Before 10pm	357 (28.9); 29.2	437 (35.4); 33.3	208 (16.8); 17.2	97 (7.8); 7.7	137 (11.1); 12.6
10 – 10:59pm	583 (26.8); 25.6	798 (36.7); 36.6	366 (16.8); 16.5	208 (9.5); 9.9	221 (10.2); 11.4
11 – 11:59pm	328 (20.8); 19.8	500 (31.7); 28.8	294 (18.7); 20.6	185 (11.7); 12.9	269 (17.1); 17.9
After midnight	45 (15.5); 14.6	79 (27.1); 26.0	46 (15.8); 14.2	36 (12.4); 13.6	85 (29.2); 31.6

Values are unweighted sample sizes (proportions); *weighted proportions based on survey weights.*

Table S10. Cross-tabulation of sleep duration and sleep latency

Boys	0-15 mins	16-30 mins	31-45 mins	46-60 mins	>60 mins
≤8 hours	160 (30.7); 28.8	128 (24.6); 22.8	72 (13.8); 14.2	42 (8.1); 9.5	119 (22.8); 24.7
>8 to 9 hours	673 (38.1); 37.8	592 (33.5); 32.8	222 (12.6); 13.1	114 (6.4); 5.5	167 (9.4); 10.8
>9 to 10 hours	875 (39.0); 40.0	730 (32.5); 31.1	354 (15.8); 16.1	143 (6.4); 5.9	142 (6.3); 6.9
>10 hours	281 (34.8); 33.5	293 (36.3); 36.9	122 (15.1); 14.8	62 (7.7); 7.4	49 (6.1); 7.4
Girls	0-15 mins	16-30 mins	31-45 mins	46-60 mins	>60 mins
≤8 hours	120 (20.7); 19.9	134 (23.1); 21.0	92 (15.9); 17.8	83 (14.3); 15.0	151 (26.0); 26.3
>8 to 9 hours	491 (27.7); 26.8	585 (33.0); 33.6	324 (18.2); 18.1	165 (9.3); 9.4	209 (11.8); 12.1
>9 to 10 hours	719 (32.3); 33.0	843 (37.8); 36.6	386 (17.3); 17.9	148 (6.7); 6.4	132 (5.9); 6.1
>10 hours	255 (36.6); 36.0	227 (32.6); 33.8	107 (15.3); 15.5	55 (7.9); 6.8	53 (7.6); 7.9

Values are unweighted sample sizes (proportions); *weighted proportions based on survey weights.*

Table S11. Cross-tabulation of sleep duration and night waking frequency

Boys	Never	A little	Sometimes	Often	Habitually
≤8 hours	155 (29.8); 26.6	150 (28.8); 28.7	86 (16.5); 16.1	46 (8.8); 9.6	84 (16.1); 19.0
>8 to 9 hours	624 (35.3); 35.3	645 (36.5); 35.4	227 (12.8); 12.7	127 (7.2); 7.5	145 (8.2); 9.1
>9 to 10 hours	857 (38.2); 38.4	794 (35.4); 33.9	313 (13.9); 14.1	130 (5.8); 5.9	150 (6.7); 7.7
>10 hours	283 (35.1); 34.5	279 (34.6); 33.8	143 (17.7); 17.0	42 (5.2); 5.0	60 (7.4); 9.7
Girls	Never	A little	Sometimes	Often	Habitually
≤8 hours	93 (16.0); 14.9	153 (26.4); 24.0	99 (17.1); 17.6	76 (13.1); 12.8	159 (27.4); 30.7
>8 to 9 hours	410 (23.1); 22.4	616 (34.7); 32.7	327 (18.4); 18.6	194 (11.0); 12.8	227 (12.8); 13.5
>9 to 10 hours	604 (27.1); 26.2	813 (36.5); 36.3	373 (16.7); 17.4	192 (8.6); 8.3	246 (11.1); 11.8
>10 hours	206 (29.5); 30.3	232 (33.3); 30.7	115 (16.5); 16.8	64 (9.2); 9.4	80 (11.5); 12.8

Values are unweighted sample sizes (proportions); *weighted proportions based on survey weights.*

Table S12. Cross-tabulation of sleep latency and night waking frequency

Boys	Never	A little	Sometimes	Often	Habitually
0-15 mins	935 (47.0); 45.9	683 (34.3); 33.3	191 (9.6); 10.7	61 (3.1); 3.3	119 (6.0); 6.8
16-30 mins	594 (34.1); 33.7	670 (38.4); 38.2	274 (15.7); 14.4	112 (6.4); 7.1	93 (5.4); 6.6
31-45 mins	206 (26.8); 28.6	274 (35.6); 34.5	155 (20.1); 18.4	70 (9.1); 8.9	65 (8.4); 9.6
46-60 mins	81 (22.4); 24.2	120 (33.2); 33.3	71 (19.7); 20.1	40 (11.1); 9.3	49 (13.6); 13.1
>60 mins	103 (21.6); 20.7	121 (25.4); 21.3	78 (16.3); 17.4	62 (13.0); 13.1	113 (23.7); 27.5
Girls	Never	A little	Sometimes	Often	Habitually
0-15 mins	616 (38.9); 37.5	579 (36.5); 34.8	185 (11.7); 12.8	88 (5.5); 6.1	117 (7.4); 8.8
16-30 mins	428 (23.9); 23.3	725 (40.5); 39.3	336 (18.8); 19.6	139 (7.8); 8.1	161 (9.0); 9.7
31-45 mins	162 (17.8); 17.9	290 (31.9); 30.8	195 (21.5); 20.4	133 (14.6); 14.1	129 (14.2); 16.8
46-60 mins	59 (13.1); 12.1	117 (25.9); 24.7	93 (20.6); 23.9	83 (18.4); 17.3	99 (22.0); 22.0
>60 mins	48 (8.8); 9.1	103 (18.9); 17.8	105 (19.3); 16.6	83 (15.2); 18.7	206 (37.8); 37.8

Values are unweighted sample sizes (proportions); *weighted proportions based on survey weights.*

Table S13. Associations of sleep onset time with adiposity – Initial model results

Boys	Before 10pm (n=1202)	10 to 10:59pm (n=2194)	11 to 11:59pm (n=1623)	After midnight (n=321)
BMI z-score	Reference	0.21 (0.10 to 0.33); <0.001	0.29 (0.17 to 0.41); <0.001	0.32 (0.09 to 0.56); 0.007
Body fat (%)	Reference	0.82 (0.17 to 1.47); 0.013	1.14 (0.45 to 1.83); 0.001	1.50 (0.05 to 2.95); 0.043
Girls	Before 10pm (n=1236)	10 to 10:59pm (n=2176)	11 to 11:59pm (n=1576)	After midnight (n=291)
BMI z-score	Reference	0.11 (0.01 to 0.21); 0.032	0.29 (0.18 to 0.40); <0.001	0.33 (0.14 to 0.52); 0.001
Body fat (%)	Reference	0.68 (0.09 to 1.28); 0.024	1.71 (1.06 to 2.35); <0.001	1.96 (0.83 to 3.09); 0.001

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement.

Table S14. Associations of sleep duration with adiposity – Initial model results

Boys	≤8 hours (n=521)	>8-9 hours (n=1768)	>9-10 hours (n=2244)	>10 hours (n=807)
BMI z-score	0.31 (0.13 to 0.49); 0.001	0.18 (0.08 to 0.28); <0.001	Reference	-0.06 (-0.20 to 0.07); 0.35
Body fat (%)	1.65 (0.55 to 2.75); 0.003	0.80 (0.18 to 1.42); 0.012	Reference	-0.11 (-0.91 to 0.70); 0.80
Girls	≤8 hours (n=580)	>8-9 hours (n=1774)	>9-10 hours (n=2228)	>10 hours (n=697)
BMI z-score	0.21 (0.07 to 0.36); 0.005	0.12 (0.03 to 0.21); 0.009	Reference	0.04 (-0.10 to 0.18); 0.55
Body fat (%)	1.32 (0.47 to 2.18); 0.003	0.72 (0.19 to 1.25); 0.008	Reference	0.35 (-0.46 to 1.17); 0.39

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement.

Table S15. Associations of sleep latency with adiposity – Initial model results

Boys	0-15 mins (n=1989)	16-30 mins (n=1743)	31-45 mins (n=770)	46-60 mins (n=361)	>60 mins (n=477)
BMI z-score	0.10 (0.00 to 0.21); 0.047	Reference	0.05 (-0.09 to 0.19); 0.46	-0.11 (-0.34 to 0.11); 0.33	0.04 (-0.15 to 0.23); 0.66
Body fat (%)	0.52 (-0.11 to 1.14); 0.11	Reference	0.39 (-0.45 to 1.23); 0.37	-0.84 (-2.18 to 0.50); 0.22	0.37 (-0.82 to 1.56); 0.54
Girls	0-15 mins (n=1585)	16-30 mins (n=1789)	31-45 mins (n=909)	46-60 mins (n=451)	>60 mins (n=545)
BMI z-score	0.07 (-0.03 to 0.17); 0.14	Reference	0.10 (-0.02 to 0.22); 0.11	0.34 (0.18 to 0.49); <0.001	0.33 (0.20 to 0.47); <0.001
Body fat (%)	0.43 (-0.15 to 1.02); 0.15	Reference	0.62 (-0.09 to 1.34); 0.088	1.92 (1.02 to 2.83); <0.001	1.82 (1.00 to 2.64); <0.001

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement.

Table S16. Associations of night waking frequency with adiposity – Initial model results

Boys	Never (n=1919)	A little (n=1868)	Sometimes (n=769)	Often (n=345)	Habitually (n=439)
BMI z-score	Reference	0.00 (-0.11 to 0.12); 0.95	0.13 (-0.01 to 0.27); 0.075	0.07 (-0.10 to 0.25); 0.42	0.07 (-0.12 to 0.27); 0.46
Body fat (%)	Reference	0.03 (-0.67 to 0.72); 0.94	0.77 (-0.07 to 1.61); 0.071	0.42 (-0.74 to 1.58); 0.48	0.47 (-0.66 to 1.61); 0.41
Girls	Never (n=1313)	A little (n=1814)	Sometimes (n=914)	Often (n=526)	Habitually (n=712)
BMI z-score	Reference	0.00 (-0.11 to 0.12); 0.98	0.10 (-0.03 to 0.22); 0.14	0.30 (0.15 to 0.45); <0.001	0.20 (0.04 to 0.36); 0.013
Body fat (%)	Reference	-0.04 (-0.71 to 0.63); 0.90	0.60 (-0.13 to 1.32); 0.11	1.75 (0.83 to 2.68); <0.001	1.33 (0.39 to 2.26); 0.006

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement.

Table S17. Associations of sleep onset time with adiposity – adjusted for screen-based behaviours other than TV viewing

Boys	Before 10pm (n=1202)	10 to 10:59pm (n=2194)	11 to 11:59pm (n=1623)	After midnight (n=321)
BMI z-score				
Model 1 + social media use	Reference	0.23 (0.12 to 0.34); <0.001	0.31 (0.18 to 0.44); <0.001	0.36 (0.14 to 0.59); 0.002
Model 2 + social media use	Reference	0.24 (0.12 to 0.35); <0.001	0.33 (0.20 to 0.45); <0.001	0.39 (0.17 to 0.62); 0.001
Model 1 + electronic games	Reference	0.23 (0.11 to 0.35); <0.001	0.32 (0.19 to 0.45); <0.001	0.40 (0.17 to 0.62); 0.001
Model 2 + electronic games	Reference	0.23 (0.12 to 0.35); <0.001	0.33 (0.21 to 0.46); <0.001	0.43 (0.21 to 0.65); <0.001
Body fat (%)				
Model 1 + social media use	Reference	0.92 (0.28 to 1.57); 0.005	1.28 (0.53 to 2.03); 0.001	1.79 (0.38 to 3.19); 0.013
Model 2 + social media use	Reference	0.94 (0.30 to 1.59); 0.004	1.34 (0.58 to 2.09); 0.001	1.89 (0.47 to 3.31); 0.009
Model 1 + electronic games	Reference	0.89 (0.22 to 1.55); 0.009	1.25 (0.52 to 1.98); 0.001	1.88 (0.50 to 3.26); 0.008
Model 2 + electronic games	Reference	0.91 (0.25 to 1.57); 0.007	1.32 (0.60 to 2.04); <0.001	1.98 (0.60 to 3.36); 0.005
Girls	Before 10pm (n=1236)	10 to 10:59pm (n=2176)	11 to 11:59pm (n=1576)	After midnight (n=291)
BMI z-score				
Model 1 + social media use	Reference	0.11 (0.01 to 0.22); 0.030	0.28 (0.16 to 0.40); <0.001	0.32 (0.12 to 0.52); 0.002
Model 2 + social media use	Reference	0.11 (0.01 to 0.22); 0.031	0.24 (0.13 to 0.36); <0.001	0.23 (0.02 to 0.44); 0.029
Model 1 + electronic games	Reference	0.11 (0.01 to 0.21); 0.032	0.29 (0.18 to 0.40); <0.001	0.37 (0.18 to 0.56); <0.001
Model 2 + electronic games	Reference	0.11 (0.01 to 0.21); 0.036	0.25 (0.14 to 0.37); <0.001	0.28 (0.08 to 0.48); 0.007
Body fat (%)				
Model 1 + social media use	Reference	0.72 (0.12 to 1.32); 0.019	1.68 (0.99 to 2.37); <0.001	1.96 (0.77 to 3.16); 0.001
Model 2 + social media use	Reference	0.72 (0.12 to 1.31); 0.019	1.48 (0.78 to 2.17); <0.001	1.47 (0.22 to 2.71); 0.021
Model 1 + electronic games	Reference	0.68 (0.08 to 1.28); 0.026	1.72 (1.07 to 2.38); <0.001	2.17 (1.02 to 3.32); <0.001
Model 2 + electronic games	Reference	0.67 (0.08 to 1.26); 0.027	1.51 (0.84 to 2.18); <0.001	1.68 (0.47 to 2.90); 0.007

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Model 1 results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement, physical activity, diet index, and social media use or electronic games as specified. Model 2 results are further adjusted for sleep latency and night waking frequency.

Table S18. Associations of sleep duration and adiposity – adjusted for screen-based behaviours other than TV viewing

Boys	≤8 hours (n=521)	>8 to 9 hours (n=1768)	>9 to 10 hours (n=2244)	>10 hours (n=807)
BMI z-score				
Model 1 + social media use	0.31 (0.13 to 0.49); 0.001	0.20 (0.10 to 0.29); <0.001	Reference	-0.07 (-0.20 to 0.07); 0.32
Model 2 + social media use	0.32 (0.14 to 0.50); <0.001	0.20 (0.11 to 0.29); <0.001	Reference	-0.06 (-0.20 to 0.07); 0.36
Model 1 + electronic games	0.32 (0.14 to 0.50); <0.001	0.19 (0.10 to 0.29); <0.001	Reference	-0.07 (-0.20 to 0.06); 0.31
Model 2 + electronic games	0.33 (0.16 to 0.51); <0.001	0.20 (0.10 to 0.29); <0.001	Reference	-0.07 (-0.20 to 0.07); 0.34
Body fat (%)				
Model 1 + social media use	1.68 (0.58 to 2.77); 0.003	0.88 (0.29 to 1.47); 0.004	Reference	-0.14 (-0.93 to 0.64); 0.72
Model 2 + social media use	1.71 (0.62 to 2.79); 0.002	0.90 (0.31 to 1.48); 0.003	Reference	-0.11 (-0.90 to 0.67); 0.78
Model 1 + electronic games	1.72 (0.62 to 2.82); 0.002	0.85 (0.26 to 1.45); 0.005	Reference	-0.14 (-0.93 to 0.65); 0.73
Model 2 + electronic games	1.75 (0.66 to 2.83); 0.002	0.87 (0.28 to 1.46); 0.004	Reference	-0.11 (-0.90 to 0.68); 0.78
Girls	≤8 hours (n=580)	>8 to 9 hours (n=1774)	>9 to 10 hours (n=2228)	>10 hours (n=697)
BMI z-score				
Model 1 + social media use	0.19 (0.04 to 0.35); 0.012	0.11 (0.01 to 0.20); 0.023	Reference	0.04 (-0.10 to 0.18); 0.56
Model 2 + social media use	0.11 (-0.04 to 0.26); 0.16	0.08 (-0.01 to 0.17); 0.091	Reference	0.03 (-0.11 to 0.17); 0.68
Model 1 + electronic games	0.22 (0.07 to 0.37); 0.003	0.12 (0.03 to 0.21); 0.013	Reference	0.04 (-0.10 to 0.17); 0.59
Model 2 + electronic games	0.13 (-0.02 to 0.29); 0.078	0.09 (-0.00 to 0.18); 0.059	Reference	0.03 (-0.11 to 0.16); 0.70
Body fat (%)				
Model 1 + social media use	1.19 (0.32 to 2.07); 0.008	0.63 (0.08 to 1.18); 0.025	Reference	0.33 (-0.48 to 1.15); 0.42
Model 2 + social media use	0.70 (-0.18 to 1.58); 0.12	0.47 (-0.08 to 1.02); 0.093	Reference	0.26 (-0.55 to 1.07); 0.53
Model 1 + electronic games	1.34 (0.47 to 2.20); 0.002	0.68 (0.14 to 1.22); 0.014	Reference	0.33 (-0.47 to 1.13); 0.42
Model 2 + electronic games	0.84 (-0.04 to 1.72); 0.060	0.52 (-0.02 to 1.06); 0.060	Reference	0.26 (-0.53 to 1.06); 0.52

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Model 1 results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement, physical activity, diet index, and social media use or electronic games as specified. Model 2 results are further adjusted for sleep latency and night waking frequency.

Table S19. Associations of sleep latency and adiposity – adjusted for screen-based behaviours other than TV viewing

Boys	0-15 mins (n=1989)	16-30 mins (n=1743)	31-45 mins (n=770)	46-60 mins (n=361)	46-60 mins (n=477)
BMI z-score					
Model 1 + social media use	0.12 (0.01 to 0.22); 0.026	Reference	0.04 (-0.09 to 0.18); 0.54	-0.11 (-0.34 to 0.11); 0.33	0.04 (-0.14 to 0.23); 0.64
Model 2 + social media use	0.12 (0.02 to 0.22); 0.022	Reference	0.03 (-0.10 to 0.17); 0.64	-0.14 (-0.37 to 0.09); 0.24	-0.04 (-0.24 to 0.16); 0.66
Model 1 + electronic games	0.10 (0.00 to 0.21); 0.046	Reference	0.03 (-0.10 to 0.17); 0.62	-0.11 (-0.32 to 0.11); 0.34	0.05 (-0.14 to 0.23); 0.62
Model 2 + electronic games	0.11 (0.01 to 0.22); 0.034	Reference	0.02 (-0.11 to 0.16); 0.74	-0.14 (-0.36 to 0.09); 0.23	-0.04 (-0.24 to 0.15); 0.66
Body fat (%)					
Model 1 + social media use	0.57 (-0.05 to 1.19); 0.074	Reference	0.33 (-0.49 to 1.15); 0.43	-0.82 (-2.16 to 0.52); 0.23	0.39 (-0.78 to 1.57); 0.51
Model 2 + social media use	0.61 (-0.01 to 1.24); 0.056	Reference	0.26 (-0.56 to 1.09); 0.53	-0.99 (-2.35 to 0.38); 0.16	-0.09 (-1.35 to 1.18); 0.90
Model 1 + electronic games	0.53 (-0.10 to 1.16); 0.097	Reference	0.29 (-0.52 to 1.11); 0.48	-0.81 (-2.13 to 0.51); 0.23	0.42 (-0.74 to 1.58); 0.48
Model 2 + electronic games	0.58 (-0.05 to 1.21); 0.069	Reference	0.23 (-0.60 to 1.05); 0.59	-1.00 (-2.34 to 0.35); 0.15	-0.06 (-1.31 to 1.19); 0.92
Girls	0-15 mins (n=1585)	16-30 mins (n=1789)	31-45 mins (n=909)	46-60 mins (n=451)	46-60 mins (n=545)
BMI z-score					
Model 1 + social media use	0.08 (-0.02 to 0.18); 0.11	Reference	0.10 (-0.02 to 0.22); 0.12	0.32 (0.17 to 0.47); <0.001	0.33 (0.19 to 0.47); <0.001
Model 2 + social media use	0.09 (-0.01 to 0.20); 0.077	Reference	0.07 (-0.05 to 0.19); 0.27	0.27 (0.12 to 0.42); 0.001	0.25 (0.10 to 0.39); 0.001
Model 1 + electronic games	0.08 (-0.02 to 0.18); 0.12	Reference	0.10 (-0.03 to 0.22); 0.12	0.32 (0.17 to 0.47); <0.001	0.34 (0.20 to 0.47); <0.001
Model 2 + electronic games	0.09 (-0.01 to 0.19); 0.089	Reference	0.07 (-0.06 to 0.19); 0.29	0.26 (0.11 to 0.42); 0.001	0.25 (0.11 to 0.39); 0.001
Body fat (%)					
Model 1 + social media use	0.47 (-0.11 to 1.05); 0.11	Reference	0.61 (-0.10 to 1.32); 0.094	1.84 (0.94 to 2.75); <0.001	1.77 (0.95 to 2.60); <0.001
Model 2 + social media use	0.55 (-0.05 to 1.15); 0.073	Reference	0.42 (-0.30 to 1.14); 0.25	1.49 (0.58 to 2.39); 0.001	1.25 (0.41 to 2.09); 0.004
Model 1 + electronic games	0.44 (-0.14 to 1.03); 0.13	Reference	0.58 (-0.13 to 1.30); 0.11	1.84 (0.94 to 2.73); <0.001	1.79 (0.97 to 2.60); <0.001
Model 2 + electronic games	0.51 (-0.09 to 1.12); 0.095	Reference	0.40 (-0.32 to 1.11); 0.28	1.47 (0.57 to 2.36); 0.001	1.23 (0.40 to 2.06); 0.004

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Model 1 results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement, physical activity, diet index, and social media use or electronic games as specified. Model 2 results are further adjusted for sleep duration and night waking frequency.

Table S20. Associations of night waking frequency and adiposity – adjusted for screen-based behaviours other than TV viewing

Boys	Never (n=1919)	A little (n=1868)	Sometimes (n=769)	Often (n=345)	Habitually (n=439)
BMI z-score					
Model 1 + social media use	Reference	-0.01 (-0.12 to 0.10); 0.91	0.12 (-0.02 to 0.26); 0.092	0.07 (-0.10 to 0.24); 0.42	0.09 (-0.10 to 0.28); 0.36
Model 2 + social media use	Reference	0.00 (-0.11 to 0.11); 0.95	0.15 (0.01 to 0.29); 0.038	0.09 (-0.09 to 0.26); 0.33	0.10 (-0.09 to 0.29); 0.31
Model 1 + electronic games	Reference	-0.01 (-0.12 to 0.10); 0.90	0.12 (-0.02 to 0.26); 0.085	0.08 (-0.10 to 0.25); 0.39	0.10 (-0.09 to 0.28); 0.31
Model 2 + electronic games	Reference	0.00 (-0.11 to 0.11); 0.99	0.15 (0.01 to 0.29); 0.038	0.09 (-0.09 to 0.27); 0.31	0.10 (-0.09 to 0.30); 0.29
Body fat (%)					
Model 1 + social media use	Reference	-0.03 (-0.71 to 0.65); 0.94	0.76 (-0.07 to 1.60); 0.074	0.45 (-0.68 to 1.58); 0.43	0.60 (-0.50 to 1.69); 0.29
Model 2 + social media use	Reference	0.02 (-0.65 to 0.69); 0.95	0.90 (0.04 to 1.75); 0.039	0.52 (-0.61 to 1.66); 0.37	0.62 (-0.50 to 1.73); 0.28
Model 1 + electronic games	Reference	-0.05 (-0.73 to 0.63); 0.88	0.75 (-0.09 to 1.58); 0.080	0.45 (-0.69 to 1.59); 0.44	0.61 (-0.47 to 1.70); 0.27
Model 2 + electronic games	Reference	-0.01 (-0.68 to 0.66); 0.97	0.87 (0.02 to 1.73); 0.045	0.52 (-0.62 to 1.66); 0.37	0.62 (-0.50 to 1.74); 0.28
Girls	Never (n=1313)	A little (n=1814)	Sometimes (n=914)	Often (n=526)	Habitually (n=712)
BMI z-score					
Model 1 + social media use	Reference	-0.01 (-0.12 to 0.11); 0.91	0.09 (-0.04 to 0.21); 0.16	0.30 (0.14 to 0.45); <0.001	0.19 (0.03 to 0.36); 0.021
Model 2 + social media use	Reference	-0.01 (-0.13 to 0.11); 0.92	0.07 (-0.06 to 0.20); 0.30	0.24 (0.08 to 0.41); 0.004	0.12 (-0.05 to 0.29); 0.16
Model 1 + electronic games	Reference	-0.01 (-0.13 to 0.10); 0.84	0.07 (-0.05 to 0.20); 0.25	0.29 (0.13 to 0.44); <0.001	0.20 (0.03 to 0.37); 0.019
Model 2 + electronic games	Reference	-0.01 (-0.13 to 0.11); 0.84	0.05 (-0.08 to 0.18); 0.44	0.23 (0.06 to 0.40); 0.008	0.12 (-0.05 to 0.30); 0.16
Body fat (%)					
Model 1 + social media use	Reference	-0.10 (-0.77 to 0.57); 0.77	0.56 (-0.16 to 1.27); 0.13	1.74 (0.80 to 2.67); <0.001	1.25 (0.28 to 2.21); 0.011
Model 2 + social media use	Reference	-0.09 (-0.78 to 0.60); 0.80	0.45 (-0.29 to 1.20); 0.24	1.46 (0.46 to 2.46); 0.004	0.88 (-0.14 to 1.89); 0.090
Model 1 + electronic games	Reference	-0.12 (-0.80 to 0.55); 0.72	0.46 (-0.26 to 1.19); 0.21	1.65 (0.69 to 2.61); 0.001	1.28 (0.29 to 2.26); 0.011
Model 2 + electronic games	Reference	-0.12 (-0.82 to 0.58); 0.73	0.35 (-0.40 to 1.10); 0.36	1.34 (0.32 to 2.37); 0.010	0.88 (-0.15 to 1.92); 0.093

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Model 1 results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement, physical activity, diet index, and social media use or electronic games as specified. Model 2 results are further adjusted for sleep duration and sleep latency.

Table S21. Associations of sleep onset time with percent body fat estimated by bioelectrical impedance analysis

Boys	Before 10pm (<i>n</i> =1187)	10 to 10:59pm (<i>n</i> =2168)	11 to 11:59pm (<i>n</i> =1595)	After midnight (<i>n</i> =315)
Log body fat				
Initial model	Reference	0.06 (0.02 to 0.10); 0.006	0.07 (0.02 to 0.11); 0.004	0.12 (0.03 to 0.20); 0.011
Model 1	Reference	0.07 (0.02 to 0.11); 0.002	0.07 (0.03 to 0.12); 0.002	0.12 (0.04 to 0.21); 0.006
Model 2	Reference	0.07 (0.02 to 0.11); 0.002	0.08 (0.03 to 0.12); 0.001	0.13 (0.04 to 0.21); 0.004
Girls	Before 10pm (<i>n</i> =1218)	10 to 10:59pm (<i>n</i> =2152)	11 to 11:59pm (<i>n</i> =1560)	After midnight (<i>n</i> =289)
Body fat (%)				
Initial model	Reference	0.59 (0.00 to 1.19); 0.050	1.68 (1.02 to 2.34); <0.001	1.96 (0.74 to 3.18); 0.002
Model 1	Reference	0.50 (-0.11 to 1.10); 0.11	1.48 (0.81 to 2.16); <0.001	1.71 (0.47 to 2.95); 0.007
Model 2	Reference	0.48 (-0.12 to 1.09); 0.11	1.27 (0.58 to 1.95); <0.001	1.23 (-0.05 to 2.51); 0.059

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Initial model results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement. Model 1 results are further adjusted for physical activity, TV viewing, diet index. Model 2 results are further adjusted for sleep latency and night waking frequency. Due to skewness results for boys are based on log transformed percent body fat.

Table S22. Associations of sleep duration with percent body fat estimated by bioelectrical impedance analysis

Boys	≤8 hours (<i>n</i> =515)	>8 to 9 hours (<i>n</i> =1747)	>9 to 10 hours (<i>n</i> =2211)	>10 hours (<i>n</i> =792)
Log body fat				
Initial model	0.07 (0.01 to 0.14); 0.028	0.06 (0.02 to 0.10); 0.002	Reference	0.01 (-0.04 to 0.06); 0.73
Model 1	0.07 (0.00 to 0.13); 0.037	0.06 (0.03 to 0.10); 0.001	Reference	0.01 (-0.04 to 0.05); 0.82
Model 2	0.07 (0.01 to 0.13); 0.035	0.06 (0.03 to 0.10); <0.001	Reference	0.01 (-0.04 to 0.06); 0.74
Girls	≤8 hours (<i>n</i> =576)	>8 to 9 hours (<i>n</i> =1756)	>9 to 10 hours (<i>n</i> =2201)	>10 hours (<i>n</i> =686)
Body fat (%)				
Initial model	1.18 (0.28 to 2.08); 0.010	0.68 (0.14 to 1.22); 0.013	Reference	0.39 (-0.41 to 1.19); 0.34
Model 1	1.07 (0.18 to 1.96); 0.019	0.59 (0.05 to 1.14); 0.033	Reference	0.45 (-0.33 to 1.22); 0.26
Model 2	0.58 (-0.32 to 1.48); 0.20	0.42 (-0.12 to 0.97); 0.13	Reference	0.37 (-0.40 to 1.15); 0.34

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Initial model results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement. Model 1 results are further adjusted for physical activity, TV viewing, diet index. Model 2 results are further adjusted for sleep latency and night waking frequency. Due to skewness results for boys are based on log transformed percent body fat.

Table S23. Associations of sleep latency with percent body fat estimated by bioelectrical impedance analysis

Boys	0-15 mins (<i>n</i> =1964)	16-30 mins (<i>n</i> =1709)	31-45 mins (<i>n</i> =761)	46-60 mins (<i>n</i> =358)	>60 mins (<i>n</i> =473)
Log body fat					
Initial model	0.04 (0.00 to 0.07); 0.049	Reference	0.03 (-0.02 to 0.08); 0.21	-0.05 (-0.13 to 0.02); 0.15	0.04 (-0.03 to 0.11); 0.29
Model 1	0.04 (-0.00 to 0.07); 0.057	Reference	0.02 (-0.02 to 0.07); 0.33	-0.06 (-0.14 to 0.02); 0.12	0.03 (-0.04 to 0.10); 0.40
Model 2	0.04 (0.00 to 0.08); 0.039	Reference	0.02 (-0.03 to 0.07); 0.40	-0.07 (-0.15 to 0.01); 0.080	0.01 (-0.07 to 0.08); 0.86
Girls	0-15 mins (<i>n</i> =1561)	16-30 mins (<i>n</i> =1772)	31-45 mins (<i>n</i> =897)	46-60 mins (<i>n</i> =445)	>60 mins (<i>n</i> =544)
Body fat (%)					
Initial model	0.32 (-0.26 to 0.90); 0.28	Reference	0.38 (-0.32 to 1.08); 0.29	1.87 (0.93 to 2.80); <0.001	1.86 (1.01 to 2.71); <0.001
Model 1	0.41 (-0.18 to 1.00); 0.17	Reference	0.34 (-0.37 to 1.04); 0.35	1.74 (0.82 to 2.66); <0.001	1.69 (0.82 to 2.56); <0.001
Model 2	0.50 (-0.11 to 1.11); 0.11	Reference	0.14 (-0.56 to 0.84); 0.69	1.37 (0.45 to 2.29); 0.004	1.17 (0.29 to 2.06); 0.010

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Initial model results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement. Model 1 results are further adjusted for physical activity, TV viewing, diet index. Model 2 results are further adjusted for sleep duration and night waking frequency. Due to skewness results for boys are based on log transformed percent body fat.

Table S24. Associations of night waking frequency with percent body fat estimated by bioelectrical impedance analysis

Boys	Never (<i>n</i> =1893)	A little (<i>n</i> =1845)	Sometimes (<i>n</i> =756)	Often (<i>n</i> =338)	Habitually (<i>n</i> =433)
Log body fat					
Initial model	Reference	-0.02 (-0.06 to 0.03); 0.44	0.04 (-0.00 to 0.09); 0.068	0.03 (-0.03 to 0.09); 0.37	0.01 (-0.05 to 0.07); 0.76
Model 1	Reference	-0.02 (-0.06 to 0.03); 0.46	0.05 (0.00 to 0.09); 0.047	0.04 (-0.03 to 0.10); 0.25	0.02 (-0.04 to 0.08); 0.51
Model 2	Reference	-0.01 (-0.06 to 0.03); 0.54	0.05 (0.01 to 0.10); 0.020	0.04 (-0.02 to 0.10); 0.17	0.02 (-0.04 to 0.09); 0.46
Girls	Never (<i>n</i> =1295)	A little (<i>n</i> =1792)	Sometimes (<i>n</i> =902)	Often (<i>n</i> =524)	Habitually (<i>n</i> =706)
Body fat (%)					
Initial model	Reference	0.11 (-0.56 to 0.78); 0.75	0.76 (0.04 to 1.49); 0.040	2.02 (1.09 to 2.95); <0.001	1.29 (0.31 to 2.27); 0.010
Model 1	Reference	0.00 (-0.68 to 0.68); 0.99	0.65 (-0.07 to 1.36); 0.075	1.94 (1.00 to 2.88); <0.001	1.25 (0.25 to 2.25); 0.014
Model 2	Reference	0.02 (-0.67 to 0.71); 0.95	0.58 (-0.16 to 1.32); 0.13	1.71 (0.70 to 2.72); 0.001	0.93 (-0.13 to 1.99); 0.084

Sample sizes are unweighted. Results are based on survey weights and are β -coefficients (95% confidence interval); p -values. Statistically significant associations ($p < 0.05$) are highlighted bold. Initial model results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement. Model 1 results are further adjusted for physical activity, TV viewing, diet index. Model 2 results are further adjusted for sleep duration and sleep latency. Due to skewness results for boys are based on log transformed percent body fat.

Table S25. Associations of sleep onset time with weight status

Boys	Before 10pm (Normal weight: <i>n</i> =858 Overweight / obese: <i>n</i> =344)	10-10:59pm (Normal weight: <i>n</i> =1473 Overweight / obese: <i>n</i> =721)	11-11:59pm (Normal weight: <i>n</i> =1015 Overweight / obese: <i>n</i> =608)	After midnight (Normal weight: <i>n</i> =197 Overweight / obese: <i>n</i> =124)
Odds ratios				
Initial model	Reference	1.35 (1.11 to 1.64); 0.003	1.56 (1.27 to 1.91); <0.001	1.61 (1.11 to 2.32); 0.012
Model 1	Reference	1.40 (1.15 to 1.72); 0.001	1.62 (1.30 to 2.01); <0.001	1.71 (1.18 to 2.48); 0.005
Model 2	Reference	1.41 (1.15 to 1.72); 0.001	1.64 (1.32 to 2.04); <0.001	1.76 (1.19 to 2.60); 0.004
Girls	Before 10pm (Normal weight: <i>n</i> =846 Overweight / obese: <i>n</i> =390)	10-10:59pm (Normal weight: <i>n</i> =1444 Overweight / obese: <i>n</i> =732)	11-11:59pm (Normal weight: <i>n</i> =935 Overweight / obese: <i>n</i> =641)	After midnight (Normal weight: <i>n</i> =174 Overweight / obese: <i>n</i> =117)
Odds ratios				
Initial model	Reference	1.12 (0.94 to 1.32); 0.21	1.48 (1.24 to 1.78); <0.001	1.36 (0.92 to 2.03); 0.13
Model 1	Reference	1.10 (0.92 to 1.30); 0.30	1.43 (1.19 to 1.73); <0.001	1.33 (0.89 to 1.99); 0.17
Model 2	Reference	1.09 (0.92 to 1.30); 0.33	1.36 (1.12 to 1.65); 0.002	1.16 (0.76 to 1.78); 0.49

Sample sizes are unweighted. Results are based on survey weights and are odds ratios (95% confidence interval); *p*-values. Statistically significant associations (*p*<0.05) are highlighted bold. Initial model results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement. Model 1 results are further adjusted for physical activity, TV viewing, diet index. Model 2 results are further adjusted for sleep latency and night waking frequency.

Table S26. Associations of sleep duration with weight status

Boys	≤8 hours (Normal weight: <i>n</i> =298 Overweight / obese: <i>n</i> =223)	>8-9 hours (Normal weight: <i>n</i> =1128 Overweight / obese: <i>n</i> =640)	>9-10 hours (Normal weight: <i>n</i> =1544 Overweight / obese: <i>n</i> =700)	>10 hours (Normal weight: <i>n</i> =573 Overweight / obese: <i>n</i> =234)
Odds ratios				
Initial model	1.79 (1.39 to 2.30); <0.001	1.34 (1.12 to 1.59); 0.001	Reference	0.95 (0.77 to 1.16); 0.60
Model 1	1.76 (1.36 to 2.29); <0.001	1.37 (1.15 to 1.62); <0.001	Reference	0.94 (0.77 to 1.15); 0.55
Model 2	1.80 (1.38 to 2.35); <0.001	1.38 (1.17 to 1.63); <0.001	Reference	0.95 (0.77 to 1.16); 0.62
Girls	≤8 hours (Normal weight: <i>n</i> =339 Overweight / obese: <i>n</i> =241)	>8-9 hours (Normal weight: <i>n</i> =1118 Overweight / obese: <i>n</i> =656)	>9-10 hours (Normal weight: <i>n</i> =1487 Overweight / obese: <i>n</i> =741)	>10 hours (Normal weight: <i>n</i> =455 Overweight / obese: <i>n</i> =242)
Odds ratios				
Initial model	1.54 (1.19 to 1.99); 0.001	1.24 (1.05 to 1.46); 0.012	Reference	1.32 (1.07 to 1.63); 0.011
Model 1	1.52 (1.18 to 1.97); 0.001	1.22 (1.03 to 1.44); 0.024	Reference	1.33 (1.07 to 1.65); 0.009
Model 2	1.38 (1.06 to 1.79); 0.016	1.17 (0.99 to 1.39); 0.065	Reference	1.31 (1.06 to 1.62); 0.014

Sample sizes are unweighted. Results are based on survey weights and are odds ratios (95% confidence interval); *p*-values. Statistically significant associations (*p*<0.05) are highlighted bold. Initial model results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement. Model 1 results are further adjusted for physical activity, TV viewing, diet index. Model 2 results are further adjusted for sleep latency and night waking frequency.

Table S27. Associations of sleep latency with weight status

Boys	0-15 mins (Normal weight: <i>n</i> =1275 Overweight / obese: <i>n</i> =714)	16-30 mins (Normal weight: <i>n</i> =1180 Overweight / obese: <i>n</i> =563)	31-45 mins (Normal weight: <i>n</i> =522 Overweight / obese: <i>n</i> =248)	46-60 mins (Normal weight: <i>n</i> =250 Overweight / obese: <i>n</i> =111)	>60 mins (Normal weight: <i>n</i> =316 Overweight / obese: <i>n</i> =161)
Odds ratios					
Initial model	1.15 (0.97 to 1.37); 0.099	Reference	1.14 (0.90 to 1.44); 0.29	0.93 (0.68 to 1.28); 0.67	1.11 (0.83 to 1.48); 0.48
Model 1	1.16 (0.97 to 1.38); 0.098	Reference	1.11 (0.88 to 1.41); 0.37	0.91 (0.65 to 1.25); 0.55	1.08 (0.82 to 1.44); 0.58
Model 2	1.16 (0.97 to 1.38); 0.10	Reference	1.10 (0.86 to 1.41); 0.44	0.86 (0.62 to 1.21); 0.40	0.94 (0.68 to 1.28); 0.69
Girls	0-15 mins (Normal weight: <i>n</i> =1057 Overweight / obese: <i>n</i> =528)	16-30 mins (Normal weight: <i>n</i> =1178 Overweight / obese: <i>n</i> =611)	31-45 mins (Normal weight: <i>n</i> =583 Overweight / obese: <i>n</i> =326)	46-60 mins (Normal weight: <i>n</i> =267 Overweight / obese: <i>n</i> =184)	>60 mins (Normal weight: <i>n</i> =314 Overweight / obese: <i>n</i> =231)
Odds ratios					
Initial model	1.05 (0.88 to 1.26); 0.60	Reference	1.14 (0.93 to 1.41); 0.20	1.53 (1.17 to 2.01); 0.002	1.57 (1.23 to 1.99); <0.001
Model 1	1.07 (0.89 to 1.29); 0.46	Reference	1.14 (0.92 to 1.40); 0.24	1.49 (1.14 to 1.96); 0.004	1.54 (1.20 to 1.96); 0.001
Model 2	1.08 (0.90 to 1.30); 0.42	Reference	1.10 (0.89 to 1.36); 0.37	1.39 (1.05 to 1.83); 0.020	1.39 (1.08 to 1.78); 0.011

Sample sizes are unweighted. Results are based on survey weights and are odds ratios (95% confidence interval); *p*-values. Statistically significant associations (*p*<0.05) are highlighted bold. Initial model results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement. Model 1 results are further adjusted for physical activity, TV viewing, diet index. Model 2 results are further adjusted for sleep duration and night waking frequency.

Table S28. Associations of night waking frequency with weight status

Boys	Never (Normal weight: <i>n</i> =1267 Overweight / obese: <i>n</i> =652)	A little (Normal weight: <i>n</i> =1278 Overweight / obese: <i>n</i> =590)	Sometimes (Normal weight: <i>n</i> =495 Overweight / obese: <i>n</i> =274)	Often (Normal weight: <i>n</i> =233 Overweight / obese: <i>n</i> =112)	Habitually (Normal weight: <i>n</i> =270 Overweight / obese: <i>n</i> =169)
Odds ratios					
Initial model	Reference	0.90 (0.75 to 1.08); 0.26	1.09 (0.86 to 1.39); 0.47	1.00 (0.72 to 1.38); 0.99	1.07 (0.80 to 1.43); 0.63
Model 1	Reference	0.88 (0.73 to 1.06); 0.19	1.09 (0.85 to 1.40); 0.49	1.01 (0.73 to 1.40); 0.96	1.12 (0.84 to 1.49); 0.43
Model 2	Reference	0.88 (0.73 to 1.07); 0.20	1.12 (0.87 to 1.44); 0.38	1.01 (0.72 to 1.41); 0.95	1.11 (0.82 to 1.50); 0.48
Girls	Never (Normal weight: <i>n</i> =871 Overweight / obese: <i>n</i> =442)	A little (Normal weight: <i>n</i> =1218 Overweight / obese: <i>n</i> =596)	Sometimes (Normal weight: <i>n</i> =577 Overweight / obese: <i>n</i> =337)	Often (Normal weight: <i>n</i> =309 Overweight / obese: <i>n</i> =217)	Habitually (Normal weight: <i>n</i> =424 Overweight / obese: <i>n</i> =288)
Odds ratios					
Initial model	Reference	0.98 (0.81 to 1.18); 0.81	1.14 (0.92 to 1.43); 0.23	1.35 (1.03 to 1.76); 0.030	1.20 (0.92 to 1.57); 0.18
Model 1	Reference	0.95 (0.78 to 1.15); 0.60	1.12 (0.90 to 1.39); 0.32	1.32 (1.01 to 1.74); 0.046	1.20 (0.91 to 1.58); 0.20
Model 2	Reference	0.95 (0.78 to 1.15); 0.59	1.07 (0.86 to 1.34); 0.54	1.21 (0.90 to 1.62); 0.21	1.05 (0.79 to 1.40); 0.73

Sample sizes are unweighted. Results are based on survey weights and are odds ratios (95% confidence interval); *p*-values. Statistically significant associations ($p < 0.05$) are highlighted bold. Initial model results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement. Model 1 results are further adjusted for physical activity, TV viewing, diet index. Model 2 results are further adjusted for sleep duration and sleep latency.

Table S29. Associations of sleep onset time with weight status – adjusted for screen-based behaviours other than TV viewing

Boys	Before 10pm (Normal weight: <i>n</i> =858 Overweight / obese: <i>n</i> =344)	10-10:59pm (Normal weight: <i>n</i> =1473 Overweight / obese: <i>n</i> =721)	11-11:59pm (Normal weight: <i>n</i> =1015 Overweight / obese: <i>n</i> =608)	After midnight (Normal weight: <i>n</i> =197 Overweight / obese: <i>n</i> =124)
Odds ratios				
Model 1 + social media use	Reference	1.42 (1.16 to 1.74); 0.001	1.66 (1.32 to 2.07); <0.001	1.77 (1.23 to 2.56); 0.002
Model 2 + social media use	Reference	1.43 (1.16 to 1.75); 0.001	1.68 (1.34 to 2.10); <0.001	1.82 (1.23 to 2.69); 0.003
Model 1 + electronic games	Reference	1.41 (1.15 to 1.72); 0.001	1.65 (1.33 to 2.04); <0.001	1.80 (1.26 to 2.59); 0.001
Model 2 + electronic games	Reference	1.41 (1.16 to 1.72); 0.001	1.67 (1.35 to 2.07); <0.001	1.85 (1.27 to 2.70); 0.001
Girls	Before 10pm (Normal weight: <i>n</i> =846 Overweight / obese: <i>n</i> =390)	10-10:59pm (Normal weight: <i>n</i> =1444 Overweight / obese: <i>n</i> =732)	11-11:59pm (Normal weight: <i>n</i> =935 Overweight / obese: <i>n</i> =641)	After midnight (Normal weight: <i>n</i> =174 Overweight / obese: <i>n</i> =117)
Odds ratios				
Model 1 + social media use	Reference	1.15 (0.96 to 1.37); 0.12	1.53 (1.25 to 1.87); <0.001	1.46 (0.97 to 2.21); 0.073
Model 2 + social media use	Reference	1.14 (0.96 to 1.36); 0.14	1.45 (1.18 to 1.78); <0.001	1.26 (0.82 to 1.95); 0.29
Model 1 + electronic games	Reference	1.12 (0.94 to 1.33); 0.22	1.49 (1.24 to 1.80); <0.001	1.48 (0.97 to 2.25); 0.066
Model 2 + electronic games	Reference	1.11 (0.93 to 1.32); 0.25	1.41 (1.16 to 1.71); 0.001	1.28 (0.83 to 1.99); 0.27

Sample sizes are unweighted. Results are based on survey weights and are odds ratios (95% confidence interval); *p*-values. Statistically significant associations (*p*<0.05) are highlighted bold. Model 1 results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement, physical activity, diet index, and social media use or electronic games as specified. Model 2 results are further adjusted for sleep latency and night waking frequency.

Table S30. Associations of sleep duration with weight status – adjusted for screen-based behaviours other than TV viewing

Boys	≤8 hours (Normal weight: <i>n</i> =298 Overweight / obese: <i>n</i> =223)	>8-9 hours (Normal weight: <i>n</i> =1128 Overweight / obese: <i>n</i> =640)	>9-10 hours (Normal weight: <i>n</i> =1544 Overweight / obese: <i>n</i> =700)	>10 hours (Normal weight: <i>n</i> =573 Overweight / obese: <i>n</i> =234)
Odds ratios				
Model 1 + social media use	1.81 (1.40 to 2.35); <0.001	1.38 (1.16 to 1.64); <0.001	Reference	0.94 (0.76 to 1.15); 0.54
Model 2 + social media use	1.85 (1.42 to 2.40); <0.001	1.39 (1.17 to 1.65); <0.001	Reference	0.95 (0.77 to 1.16); 0.61
Model 1 + electronic games	1.82 (1.41 to 2.36); <0.001	1.37 (1.15 to 1.62); <0.001	Reference	0.94 (0.76 to 1.15); 0.52
Model 2 + electronic games	1.86 (1.43 to 2.41); <0.001	1.38 (1.17 to 1.64); <0.001	Reference	0.95 (0.77 to 1.16); 0.59
Girls	≤8 hours (Normal weight: <i>n</i> =339 Overweight / obese: <i>n</i> =241)	>8-9 hours (Normal weight: <i>n</i> =1118 Overweight / obese: <i>n</i> =656)	>9-10 hours (Normal weight: <i>n</i> =1487 Overweight / obese: <i>n</i> =741)	>10 hours (Normal weight: <i>n</i> =455 Overweight / obese: <i>n</i> =242)
Odds ratios				
Model 1 + social media use	1.55 (1.20 to 2.1); 0.001	1.23 (1.04 to 1.45); 0.017	Reference	1.28 (1.03 to 1.59); 0.024
Model 2 + social media use	1.40 (1.07 to 1.82); 0.013	1.18 (1.00 to 1.40); 0.050	Reference	1.26 (1.02 to 1.57); 0.035
Model 1 + electronic games	1.57 (1.22 to 2.03); 0.001	1.23 (1.04 to 1.45); 0.018	Reference	1.30 (1.05 to 1.61); 0.016
Model 2 + electronic games	1.41 (1.08 to 1.84); 0.010	1.18 (1.00 to 1.40); 0.052	Reference	1.28 (1.03 to 1.59); 0.023

Sample sizes are unweighted. Results are based on survey weights and are odds ratios (95% confidence interval); *p*-values. Statistically significant associations (*p*<0.05) are highlighted bold. Model 1 results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement, physical activity, diet index, and social media use or electronic games as specified. Model 2 results are further adjusted for sleep latency and night waking frequency.

Table S31. Associations of sleep latency with weight status – adjusted for screen-based behaviours other than TV viewing

Boys	0-15 mins (Normal weight: <i>n</i> =1275 Overweight / obese: <i>n</i> =714)	16-30 mins (Normal weight: <i>n</i> =1180 Overweight / obese: <i>n</i> =563)	31-45 mins (Normal weight: <i>n</i> =522 Overweight / obese: <i>n</i> =248)	46-60 mins (Normal weight: <i>n</i> =250 Overweight / obese: <i>n</i> =111)	>60 mins (Normal weight: <i>n</i> =316 Overweight / obese: <i>n</i> =161)
Odds ratios					
Model 1 + social media use	1.18 (0.99 to 1.40); 0.068	Reference	1.11 (0.88 to 1.41); 0.38	0.93 (0.67 to 1.28); 0.65	1.11 (0.84 to 1.48); 0.46
Model 2 + social media use	1.17 (0.98 to 1.40); 0.079	Reference	1.10 (0.86 to 1.40); 0.45	0.89 (0.63 to 1.24); 0.48	0.96 (0.70 to 1.31); 0.78
Model 1 + electronic games	1.16 (0.98 to 1.38); 0.092	Reference	1.11 (0.87 to 1.40); 0.40	0.93 (0.68 to 1.28); 0.67	1.11 (0.84 to 1.48); 0.47
Model 2 + electronic games	1.16 (0.97 to 1.38); 0.096	Reference	1.09 (0.86 to 1.39); 0.47	0.89 (0.64 to 1.23); 0.48	0.95 (0.70 to 1.31); 0.77
Girls	0-15 mins (Normal weight: <i>n</i> =1057 Overweight / obese: <i>n</i> =528)	16-30 mins (Normal weight: <i>n</i> =1178 Overweight / obese: <i>n</i> =611)	31-45 mins (Normal weight: <i>n</i> =583 Overweight / obese: <i>n</i> =326)	46-60 mins (Normal weight: <i>n</i> =267 Overweight / obese: <i>n</i> =184)	>60 mins (Normal weight: <i>n</i> =314 Overweight / obese: <i>n</i> =231)
Odds ratios					
Model 1 + social media use	1.06 (0.88 to 1.26); 0.55	Reference	1.15 (0.93 to 1.41); 0.21	1.51 (1.15 to 1.98); 0.003	1.58 (1.24 to 2.01); <0.001
Model 2 + social media use	1.06 (0.89 to 1.28); 0.51	Reference	1.11 (0.90 to 1.37); 0.33	1.40 (1.06 to 1.86); 0.017	1.43 (1.11 to 1.84); 0.005
Model 1 + electronic games	1.06 (0.88 to 1.27); 0.54	Reference	1.14 (0.93 to 1.42); 0.21	1.50 (1.15 to 1.97); 0.003	1.59 (1.24 to 2.03); <0.001
Model 2 + electronic games	1.06 (0.88 to 1.27); 0.52	Reference	1.11 (0.90 to 1.37); 0.33	1.40 (1.06 to 1.84); 0.017	1.43 (1.12 to 1.84); 0.005

Sample sizes are unweighted. Results are based on survey weights and are odds ratios (95% confidence interval); *p*-values. Statistically significant associations (*p*<0.05) are highlighted bold. Model 1 results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement, physical activity, diet index, and social media use or electronic games as specified. Model 2 results are further adjusted for sleep duration and night waking frequency.

Table S32. Associations of night waking frequency with weight status – adjusted for screen-based behaviours other than TV viewing

Boys	Never (Normal weight: <i>n</i> =1267 Overweight / obese: <i>n</i> =652)	A little (Normal weight: <i>n</i> =1278 Overweight / obese: <i>n</i> =590)	Sometimes (Normal weight: <i>n</i> =495 Overweight / obese: <i>n</i> =274)	Often (Normal weight: <i>n</i> =233 Overweight / obese: <i>n</i> =112)	Habitually (Normal weight: <i>n</i> =270 Overweight / obese: <i>n</i> =169)
Odds ratios					
Model 1 + social media use	Reference	0.88 (0.73 to 1.06); 0.19	1.09 (0.85 to 1.40); 0.50	1.00 (0.72 to 1.39); 0.99	1.13 (0.85 to 1.51); 0.40
Model 2 + social media use	Reference	0.89 (0.73 to 1.07); 0.21	1.12 (0.87 to 1.45); 0.38	1.00 (0.72 to 1.40); 0.99	1.12 (0.83 to 1.51); 0.47
Model 1 + electronic games	Reference	0.88 (0.73 to 1.06); 0.19	1.09 (0.84 to 1.41); 0.51	1.01 (0.73 to 1.41); 0.93	1.13 (0.85 to 1.50); 0.39
Model 2 + electronic games	Reference	0.88 (0.73 to 1.07); 0.20	1.11 (0.86 to 1.44); 0.41	1.01 (0.72 to 1.41); 0.94	1.12 (0.83 to 1.50); 0.47
Girls	Never (Normal weight: <i>n</i> =871 Overweight / obese: <i>n</i> =442)	A little (Normal weight: <i>n</i> =1218 Overweight / obese: <i>n</i> =596)	Sometimes (Normal weight: <i>n</i> =577 Overweight / obese: <i>n</i> =337)	Often (Normal weight: <i>n</i> =309 Overweight / obese: <i>n</i> =217)	Habitually (Normal weight: <i>n</i> =424 Overweight / obese: <i>n</i> =288)
Odds ratios					
Model 1 + social media use	Reference	0.96 (0.79 to 1.17); 0.71	1.13 (0.91 to 1.41); 0.27	1.36 (1.04 to 1.80); 0.026	1.19 (0.90 to 1.58); 0.22
Model 2 + social media use	Reference	0.96 (0.79 to 1.17); 0.67	1.08 (0.86 to 1.35); 0.52	1.23 (0.92 to 1.65); 0.16	1.03 (0.77 to 1.38); 0.83
Model 1 + electronic games	Reference	0.95 (0.78 to 1.15); 0.60	1.10 (0.88 to 1.37); 0.41	1.31 (1.00 to 1.73); 0.054	1.20 (0.90 to 1.59); 0.21
Model 2 + electronic games	Reference	0.94 (0.77 to 1.15); 0.56	1.04 (0.83 to 1.31); 0.71	1.18 (0.88 to 1.58); 0.27	1.04 (0.78 to 1.39); 0.80

Sample sizes are unweighted. Results are based on survey weights and are odds ratios (95% confidence interval); *p*-values. Statistically significant associations (*p*<0.05) are highlighted bold. Model 1 results are adjusted for age, ethnicity, net disposal household income, birth weight, breast feeding history, season of measurement, physical activity, diet index, and social media use or electronic games as specified. Model 2 results are further adjusted for sleep duration and sleep latency.