

Table e-1. Baseline neuropsychiatric inventory scores according to clinical AD stage.

	Preclinical AD (n=113)			MCI due to AD (n=321)			AD dementia (n=1,090)		
	Pre	Fre	Sev	Pre	Fre	Sev	Pre	Fre	Sev
Apathy	42.5%	1.0 (1.3)	0.7 (0.9)	49.2%	1.1 (1.2)	0.8 (0.9)	60.2% ^{a, e}	1.5 (1.4) ^{a, b}	1.1 (1.0) ^{a, b}
Irritability	53.1%	1.2 (1.2)	0.9 (1.0)	44.9%	1.0 (1.2)	0.8 (0.9)	42.9%	0.9 (1.2)	0.7 (1.0)
Depression	31.0%	0.5 (0.9)	0.4 (0.7)	26.5%	0.5 (1.0)	0.4 (0.8)	32.1%	0.7 (1.1)	0.5 (0.8)
Anxiety	21.2%	0.4 (0.9)	0.3 (0.6)	23.7%	0.4 (0.8)	0.4 (0.8)	31.6% ^b	0.7 (1.1) ^{c, d}	0.5 (0.9) ^{d, e}
Eating behaviors	21.2%	0.4 ± 0.9	0.3 ± 0.7	20.1%	0.4 ± 0.9	0.3 ± 0.7	31.4% ^e	0.7 ± 1.1 ^b	0.5 ± 0.8 ^b
Night-time behaviors	30.1%	0.7 ± 1.1	0.5 ± 0.8	20.1%	0.4 ± 0.9	0.3 ± 0.7	20.0%	0.5 ± 1.0	0.3 ± 0.7
Agitation	14.2 %	0.3 (0.7)	0.3 (0.7)	15.3%	0.3 (0.9)	0.3 (0.7)	16.2%	0.3 (0.9)	0.3 (0.7)
AMB	8.0%	0.2 ± 0.7	0.1 ± 0.4	12.1%	0.3 ± 0.8	0.2 ± 0.5	18.9% ^{d, e}	0.4 ± 1.0 ^{d, e}	0.3 ± 0.7 ^{d, e}
Disinhibition	9.7%	0.2 (0.7)	0.2 (0.6)	13.7%	0.2 ± 0.6	0.2 ± 0.6	12.4%	0.2 ± 0.6	0.2 ± 0.6
Delusions	3.5%	0.1 ± 0.4	0.1 ± 0.4	4.7%	0.1 ± 0.5	0.1 ± 0.3	9.3% ^{d, e}	0.2 ± 0.7 ^e	0.2 ± 0.5 ^e
Euphoria	4.4%	0.1 (0.3)	0.1 (0.4)	5.3%	0.1 (0.5)	0.1 (0.4)	6.0%	0.1 (0.4)	0.1 (0.3)
Hallucinations	5.3%	0.1 (0.6)	0.1 (1.1)	1.2%	0.0 ± 0.2	0.0 ± 0.2	4.6% ^e	0.1 ± 0.4	0.1 ± 0.3
Overall	81.4%	5.1 ± 4.6	3.8 ± 3.6	81.2%	4.8 ± 4.5	3.8 ± 3.6	88.7% ^e	6.3 ± 5.1 ^{b, d}	4.8 ± 3.9 ^{b, d}

Notes. AD = Alzheimer's disease, MCI = mild cognitive impairment, Pre = prevalence (NPI frequency \times severity scores ≥ 1), Fre = NPI frequency score (range 0–4), Sev = NPI severity score (range 0–3), AMB = aberrant motor behaviors. Differences in prevalence rates tested using Chi-Square Tests, differences in frequency and severity scores tested using Kruskal-Wallis Tests.

^a preclinical AD < AD dementia, $p < 0.001$ after correcting for false discovery rate.

^b MCI due to AD < AD dementia, $p < 0.001$ after correcting for false discovery rate.

^c MCI due to AD < AD dementia, $p < 0.01$ after correcting for false discovery rate.

^d preclinical AD < AD dementia, $p < 0.05$ after correcting for false discovery rate.

^e MCI due to AD < AD dementia, $p < 0.05$ after correcting for false discovery rate.

Table e-2. Within-subject and between-subject variances of cognitive and neuropsychiatric measures over time according to clinical AD stage.

	Preclinical AD at baseline		MCI due to AD at baseline		AD dementia at baseline	
	<i>BS variance</i>	<i>WS variance</i>	<i>BS variance</i>	<i>WS variance</i>	<i>BS variance</i>	<i>WS variance</i>
	<i>(% explained)</i>	<i>(% explained)</i>	<i>(% explained)</i>	<i>(% explained)</i>	<i>(% explained)</i>	<i>(% explained)</i>
Cognitive measures						
MMSE	1.42 (52.40%)	1.29 (47.60%)	3.51 (47.37%)	3.90 (52.63%)	14.55 (54.29%)	12.25 (45.71%)
Memory	0.64 (64.65%)	0.35 (35.35%)	2.09 (59.71%)	1.41 (40.29%)	2.75 (60.04%)	1.83 (36.96%)
Attention	0.48 (64.86%)	0.26 (35.14%)	1.07 (56.91%)	0.81 (43.09%)	10.73 (62.60%)	6.41 (37.40%)
Executive functions	0.61 (78.21%)	0.17 (21.79%)	0.87 (74.36%)	0.30 (25.64%)	2.14 (70.86%)	0.88 (29.14%)
Language	0.19 (61.29%)	0.12 (38.71%)	0.36 (61.02%)	0.23 (38.98%)	1.86 (64.36%)	1.03 (35.64%)
Visuospatial	0.08 (17.78%)	0.37 (82.22%)	0.49 (37.12%)	0.83 (62.88%)	11.72 (73.76%)	4.17 (26.24%)
Average % explained	56.53%	43.47%	56.08%	43.90%	64.32%	35.18%
NPS measures						
NPI total score	36.11 (46.16%)	42.12 (56.84%)	29.70 (37.85%)	48.77 (62.15%)	41.03 (37.25%)	69.12 (62.75%)
NPI-apathy	0.84 (13.93%)	5.19 (86.07%)	1.40 (22.29%)	4.88 (77.71%)	2.24 (20.25%)	8.82 (79.75%)

NPI-depression	0.76 (25.25%)	2.25 (74.75%)	1.45 (31.94%)	3.09 (68.06%)	1.40 (29.60%)	3.33 (70.40%)
NPI-irritability	2.23 (27.23%)	5.96 (72.77%)	1.43 (23.40%)	4.68 (76.60%)	2.16 (30.00%)	5.04 (70.00%)
Average % explained	28.14%	72.61%	28.87%	71.13%	29.28%	70.73%

Abbreviations: BS = between-subject, WS = within-subject, AD = Alzheimer's disease, MCI = mild cognitive impairment, MMSE = Mini-Mental State Examination.

Table e-3. Associations between baseline NPS and concurrent cognitive performance according to clinical AD stage.

Preclinical AD (n=113)						
	MMSE	Memory	Attention	EF	Language	Visuospatial
Agitation	0.22 [-0.25, 0.69]	0.43 [-0.08, 0.93]	-0.02 [-0.51, 0.48]	-0.20 [-0.68, 0.27]	-0.35 [-0.86, 0.16]	0.35 [-0.09, 0.80]
AMB	0.01 [-0.61, 0.63]	0.31 [-0.37, 0.99]	0.32 [-0.32, 0.96]	-0.20 [-0.81, 0.42]	0.11 [-0.59, 0.80]	0.30 [-0.31, 0.92]
Anxiety	0.12 [-0.41, 0.65]	0.05 [-0.46, 0.55]	0.36 [-0.15, 0.86]	0.14 [-0.32, 0.60]	0.05 [-0.47, 0.58]	0.22 [-1.03, 1.46]
Delusions	1.19 [0.06, 2.32] ^a	0.47 [-0.67, 1.61]	1.22 [0.11, 2.33] ^a	1.01 [-0.03, 2.06]	-0.45 [-1.62, 0.72]	1.25 [-0.90, 3.40]
Disinhibition	-0.45 [-0.97, 0.07]	-0.12 [-0.72, 0.47]	-0.21 [-0.78, 0.36]	-0.25 [-0.79, 0.29]	0.02 [-0.58, 0.62]	0.25 [-0.27, 0.78]
Eating behaviors	0.21 [-0.22, 0.63]	0.05 [-0.39, 0.49]	0.05 [-0.38, 0.49]	-0.09 [-0.50, 0.32]	-0.13 [-0.59, 0.32]	0.57 [0.14, 1.01] ^a
Euphoria	0.30 [-0.46, 1.06]	0.35 [-0.47, 1.18]	0.40 [-0.39, 1.20]	0.42 [-0.34, 1.18]	-0.27 [-1.10, 0.56]	0.15 [-0.54, 0.85]
Hallucinations	0.36 [-0.52, 1.23]	0.22 [-0.71, 1.16]	0.24 [-0.67, 1.15]	-0.24 [-1.11, 0.62]	0.19 [-0.75, 1.14]	0.71 [-0.12, 1.55]
Night-time behaviors	0.09 [-0.26, 0.45]	0.42 [0.04, 0.79]	0.25 [-0.11, 0.62]	0.39 [0.04, 0.73] ^a	0.17 [-0.21, 0.55]	-0.15 [-0.50, 0.20]
MCI due to AD (n=321)						
	MMSE	Memory	Attention	EF	Language	Visuospatial
Agitation	0.10 [-0.16, 0.36]	0.11 [-0.17, 0.40]	-0.23 [-0.51, 0.05]	-0.21 [-0.47, 0.06]	-0.02 [-0.32, 0.28]	-0.01 [-0.30, 0.28]

AMB	0.01 [-0.30, 0.31]	-0.10 [-0.44, 0.23]	-0.36 [-0.70, -0.03] ^a	0.00 [-0.32, 0.31]	-0.19 [-0.54, 0.16]	0.17 [-0.19, 0.53]
Anxiety	0.13 [-0.09, 0.35]	0.00 [-0.24, 0.25]	-0.05 [-0.29, 0.19]	-0.02 [-0.25, 0.21]	-0.10 [-0.35, 0.16]	0.04 [-0.21, 0.29]
Delusions	-0.12 [-0.57, 0.32]	-0.51 [-0.99, -0.04] ^a	-0.24 [-0.72, 0.24]	0.00 [-0.45, 0.46]	0.20 [-0.31, 0.71]	0.18 [-0.30, 0.65]
Disinhibition	0.13 [-0.14, 0.41]	0.41 [0.11, 0.71] ^a	-0.20 [-0.50, 0.10]	-0.03 [-0.31, 0.25]	-0.17 [-0.49, 0.15]	0.17 [-0.14, 0.48]
Eating behaviors	0.00 [-0.24, 0.24]	-0.04 [-0.30, 0.22]	-0.21 [-0.47, 0.05]	-0.23 [-0.47, 0.02]	-0.41 [-0.69, -0.14] ^a	-0.10 [-0.38, 0.18]
Euphoria	0.34 [-0.07, 0.74]	0.20 [-0.23, 0.64]	0.08 [-0.35, 0.52]	0.08 [-0.33, 0.50]	0.20 [-0.26, 0.66]	-0.36 [-0.80, 0.08]
Hallucinations	-0.29 [-1.21, 0.62]	-0.62 [-1.55, 0.31]	-1.73 [-2.68, -0.79]^b	-1.16 [-2.04, -0.29] ^a	-0.28 [-1.28, 0.72]	-0.58 [-1.69, 0.53]
Night-time behaviors	0.25 [-0.02, 0.48]	0.11 [-0.14, 0.37]	-0.03 [-0.28, 0.22]	0.06 [-0.18, 0.30]	0.16 [-0.11, 0.43]	0.00 [-0.27, 0.27]

AD dementia (n=1,090)

	MMSE	Memory	Attention	EF	Language	Visuospatial
Agitation	-0.27 [-0.43, -0.11]^b	-0.01 [-0.19, 0.17]	-0.11 [-0.28, 0.05]	-0.17 [-0.34, 0.00]	-0.08 [-0.25, 0.10]	-0.09 [-0.28, 0.10]
AMB	-0.28 [-0.43, -0.13]^b	-0.16 [-0.33, 0.01]	-0.22 [-0.37, 0.06]	-0.09 [-0.25, 0.07]	-0.05 [-0.21, 0.12]	-0.20 [-0.38, -0.02] ^a
Anxiety	-0.13 [-0.26, 0.00]	-0.08 [-0.22, 0.06]	-0.13 [-0.26, 0.00]	-0.03 [-0.16, 0.11]	-0.01 [-0.15, 0.13]	-0.20 [-0.34, -0.05] ^a
Delusions	-0.16 [-0.38, 0.06]	-0.15 [-0.39, 0.10]	-0.26 [-0.49, -0.04] ^a	-0.18 [-0.40, 0.05]	-0.13 [-0.37, 0.10]	-0.41 [-0.65, -0.16] ^a
Disinhibition	0.02 [-0.16, 0.19]	0.11 [-0.08, 0.30]	0.08 [-0.10, 0.27]	-0.03 [-0.21, 0.16]	0.03 [-0.17, 0.22]	-0.07 [-0.27, 0.13]

Eating behaviors	-0.07 [-0.20, 0.05]	0.01 [-0.13, 0.15]	-0.08 [-0.21, 0.05]	-0.08 [-0.22, 0.05]	-0.03 [-0.17, 0.11]	-0.06 [-0.20, 0.09]
Euphoria	-0.27 [-0.41, -0.12]^b	-0.17 [-0.33, 0.00]	-0.17 [-0.33, -0.02] ^a	-0.07 [-0.23, 0.08]	-0.06 [-0.22, 0.10]	-0.19 [-0.37, -0.02] ^a
Hallucinations	-0.41 [-0.70, -0.12] ^a	-0.02 [-0.36, 0.31]	-0.35 [-0.65, -0.05] ^a	-0.37 [-0.68, -0.06] ^a	-0.09 [-0.40, 0.22]	-0.29 [-0.66, 0.07]
Night-time behaviors	-0.13 [-0.27, 0.01]	0.07 [-0.09, 0.23]	-0.05 [-0.20, 0.10]	0.01 [-0.14, 0.16]	-0.29[-0.44, -0.13]^a	-0.07 [-0.07, 0.09]

Abbreviations: AD = Alzheimer's disease, MCI = mild cognitive impairment, MMSE = Mini-mental state examination, EF = executive functions, AMB = aberrant motor behaviors.

Data are standardized estimated annual β [95% confidence interval] corrected for sex, age, and education.

^a $p < 0.05$.

^b $p < 0.05$ after correcting for false discovery rate.

Table e-4. Associations between baseline NPS and cognitive decline over time according to clinical AD stage.

Preclinical AD (n=53)						
	MMSE	Memory	Attention	EF	Language	Visuospatial
Agitation*Time	0.11 [-0.24, 0.46]	-0.03 [-0.30, 0.24]	-0.03 [-0.33, 0.27]	-0.23 [-0.47, 0.01]	0.00 [-0.31, 0.31]	0.03 [-0.40, 0.46]
AMB*Time	0.28 [-0.12, 0.68]	0.24 [-0.09, 0.58]	-0.19 [-0.55, 0.17]	-0.17 [-0.47, 0.13]	0.30 [-0.07, 0.67]	0.05 [-0.45, 0.56]
Anxiety*Time	0.01 [-0.60, 0.63]	-0.18 [-0.66, 0.30]	0.74 [0.25, 1.24] ^a	0.29 [-0.15, 0.73]	0.05 [-0.49, 0.59]	-0.13 [-2.05, 1.79]
Delusions*Time	0.57 [-0.90, 2.03]	0.12 [-0.97, 1.22]	0.06 [-1.16, 1.28]	0.14 [-0.86, 1.15]	-0.62 [-1.87, 0.62]	1.96 [-1.67, 5.59]
Disinhibition* *Time	0.12 [-0.22, 0.47]	0.05 [-0.23, 0.32]	-0.12 [-0.43, 0.18]	-0.19 [-0.44, 0.06]	0.14 [-0.18, 0.45]	0.03 [-0.40, 0.46]
Eating behaviors*Time	0.25 [-0.15, 0.64]	0.17 [-0.15, 0.48]	0.04 [-0.30, 0.39]	0.19 [-0.10, 0.47]	0.00 [-0.36, 0.35]	0.02 [-0.47, 0.51]
Euphoria*Time	-0.08 [-0.61, 0.45]	-0.19 [-0.59, 0.21]	-0.03 [-0.47, 0.42]	-0.12 [-0.49, 0.25]	-0.24 [-0.70, 0.21]	-0.02 [-0.66, 0.62]
Hallucinations* Time	0.13 [-0.43, 0.68]	0.01 [-0.42, 0.44]	-0.03 [-0.50, 0.45]	0.50 [0.12, 0.88] ^a	0.15 [-0.33, 0.64]	-0.11 [-0.79, 0.58]
Night-time behaviors*Time	0.02 [-0.23, 0.28]	0.18 [-0.02, 0.38]	0.14 [-0.08, 0.36]	0.12 [-0.06, 0.30]	0.12 [-0.11, 0.35]	0.13 [-0.20, 0.47]
MCI due to AD (n=142)						
	MMSE	Memory	Attention	EF	Language	Visuospatial

Agitation*Time	0.03 [-0.13, 0.20]	0.05 [-0.10, 0.19]	0.00 [-0.16, 0.16]	0.03 [-0.09, 0.15]	0.07 [-0.09, 0.22]	-0.03 [-0.24, 0.18]
AMB*Time	0.08 [-0.17, 0.32]	0.15 [-0.05, 0.36]	-0.09 [-0.33, 0.15]	0.11 [-0.07, 0.29]	0.05 [-0.18, 0.28]	0.07 [-0.25, 0.38]
Anxiety*Time	0.14 [-0.02, 0.30]	-0.04 [-0.17, 0.10]	0.05 [-0.11, 0.20]	0.08 [-0.04, 0.20]	0.03 [-0.13, 0.18]	0.10 [-0.09, 0.29]
Delusions*Time	0.05 [-0.30, 0.40]	0.06 [-0.23, 0.35]	0.06 [-0.28, 0.39]	-0.04 [-0.30, 0.22]	-0.03 [-0.37, 0.30]	0.03 [-0.39, 0.45]
Disinhibition* Time	0.01 [-0.19, 0.22]	0.14 [-0.04, 0.33]	0.07 [-0.14, 0.27]	0.11 [-0.04, 0.27]	0.15 [-0.06, 0.35]	0.08 [-0.17, 0.33]
Eating behaviors*Time	0.01 [-0.16, 0.17]	-0.04 [-0.18, 0.10]	0.00 [-0.16, 0.16]	-0.07 [-0.19, 0.06]	-0.25 [-0.41, -0.08] ^a	-0.10 [-0.30, 0.11]
Euphoria*Time	-0.20 [-0.42, 0.02]	-0.14 [-0.32, 0.04]	-0.27 [-0.48, -0.06] ^a	-0.26 [-0.42, -0.10] ^a	-0.07 [-0.28, 0.14]	-0.23 [-0.49, 0.04]
Hallucinations* Time	N/A [#]	N/A [#]	N/A [#]	N/A [#]	N/A [#]	N/A [#]
Night-time behaviors*Time	0.14 [-0.01, 0.30]	0.06 [-0.07, 0.19]	0.02 [-0.13, 0.17]	-0.09 [-0.21, 0.02]	0.07 [-0.09, 0.22]	-0.09 [-0.28, 0.10]

AD dementia (n=325)

	MMSE	Memory	Attention	EF	Language	Visuospatial
Agitation*Time	-0.07 [-0.18, 0.04]	0.00 [-0.11, 0.12]	-0.04 [-0.15, 0.07]	0.00 [-0.10, 0.10]	0.07 [-0.04, 0.18]	-0.08 [-0.18, 0.02]
AMB*Time	-0.03 [-0.14, 0.07]	-0.02 [-0.13, 0.08]	-0.15 [-0.25, -0.05] ^a	-0.09 [-0.18, 0.00]	0.05 [-0.06, 0.15]	-0.02 [-0.12, 0.08]

Anxiety*Time	0.02 [-0.06, 0.10]	-0.02 [-0.10, 0.07]	-0.06 [-0.14, 0.02]	0.01 [-0.06, 0.08]	0.03 [-0.05, 0.11]	-0.01 [-0.08, 0.07]
Delusions*Time	0.12 [-0.07, 0.32]	-0.14 [-0.35, 0.06]	-0.14 [-0.33, 0.06]	-0.03 [-0.20, 0.14]	-0.01 [-0.21, 0.19]	-0.24 [-0.41, -0.07]
Disinhibition* Time	0.09 [-0.03, 0.21]	-0.09 [-0.22, 0.04]	-0.02 [-0.14, 0.10]	0.01 [-0.10, 0.12]	0.13 [0.01, 0.25] ^a	0.09 [-0.02, 0.20]
Eating behaviors*Time	0.01 [-0.07, 0.08]	0.00 [-0.08, 0.08]	0.00 [-0.08, 0.07]	-0.01 [-0.08, 0.06]	0.05 [-0.03, 0.13]	0.04 [-0.03, 0.11]
Euphoria*Time	-0.06 [-0.22, 0.11]	-0.24 [-0.44, -0.05] ^a	-0.09 [-0.28, 0.09]	-0.19 [-0.35, -0.03] ^a	0.12 [-0.06, 0.31]	-0.02 [-0.17, 0.14]
Hallucinations* Time	-0.09 [-0.29, 0.10]	0.10 [-0.11, 0.32]	0.05 [-0.15, 0.26]	0.09 [-0.09, 0.28]	0.04 [-0.16, 0.24]	0.06 [-0.13, 0.25]
Night-time behaviors*Time	0.03 [-0.06, 0.12]	-0.02 [-0.11, 0.07]	0.12 [0.03, 0.20] ^a	0.06 [-0.02, 0.14]	0.04 [-0.05, 0.13]	0.01 [-0.07, 0.09]

Abbreviations: AD = Alzheimer's disease, MCI = mild cognitive impairment, MMSE = Mini-mental state examination, EF = executive functions, AMB = aberrant motor behaviors.

Data are standardized estimated annual β [95% confidence interval] corrected for sex, age, and education.

^a $p < 0.05$.

^b $p < 0.05$ after correcting for false discovery rate.

[#] No patients with MCI and hallucinations at baseline had follow-up assessments available.