

COVID-19 Risk Factors and Mortality among Native Americans

Katherine Leggat-Barr¹

Fumiya Uchikoshi²

Noreen Goldman³

¹ Princeton School for Public and International Affairs and Global Health and Health Policy Program, Princeton University, Princeton, New Jersey

² Office of Population Research and Department of Sociology, Princeton University, Princeton, New Jersey

³ Office of Population Research and Princeton School of Public and International Affairs, Princeton University, Princeton, New Jersey

Abstract

BACKGROUND: Academic research on the disproportionate impact of COVID-19 among Native Americans has largely been restricted to particular indigenous groups or reservations.

OBJECTIVE: We estimate COVID-19 mortality for Native Americans relative to other racial/ethnic groups and explore how state-level mortality is associated with known risk factors.

METHODS: We use the Standard Mortality Ratio (SMR), adjusted for age and county, to estimate COVID-19 mortality by racial/ethnic groups for the U.S. and 10 selected states. The prevalence of risk factors is derived from the American Community Survey and the Behavioral Risk Factor Surveillance System.

RESULTS: The SMR for Native Americans greatly exceeds those for Black and Latino populations and varies enormously across states. There is a strong correlation between the share of Native Americans living on a reservation and the SMR. The SMR for Native Americans is also highly correlated with the income-poverty ratio and the prevalence of multigenerational families, crowded housing, frontline worker status, and health insurance (excluding the Indian Health Service). Risk factors associated with socioeconomic status and co-morbidities are generally more prevalent for Native Americans living on homelands, a proxy for reservation status, than for those living elsewhere.

CONCLUSIONS: Most risk factors for COVID-19 are disproportionately high among Native Americans, particularly for those living on homelands. Reservation life appears to increase the risk of COVID-19 mortality.

CONTRIBUTION: We assemble and analyze a broader set of COVID-19-related risk factors for Native Americans than previous studies, a critical step toward understanding the exceptionally high COVID-19 death rates in this population.

Introduction

One of the many egregious consequences of the COVID-19 pandemic has been the disproportionate impact of morbidity and mortality on already disadvantaged populations (Thakur et al. 2020; Andrasfay and Goldman 2021). Although the full effect of the coronavirus for all racial and ethnic groups has yet to be determined, high quality estimates for Native Americans remain particularly scarce. Despite extensive media attention to Native Americans, academic research on COVID-19 in this population has been much less common than for Black and Latino Americans, and along with media reports, has been largely restricted to particular indigenous groups or reservations. This limitation is due in large part to the deficiencies and paucity of data for Native Americans (Hatcher et al. 2020).

Native Americans, defined as those who self-identify in the U.S. Census as American Indian or Alaska Native (AI/AN) alone or in conjunction with other racial or ethnic groups, comprise about 6 million individuals or two percent of the U.S. population. About half of this group identifies as single-race Native American (Norris, Vines, and Hoeffel 2012). The vast majority of these individuals self-identify as American Indian rather than Alaska Native.¹ We use the term Native American to refer to estimates for the combined group AI/AN.

As with other marginalized groups, high rates of vulnerability to COVID-19 among Native Americans have been attributed to long-standing structural inequalities and racism that include poverty, inadequate health care and health insurance, a high prevalence of co-morbidities that increase susceptibility to COVID-19, and poor living conditions including inadequate plumbing and crowded living quarters (Kakol, Upson, and Sood 2020). Some or all of these factors may be especially prevalent on Native American reservations (Deweese and Marks, 2017; Yellow Horse, Deschine, Parkhurst, and Huyser 2020; Yellow Horse, Yang, and Huyser 2021). Nevertheless, the importance of these risk factors for COVID-19 mortality among Native Americans remains largely speculative.

Our paper has several objectives. The first is to provide improved estimates of COVID-19 death rates for Native Americans in comparison with the Black, Latino and White populations. The most frequently reported measures, such as the proportion of the group dying from COVID-19 in relation to the group's representation in the population, fail to adjust for the large differences in underlying age distributions across racial and ethnic groups. Although several recent estimates account for these age differences (Bassett et al. 2020; Xu et al. 2021), our estimates control for variation in geographic distribution in addition to age.

A second goal is to determine the prevalence and potential importance of risk factors that may be related to COVID-19 mortality. We measure a broader set of risk factors compared with most earlier studies on Native American vulnerability: poverty, frequency of multigenerational households and residential crowding, health insurance coverage, frontline worker status, smoking status, the prevalence of seven health conditions that may increase the severity of COVID-19 infection and risk of death, and state policies to mitigate viral transmission. We estimate these COVID-19 death rates and prevalence of risk factors at both a national level and for the 10 states with the largest number of COVID-19 deaths among Native Americans:

¹ About 95% of those reporting a single-race AI/AN for the U.S. and over 99% for most states in our analysis report American Indian race rather than Alaskan Native.

Arizona, California, Minnesota, Mississippi, New Mexico, New York, Oklahoma, South Dakota, Utah, and Washington.²

The final objective of this paper is to assess whether vulnerability to COVID-19 varies between Native Americans who live on vs. off reservations and whether reservation status is associated with COVID-19 death rates across states.

In order to address these questions, we analyze several data sources: provisional counts of COVID-19 deaths during 2020 from the Centers for Disease Control, recent population counts for the U.S., and potential risk factors from both the American Community Survey (ACS) and the Behavioral Risk Factor Surveillance Survey (BRFSS).

Background

SARS-CoV-2, the virus responsible for COVID-19, is transmitted primarily through respiratory droplets and aerosols when an infected person sneezes, coughs, talks or breathes in close proximity to others (Wilson, Corbett, and Tovey 2020). Research has identified a diverse set of socioeconomic and health-related risk factors that increase the likelihood of viral infection, illness severity, hospitalization and mortality. Many of these factors are likely to be especially prevalent among Native Americans, rendering them socially vulnerable to the coronavirus (Hathaway 2020; Yellow Horse, Yang, and Huyser 2021).

Both individual and community-level poverty are risk factors for COVID-19 transmission and severity (Patel et al. 2020; Williamson et al. 2020). Poverty status operates through multiple exposures including employment status and job characteristics, housing and living conditions, health-related behaviors and health status. Studies have pointed to likely higher risks among frontline and essential workers, who are often drawn disproportionately from disadvantaged racial and ethnic groups, and have jobs involving close indoor contact with others, often without adequate protective measures and without the possibility of being performed remotely (Goldman et al. 2020; Selden and Berdahl 2020). Frequent reliance on public transport exacerbates the infection risks faced by these workers (McLaren 2020). Because household transmission has been a significant source of COVID-19 spread (Haroon et al. 2020; Shen et al. 2020), multigenerational households, particularly those in crowded living spaces, enhance COVID-19 transmission and are particularly dangerous for the elderly (Jing et al. 2020; Chen and Kreiger 2021).

One reason for the much higher death rates among older individuals is the increasing prevalence with age of co-morbidities that are believed to raise the risk of dying from the virus. According to the Centers for Disease Control, the health-related conditions in the highest risk category include obesity, cardiovascular disease, cancer, chronic obstructive pulmonary disease, (type 2) diabetes, and chronic kidney disease (Centers for Disease Control and Prevention 2021b). Although there is some evidence that asthma is also a risk factor for COVID-19 severity and fatality, findings to date are mixed (Williamson et al. 2020). Consistent with earlier evidence for influenza and other respiratory diseases, a recent review of several studies suggests that smoking is associated with worse outcomes from COVID-19 infection (Vardabas and Nikitara 2020). Access to high-quality health care, which depends in part on health

² We chose the 10 states with the largest number of COVID-19 deaths among Native Americans as of July 2020, when we began the analysis. Nine of these states are among the 13 states with the largest number of COVID-19 deaths among Native Americans as of March 3, 2021 (New York is the exception).

insurance coverage as well as reliable and comprehensible health information, is an additional important determinant of mortality risk in general as well as from COVID-19 (Thakur et al. 2020).

An obvious risk factor for COVID-19 is the prevalence of infection in the local community, which affects the likelihood that an individual's day-to-day contacts with others will result in viral transmission. By modifying individual and group behaviors, local and state policies can have a substantial impact on these risks. Strong policy responses, such as lockdowns of businesses, restrictions on size of gatherings and travel, border closures, and quarantines have been associated with reduced COVID-19 transmission (Hsiang et al. 2020; Chaudhry et al. 2020).

Reservation Status

Tribal nations appear to be especially vulnerable to COVID-19. The high rates of poverty and generally poor living conditions characterizing reservation life have been implicated in excessive rates of infectious disease and mortality among American Indians for two hundred years or more before the arrival of COVID-19 (Jones 2006).

There are currently 326 Federal Indian reservations that are sovereign nations within the U.S. (Rodriguez-Lonebear et al. 2020). Until the 1950s, most Native Americans lived on reservations, but many relocated voluntarily or were coerced to move subsequently. A large number of reservations are located in rural areas. Almost four-fifths of Native Americans live off the reservation, largely in metropolitan areas (U.S. Department of Health and Human Services 2018). However, this proportion varies substantially across states with, for example, over 60% of Native Americans living on reservations in South Dakota in contrast to fewer than 10% in New York and California (see Table 1).³

Native American reservations are affected by many of the risk factors for infectious disease described earlier. The poverty rate far exceeds that of the general population, approaching 40% for some Native groups, with low levels of education and high rates of unemployment and food insecurity on reservations compared with the general population (Sarche and Spicer 2008; Jernigan et al. 2012). Many studies have pointed to lack of indoor running water or, more generally, inadequate household plumbing (Deitz and Meehan 2019; Rodriguez-Lonebear et al. 2020), a serious concern for quality of life in general but a particularly critical risk factor for the spread of infectious diseases like COVID-19. The Navajo Nation, the largest and most populous reservation in the U.S., and one that has garnered considerable media attention due to its high rates of COVID-19 infection and mortality, has some of the highest water and plumbing insecurity in the U.S. with about 40% of families lacking running water in their homes (Deitz and Meehan 2019). In a study of reported cases of COVID-19 on reservations during the early stages of the pandemic, the proportion of homes lacking running water was more strongly associated with COVID-19 case rates than other household variables (Rodriguez-Lonebear et al. 2020).

Inadequate access to and poor quality of health services for Native Americans has been a serious problem for many years. During 2020, the problems have become particularly acute because of obstacles that Native Americans have faced in obtaining COVID-19 tests, and the challenges that have affected treatment, hospitalization and recovery among those infected. Federally recognized tribes in the American Indian and Alaskan Native populations typically

³ This is based on Native Americans who classify themselves with a single race in the American Community Survey.

receive health care from the Indian Health Service (IHS), which has three branches: (1) provision of free health care primarily to those residing on or nearby reservations, (2) independent health services managed by the tribes but federally funded by the IHS, and (3) urban health services serving Native Americans living in urban areas (Sequist et al. 2011; Institute of Medicine 2003; Joe 2003).

The IHS and tribal health care systems are chronically underfunded. Many provider positions in the IHS remain unfilled, with vacancy rates of 25% in 2018 across the country and 30% for the Navajo Nation (U.S. Government Accountability Office 2018), and physicians report limited access to essential services as well as lack of high quality specialists, mental health services and diagnostic imaging (Sequist et al. 2011). Moreover, many Native Americans have no health insurance other than the IHS, Medicaid is underutilized in this population, and Native Americans frequently face discrimination and language barriers in non-IHS and non-tribal health facilities (Institute of Medicine 2003; Joe 2003; Cromer, Wofford, and Wyant 2019). Native Americans living on reservations (and in rural areas more generally) face especially severe challenges because of the extensive distance, travel time and cost involved in accessing health care and the lack of public health preparedness and infrastructure on reservations that pose a serious threat during a pandemic (Cromer, Wofford, and Wyant 2019; Groom et al. 2009; Rodriguez-Lonebear et al. 2020).

Although reservation life encompasses many challenges during this period of high viral transmission, there are some health-related advantages, including a strong sense of community and preservation of culture that may offer strong social support along with protection from psychological distress (Huyser et al. 2018). In addition, COVID-19 policies on some reservations have been much stricter than those in the states where these reservations are located. For example, the Navajo Nation, which is located in New Mexico, Arizona, and Utah, has had some of the most aggressive lockdown and curfew policies in the U.S. since the start of the pandemic along with high testing rates for the coronavirus (Navajo Nation Department of Health n.d.). Nevertheless, the analysis below suggests that these advantages may be outweighed by the many risk factors for COVID-19 infection and severity that are disproportionately present on reservations.

Data

We calculated death rates from COVID-19 from two sources. Provisional counts of deaths from COVID-19 by age group, race/ethnicity, and location (state and county) were taken from the National Center for Health Statistics (NCHS) for deaths through January 16, 2021 (we refer to these as 2020 deaths; Center for Disease Control and Prevention 2021a). Midyear population counts for 2019 for these same characteristics were obtained from the U.S. Census Bureau (2020). Numbers of deaths and population counts for the 10 states in the analysis are presented in Table 1.

We used two large, national surveys, the American Community Survey (ACS) and the Behavioral Risk Factor Surveillance System (BRFSS), to estimate potential risk factors for viral transmission and severity. The 2014-2018 ACS provided information on type of health insurance, the income-poverty ratio, and household information on living arrangements (whether the household is multigenerational and number of persons per room). Because the ACS asks about occupations in the past five years, we used the most recent ACS (2018) to determine frontline worker status. In an effort to obtain sufficiently large sample sizes of Native Americans, particularly at the state-level, while also ensuring consistency of question wording, we pooled data from eight years of the BRFSS (2011-2018) for estimates of smoking status and the

prevalence of seven health conditions: asthma, chronic obstructive pulmonary disease (COPD), kidney disease, cancer (excluding skin), heart disease, diabetes, and obesity. Sample sizes for each of the health conditions vary slightly due to different degrees of non-response.

Information on the extent to which each of the 10 states in our analysis remained open for business during the pandemic period was obtained from MultiState (Multistate 2020). In an effort to examine responses to the coronavirus, MultiState has measured the extent of lockdown of businesses and subsequent reopenings of the economy for state and local areas during much of the pandemic period.

Variables

Racial classification and reservation status

We consider four mutually exclusive racial/ethnic groups: non-Latino Whites, non-Latino Blacks, Latinos, and non-Latino Native Americans (American Indians and Alaskan Natives). For convenience, we drop the term “non-Latino” from subsequent references to these groups. Information on race and ethnicity for COVID-19 fatalities are drawn from the death certificates, whereas data in the surveys derive from self-reports. In the ACS, where finer racial distinctions are available, we consider both Native Americans with single-race identification and, separately, those with multiple-race identification (excluding those reporting Latino ethnicity). Estimates from the BRFSS are based only on self-reported single-race Native Americans.

We used My Tribal Area tool, which compiles 2014-2018 ACS estimates of reservation population by race, to calculate the number of single-race Native Americans residing on reservations or off-reservation trust land within the states in our analysis (U.S. Census Bureau n.d.); for simplicity, we refer to these estimates as living on a reservation. Because reservation status is not available on the microdata files from the 2014-2018 ACS obtained from IPUMS, we used homeland status as a proxy for reservation status in the state-specific estimates of risk factors. Homelands are similar to reservations,⁴ but include statistical lands in addition to legal ones; these statistical lands are intended to be “substantially meaningful” to local Native Americans (IPUMS USA n.d.). The variable in the ACS indicates whether the household is in a public use microdata area that includes any census block designated as a homeland area for American Indians, Alaska Natives, or Native Hawaiians. In July, 2020 the U.S. Supreme Court reclassified most tribal statistical land in Oklahoma as reservations. Because My Tribal Area tool has not been updated since that time and current status estimates would not in any case be an adequate measure of exposure in the recent past, Oklahoma is excluded from estimates that require reservation status but is included in estimates based on homeland status (McGirt vs. Oklahoma 2020).

Socioeconomic factors related to vulnerability

The measure of household crowding is determined from the number of persons and number of rooms in the household in the ACS. We use the proportion of households with more than one person per room, a standard assessment of overcrowding that has been applied in other studies (Blake, Kellerson and Simic 2007).

Estimates of the frequency of frontline workers, which are based on data in the 2018 ACS, include only those individuals who held any job in the five-year period prior to the survey, regardless of age, and who reported their occupation. About 88% of these individuals reported a

⁴ The median difference in the percent of single-race Native Americans living on reservations vs. living on homeland areas across the nine states (excluding Oklahoma) equals 1.3 percentage points.

job held during the single-year prior to the ACS. Our definition of frontline workers is based on the definition offered by Dingel and Neiman (2020) and used by Blau, Koebe, and Meyerhofer (2020): occupations in which one-third or fewer workers can feasibly work from home, ascertained from responses to 15 questions in the Occupational Information Network (O*NET, n.d.).⁵

Health-related variables

Because the prevalence of risk factors considered in this analysis may vary substantially by age, particularly for the health-related variables, and because the racial/ethnic groups differ in their age distribution, we restricted the age range to 18-60 for the 2014-2018 ACS and 18-59 for the BRFSS. To further reduce potential bias, we age-standardized all health-related variables derived from the BRFSS.⁶

Each of the six illness variables is based on a question in the BRFSS asking respondents whether a medical professional ever told them that they had the particular condition. The measure of obesity is based on a calculation of BMI ($BMI \geq 30$), obtained from self-reported height and weight. A score was calculated as a count of the number of health conditions reported by the individual among these seven variables.

All estimates from the ACS and BRFSS are based on individual-level data, with the exception of the frequency of households comprising at least three generations and the measure of crowding, both of which are derived from household-level data.

State openness

To determine the degree of state openness during the pandemic, we used scores developed by MultiState for the period between May 4, 2020 and December 30, 2020. The scores were based on assessments five days per week, ranged between 0 (full lockdown) to 100 (fully open), and were derived from 11 variables pertaining to stay-at-home orders: the definition of “essential business;” openings or operation of non-essential offices, non-essential retail, construction sites, personal care services, physical fitness businesses, bars, and venues servicing large groups; and approaches to reopening within the state, region and county (Multistate 2020).

Analytic Strategy

The analysis is primarily descriptive. We begin by comparing age-place standardized 2020 COVID-19 mortality measures across the four ethnic/racial groups for the U.S. as a whole and examine variability in these rates across the 10 states. Next, to understand this variability, we compare potential risk factors across racial/ethnic groups, nationally and for the selected states. Because Native Americans who classify themselves with a single racial category are more likely to live on reservations than those who use multiple categories (Norris, Vines, and Hoeffel 2012), and because we anticipate that reservation status in itself may reflect higher levels of risk factors, we examine differences in potential vulnerability to the virus between single-race and multiracial Native American groups at the national level. At the state level, we directly compare risk factors between Native Americans residing on homelands and those living in other areas, as well as between Native Americans and other racial/ethnic groups. In the final stage of the

⁵ O*NET is a database of work characteristics by detailed occupation collected by the U.S. Department of Labor. See Goldman et al. (2020) for details of the construction of the frontline status variable.

⁶ The standardization is based on 2019 population estimates for the entire U.S. based on the following age groups: 18-29, 30-39, 40-49, and 50-59.

analysis, we explore the relationship between potential risk factors and COVID-19 mortality across states for Native Americans.

The racial and ethnic groups in this analysis differ in their underlying age distributions as well as their patterns of geographic residence across the U.S. (e.g., rural vs. urban and region). SARS-CoV-2 infection and death rates also vary by age and by location. Thus, in order to compare rates across groups, we standardize our estimates by age and county of residence. Because age- and race-specific death rates from COVID-19 are not currently available by county, we use indirect standardization techniques to calculate the Standardized Mortality Ratio (SMR) for each racial/ethnic group. We follow the procedure used to examine racial/ethnic disparities in COVID-19 mortality early in the pandemic (Goldstein and Atherwood 2020). The SMR is a ratio of the observed number of COVID-19 deaths in a particular racial/ethnic group (during 2020) to the “expected” number. The expected number for each group reflects the number of deaths that would have occurred during 2020 had the group experienced the COVID-19 death rates by age and county of the entire US standard population⁷ but retained the observed population distribution by age and county of the particular racial/ethnic group. In addition to calculating the SMRs for the U.S., we obtain the corresponding numbers for the 10 states in the analysis based on age and county information for the state, using the same standard schedule as for the nationwide calculation.

In the first part of the analysis, we examine the SMR for each of the three disadvantaged groups (Native American, Black and Latino populations) relative to (i.e., divided by) the SMR for Whites. To illustrate the importance of standardization, we also calculate the crude death rates from COVID-19 for each of these groups relative to the crude death rate for Whites. The crude death rate, so-called because it does not take age into account, is defined simply as the number of deaths from COVID-19 for a particular group relative to the total population for that group.

Analyses that examine potential risk factors for the virus across racial/ethnic groups are based on the ACS and BRFSS (the openness scores are an exception). The estimates of the prevalence of risk factors are weighted based on the sampling weights in the ACS and BRFSS, but numbers of survey participants and households shown in the tables are unweighted. In order to minimize our inclusion of unreliable estimates, state estimates from the ACS are not shown when the number of Native American households or individuals by homeland status falls below 50. In estimated associations that examine only Native Americans (i.e., correlations of risk factors with COVID-19 mortality across the states), we measure COVID-19 mortality with the SMR for Native Americans. Because our dataset includes only 10 states, we restrict statistical measures to simple correlations between the SMR and risk factors rather than undertake multiple regression analysis.

Results

Figure 1 shows the crude death rates and the SMRs for COVID-19 in the U.S. during 2020 for each of the Native American, Black and Latino populations relative to Whites. These estimates reveal that, when age and county of residence are taken into account, COVID-19 mortality among Native Americans relative to Whites is substantially higher than suggested by crude

⁷ The standard mortality schedule is derived from COVID-19 deaths for 2020 for all races and ethnic groups combined and the entire U.S. population for 2019, with deaths and population based on 10-year age groups after the first age group of 0-4 (U.S. Census Bureau 2019).

death rates. The SMRs for Native Americans are 2.8 times as high as among Whites in comparison with the corresponding values of 1.6 and 1.8 for Blacks and Latinos respectively.

Figure 2 presents the SMRs for the Black, Latino and Native American populations relative to Whites in the 10 states. The estimates underscore the huge geographic variability in COVID-19 mortality. For the Native American population, these values are six or greater in half of the states, ranging from below one in New York to almost ten in South Dakota.

The scatterplots in Figure 3 show the SMRs for Native Americans in comparison with the percent of Native Americans residing on reservations (given in Table 1) across the states; the size of each circle is proportional to the population of Native Americans in the state.⁸ The pattern suggests a strong correlation ($r=0.80$) between living on a reservation and the SMR for COVID-19, with South Dakota having extremely large values on both of these measures in contrast to New York and California where few Native Americans reside on reservations and the SMR is relatively low. We examined the sensitivity of this correlation to two alternative measures of COVID-19 mortality among Native Americans: the SMR controlling only for age ($r=0.87$) and the SMR for Native Americans controlling for age and county relative to the corresponding SMR for whites ($r=0.88$). Nevertheless, we recognize that estimates based on such a small sample size may not be robust and statistical testing could be misleading.

Table 2 presents the risk factors derived from the ACS and the BRFSS for the four racial/ethnic groups for the U.S. Corresponding values for the 10 states are given in the appendix. The estimates in Table 2 indicate that potential exposure to COVID-19 among Native Americans is higher than that among the other racial/ethnic groups and that single-race Native Americans appear to be at higher risk for COVID-19 exposure than their multiple-race counterparts. Single-race Native Americans have the lowest incomes, on average, and are most likely (along with Latinos) to be without health insurance (excluding the IHS). Second only to Latinos, single-race Native Americans are the most likely group to hold frontline occupations and to live in multigenerational and crowded households. The most common frontline occupations held by Native Americans for the period prior to the pandemic included cashiers, janitors and building cleaners, maids, construction and other laborers, and drivers, all relatively low paying positions that require face-to-face exposure with others (authors' calculations). Workers in these low-income jobs are not typically provided with personal protective equipment and are likely to encounter weak safety standards in their places of employment (Carlsten et al. 2021; Gaitens et al. 2021).

Native Americans also have the poorest health profile in terms of COVID-19 susceptibility as demonstrated by the age-standardized proportions in the lower panel of Table 2. They have the highest prevalence of smoking, COPD, kidney disease, cancer, heart disease, and diabetes among the four groups and the level of obesity among Native Americans is exceeded only by the Black population. For the most part, the corresponding state-specific estimates, presented in the appendix, also show higher levels of co-morbidities among Native Americans relative to the other groups. It is quite likely that some of these estimates are too low, especially for the Native American and Latino populations, because they depend on whether the a health professional told the respondent that he/she had the condition and, thus, on access to medical care.

Figure 4 presents scatterplots between the SMRs and state-specific risk factors for Native Americans, including the average openness scores in Table 3 and eight of the risk factors in the appendix. The graphs are not presented for several health conditions with low prevalence

⁸ For reasons discussed earlier, Oklahoma is not included in this graph.

(cancer, heart disease, COPD and kidney disease) nor for asthma, which does not have a well-established link with COVID-19 at the present time. The scatterplots reveal particularly strong correlations (>0.8 in absolute value) with the SMR across states for the income-poverty ratio and the prevalence of multigenerational families, crowded housing, frontline worker status, and health insurance (excluding the IHS). South Dakota, the state with the highest SMR, has one of the most extreme values on all of these risk factors as well as the highest proportion of Native Americans residing on reservations.

As noted earlier, we do not fit multiple regression models, both because we have only 10 observations and because many of the risk factors are highly correlated with one another, suggesting the co-occurrence of risk factors for COVID-19 and rendering it impossible to determine which are most salient. The correlations among the risk factors, as well as between the risk factors and the SMR, are shown in a heatmap in Figure 5. Not surprisingly, the income-poverty ratio is highly correlated with the presence of multigenerational households and household crowding as well as with the proportion of Native Americans with health insurance (excluding the IHS) and the proportion with frontline occupations. These findings suggest that Native Americans are at risk of dying from COVID-19 through multiple exposures.

To establish whether Native Americans living on homelands face higher risks than others, the state tables in the appendix stratify the ACS estimates for single-race Native Americans by homeland status. Although we cannot consider the association between risk factors and COVID-19 mortality separately by homeland status (because the COVID-19 death information does not include reservation or homeland status), these data permit us to assess the extent to which risk factors vary between those residing on vs. off homelands. For the vast majority of estimates across the states, the prevalence of COVID-19 risk factors is higher for Native Americans living on homelands than for those living off homelands. The differentials are particularly large for living conditions, i.e., multigenerational households and crowding. For example, households comprising at least three generations are about 50% or more frequent for those residing on homelands in several of the states. Not surprisingly given that IHS facilities are typically situated on or near Native lands, Native Americans on homeland areas have higher rates of health insurance when the IHS is included in the measure. However, in all states, possession of health insurance other than the IHS is more common among Native Americans residing off homelands.

Discussion

From the early days of the pandemic, the media has drawn attention to high mortality from the coronavirus among Native Americans, particularly those on select reservations. The analysis presented here measures this high fatality in a way that enhances comparability across racial and ethnic groups. Once age and geographic distribution are taken into account, the national 2020 COVID-19 mortality level for Native Americans is almost three times that for Whites, an estimate considerably higher than the corresponding values for Blacks and Latinos. In half of the states in our analysis, Native Americans experience a mortality level from COVID-19 at least six times that of the White population.

An examination of risk factors for COVID-19 points to several related variables that are particularly prevalent among Native Americans and that have been previously identified as risk factors for exposure to the virus or for dying among those infected. Most notable among these are high rates of poverty, which, in turn, are correlated with high density housing (high levels of crowding and multigenerational families residing in a single household), employment as frontline workers, prevalence of co-morbidities (especially obesity and diabetes) that are associated with

hospitalization and mortality from the virus, and low rates of health insurance other than the IHS. These risk factors are more prevalent among Native Americans who self-identify as a single-race as well as those who reside on homelands compared with their respective counterparts, underscoring potential associations between reservation life and both viral exposure and disease severity. At the same time, it is important to recognize that, although they act as risk factors for viral transmission, multigenerational households and large family compounds may also serve important social and cultural functions that could be protective of health.

Further evidence that reservation life may increase risk for COVID-19 mortality comes from the extremely high correlation between the percent of Native Americans in a given state residing on reservations and the SMR for Native Americans in that state. This association suggests that large differences in the pervasiveness of reservation life may drive much of the statewide variation in COVID-19 mortality. However, we recognize that our ecological analysis cannot provide sufficient evidence of a causal connection between reservation life and the risk of COVID-19 mortality. Moreover, given that the variables in our analysis capture only some of the important components of risk, our efforts to identify the most important facets of reservation life underlying the high mortality of Native Americans remain uncertain.

A particular problem in our analysis is that possession of health insurance is a poor proxy for access to quality healthcare, a critical variable during this pandemic. Despite the establishment of treaties several generations ago, the federal government has largely neglected its obligation to provide suitable healthcare to Native Americans (Warne and Frizzel 2014). The Indian Health Service has been and continues to be woefully underfunded, a problem that is likely to be more severe among those residing on reservations who depend on the IHS for healthcare (Congressional Research Service 2016). In 2019, per capita spending for the Indian Health Service was less than half of the per capita spending on healthcare for the general U.S. population (Indian Health Service n.d.), making it extremely difficult for health facilities to acquire the resources needed to mount an effective pandemic response. For example, the IHS has few ICUs and ventilators or appropriate medical providers to treat critically ill COVID-19 patients (Schultz 2020). The historical legacies of injustice, many of which resulted from federal government actions, have created mistrust among Native Americans particularly toward the Indian Health Service and other medical institutions (Guadagnolo et al. 2009; Pacheco et al. 2013). Past experiences of discrimination, culturally and linguistically insensitive care, lack of reliable internet and phone service, and language and transportation barriers may have further reduced the likelihood that Native Americans received timely and comprehensible information about the pandemic and effective healthcare when needed (Guadagnolo et al. 2009; Joe 2003).

Another risk factor that is only partly taken into account in our analysis is inadequate housing infrastructure. Beyond the issue of crowding, Native Americans are much more likely than other groups to lack adequate plumbing in their households. About 2.5% of single-race Native American households lack hot and cold running water (in contrast to 0.2-0.4% among the other groups; authors' calculations from the ACS). An analysis of "plumbing poverty" demonstrates that lack of complete plumbing is both spatially and racially clustered, with an especially high prevalence among Native Americans and a concentration in areas containing reservations (as high as 73 percent in one neighborhood in northern Arizona that includes five reservations; Deitz and Meehan 2019). Such water insecurity presents an obvious risk to the spread of communicable diseases and a barrier for adherence to government guidelines regarding handwashing and sanitation during the COVID-19 pandemic.

Health disparities between Native Americans and other groups have persisted for centuries and are reflected today in higher death rates among Native Americans for almost all major causes of death, infectious and chronic (Jones 2006). The extraordinary COVID-19 fatality rates among Native Americans are due, at least in part, to a high prevalence of illnesses and health conditions, many of which are known risk factors for COVID-19 infection and severity. In particular, extremely high rates of obesity and diabetes, which result partly from high levels of food insecurity in this population (Jernigan et al. 2013), appear to increase the risk that Native Americans acquire SARS-CoV-2 infections, develop complications and die (Zhou et al. 2020). In the current pandemic, these co-morbidities portend a poor prognosis from COVID-19.

There are several additional limitations to this analysis. As mentioned previously, the lack of individual data does not permit us to assess the relative importance of the risk factors investigated here or to estimate the magnitude of risk for those residing on vs. off reservations. Furthermore, the absence of reliable data on COVID-19 cases prevents us from distinguishing the extent to which Native Americans are more susceptible than other groups to infection versus the degree to which infections in this group are more likely to result in fatalities. An additional concern relates to inaccuracies in the data on deaths from COVID-19 due to misattribution of cause of death on death certificates, exclusion of deaths indirectly attributable to COVID-19 (e.g., from delayed or foregone care from other health conditions), and errors in recording of age, race and ethnicity. Misclassification of race and ethnicity is a particularly severe problem among Native Americans, much more so than among other groups, with previous studies identifying substantial misclassification (underreporting in particular) of Native American race on death certificates as well as in survey data (Espey et al. 2014; Haozous et al. 2014). Moreover, in a recent study, Becker and colleagues underscore the huge variation in definitions of the Native American population across health surveys in the U.S. and the implications of such inconsistency for measurement of health disparities in the Native American population (Becker et al. 2021).

Elevated COVID-19 mortality serves as a stark reminder of the legacies of historical mistreatment of Native Americans and the continued failure of the federal, state and local governments to meet the most basic needs of this population. Our analysis shows that the death rates from COVID-19 vary substantially across states and suggests differences between those living on vs. off reservations, but there is undoubtedly further variability that we cannot assess. Previous studies have underscored heterogeneity among reservations, with substantial variability in size and, as a consequence, in the ability of tribal governments to govern effectively. There are also large differences in the prevalence of health conditions and access to healthcare (Department of the Interior n.d.; U.S. Department of Health and Human Services; Indian Health Service 2014). These disparities almost certainly have led to differential infection and fatality rates across reservations during the pandemic, above and beyond the infection rates in the broader geographic areas. Consistent with this variability, research has underscored the huge diversity among Native Americans and the unique vulnerabilities of different Native American groups (Cheek et al. 2014; Groom et al. 2009; Yellow Horse et al. 2021).

Despite these critical differences in susceptibility within the Native American population, the overall set of factors associated with high risk for COVID-19 in this population are largely the same as those that have been identified as increasing the vulnerability of Native Americans to a host of diseases throughout generations, factors that reflect long-standing structural inequalities and racism. As many researchers and policy experts have noted, increased support for Native American communities is required at multiple levels, from distal to proximate variables that affect both exposure to and treatment of disease. A desperately needed first step to reduce COVID-19 fatalities among Native Americans during the ongoing pandemic, and to avert a

disproportionate impact in the next one, is increased funding and coordinated policies for improvements in public health, basic infrastructure, and access to quality healthcare.

Acknowledgments

We would like to acknowledge financial support from the Center for Health and Wellbeing at Princeton University, assistance with the frontline calculations from Keunbok Lee, helpful comments on an earlier draft from Theresa Andrasfay, Dana Glej, Gracie Himmelstein and Anne Pebley, and assistance with the tables from Kristen Cuzzo.

References

- Andrasfay, T. and Goldman, N. (2021). Reductions in 2020 US life expectancy due to COVID-19 and the disproportionate impact on the Black and Latino populations. *Proceedings of the National Academy of Sciences* 118(5).
- Becker, T., Babey, S.H., Dorsey, R., and Ponce, N.A. (2021). Data Disaggregation with American Indian/Alaska Native Population Data. *Population Research and Policy Review* 40(1):103–125. doi:10.1007/s11113-020-09635-2.
- Blake, K., Kellerson, R., and Simic, A. (2007). *Measuring overcrowding in housing*. Bethesda, Maryland: US Department of Housing and Urban Development, Office of Policy Development and Research.
- Blau, F.D., Koebe, J., and Meyerhofer, P.A. (2020). Who are the essential and frontline workers? National Bureau of Economic Research (Working Paper 27791).
- Carlsten, C., Gulati, M., Hines, S., Rose, C., Scott, K., Tarlo, S.M., Torén, K., Sood, A., and Hoz, R.E. (2021). COVID-19 as an occupational disease. *American Journal of Industrial Medicine*: ajim.23222.
- Center for Disease Control and Prevention (2021a). COVID-19 provisional counts - weekly updates by select demographic and geographic characteristics. [electronic resource] Atlanta: National Center for Chronic Diseases Prevention and Health Promotion https://www.cdc.gov/nchs/nvss/vsrr/covid_weekly/index.htm.
- Centers for Disease Control and Prevention (2021b). COVID-19 and your health. [electronic resource]. Atlanta: National Center for Chronic Diseases Prevention and Health Promotion <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html>.
- Chaudhry, R., Dranitsaris, G., Mubashir, T., Bartoszko, J., and Riazi, S. (2020). A country level analysis measuring the impact of government actions, country preparedness and socioeconomic factors on COVID-19 mortality and related health outcomes. *EClinicalMedicine* 25.
- Cheek, J.E., Holman, R.C., Redd, J.T., Haberling, D., and Hennessy, T.W. (2014). Infectious disease mortality among American Indians and Alaska Natives, 1999–2009. *American Journal of Public Health* 104: S446–S452.
- Chen, J.T., and Krieger, N., (2021). Revealing the unequal burden of COVID-19 by income, race/ethnicity, and household crowding: US county versus zip code analyses. *Journal of Public Health Management Practices* 27(1): S43-S56.
- Congressional Research Service. (2016). The Indian Health Service (IHS): an overview [electronic resource] <https://crsreports.congress.gov/product/pdf/R/R43330>.
- Cromer, K. J., Wofford, L., and Wyant, D. K. (2019). Barriers to healthcare access facing American Indian and Alaska Natives in rural America. *Journal of Community Health Nursing* 36(4): 165–187.
- Deitz, S., and Meehan, K. (2019). Plumbing poverty: mapping hot spots of racial and geographic inequality in U.S. household water insecurity. *Annals of the American Association of Geographers* 109(4): 1092–1109.
- Deweese, S. and Marks, B. (2017). *Twice Invisible: Understanding Rural Native America*. First Nations Development Institute.
- Dingel, J.I., and Neiman, B. (2020). How many jobs can be done at home? National Bureau of Economic Research (Working Paper 26948).
- Department of the Interior (n.d.). Fractionation [electronic resource]. <https://www.doi.gov/buybackprogram/fractionation>.
- Espey, D. K., Jim, M. A., Richards, T. B., Begay, C., Haverkamp, D., and Roberts, D. (2014).

- Methods for improving the quality and completeness of mortality data for American Indians and Alaska Natives. *American Journal of Public Health* 104(S3): S286–S294.
- Gaitens, J., Condon, M., Fernandes, E., and McDiarmid, M. (2021). COVID-19 and essential workers: A narrative review of health outcomes and moral injury. *International Journal of Environmental Research and Public Health* 18(4):1446.
- Goldman, N., Pebley, A., Lee, K., Andrasfay, T., Pratt, B. (2020). Racial and ethnic differentials in COVID-19-related job exposures by occupational status in the US. *medRxiv*. 1-20. <https://doi.org/10.1101/2020.11.13.20231431>
- Goldstein, J.R., and Atherwood, S. (2020). Improved measurement of racial/ethnic disparities in COVID-19 mortality in the United States. *medRxiv*. <https://doi.org/10.1101/2020.05.21.20109116>
- Groom, A.V., Jim, C., LaRoque, M., Mason, C., McLaughlin, J., Neel, L., Powell, T., Weiser, T., and Bryan, R.T. (2009). Pandemic influenza preparedness and vulnerable populations in tribal communities. *American Journal of Public Health* 99: S271–S278.
- Guadagnolo, B.A., Cina, K., Helbig, P., Molloy, K., Reiner, M., Cook, E.F., and Petereit, D.G., (2009). Medical mistrust and less satisfaction with health care among Native Americans presenting for cancer treatment. *Journal of Health Care for the Poor and Underserved* 20: 210–226.
- Haozous, E.A., Strickland, C.J., Palacios, J.F., and Solomon, T.G.A. (2014). Blood politics, ethnic identity, and racial misclassification among American Indians and Alaska Natives. *Journal of Environmental and Public Health* 2014: 1-9.
- Haroon, S., Chandan, J.S., Middleton, J., and Cheng, K.K. (2020). Covid-19: breaking the chain of household transmission. *British Medical Journal* 370 (m3181).
- Hathaway, E.D. (2020). American Indian and Alaska Native people: social vulnerability and COVID-19. *The Journal of Rural Health* 37(1): 256–259.
- Hatcher, S.M., Agnew-Brune, C., Anderson, M., Zambrano, L., Rose, C., Jim, M., Baugher, A., Liu, G., Patel, S., Evans, M., Pindyck, T., Dubray, C., Rainey, J., Chen, J. Sadowski, C., Winglee, K., Penman-Aguilar, A., Dixit, A., Claw, E., Pashall, C., Provost, E., Ayala, A., Gonzalez, G., Ritchey, J., Davis, J., Warren-Mears, V., Joshi, S., Weiser, T., Echo-Hawk, A., Dominguez, A., Poel, A., Duke, C., Ransby, I., Apostolou, A., and McCollum, J. (2020). COVID-19 among American Indian and Alaska Native persons — 23 states, January 31–July 3, 2020. *MMWR Morbidity and Mortality Weekly Report* 69: 1166–1169
- Hsiang, S., Allen, D., Annan-Phan, S., Bell, K., Bolliger, I., Chong, T., Druckenmiller, H., Huang, L.Y., Hultgren, A., Krasovich, E., Lau, P., Lee, J., Rolf, E., Tseng, J., and Wu, T. (2020). The effect of large-scale anti-contagion policies on the COVID-19 pandemic. *Nature* 584: 262–267.
- Huyser, K.R., Angel, R.J., Beals, J., Cox, J.H., Hummer, R.A., Sakamoto, A., Manson, S.M., and the AI-SUPERPFP Team (2018). Reservation lands as a protective social factor: An analysis of psychological distress among two American Indian tribes. *Socius: Sociological Research for a Dynamic World* 4:237802311880702.
- Indian Health Service. (n.d.). IHS Profile | fact sheets newsroom. [electronic resource] <https://www.ihs.gov/newsroom/factsheets/ihsprofile/>.
- Institute of Medicine. (2003). *Unequal treatment: confronting racial and ethnic disparities in health care*. The National Academies Press.
- IPUMS USA. (n.d.) HOMELAND [electronic resource] https://usa.ipums.org/usa/action/variables/HOMELAND#description_section.
- Jernigan, V.B.B., Garrouette, E., Krantz, E.M., and Buchwald, D. (2013). Food insecurity and obesity among American Indians and Alaska Natives and Whites in California. *Journal of Hunger and Environmental Nutrition* 8: 458–471.
- Jernigan, B. B. V., Salvatore, A. L., Styne, D. M., and Winkleby, M. (2012). Addressing food

- insecurity in a Native American reservation using community-based participatory research. *Health Education Research* 27(4): 645–655.
- Jing, Q.-L., Liu, M.-J., Zhang, Z.-B., Fang, L.-Q., Yuan, J., Zhang, A.-R., Dean, N.E., Luo, L., Ma, M.-M., Longini, I., Kenah, E., Lu, Y., Ma, Y., Jalali, N., Yang, Z.-C., and Yang, Y. (2020). Household secondary attack rate of COVID-19 and associated determinants in Guangzhou, China: a retrospective cohort study. *The Lancet Infectious Diseases* 20: 1141–1150.
- Joe, J. R. (2003). The rationing of healthcare and health disparity for the American Indians/Alaska Natives. In: Institute of Medicine (ed.), *Unequal Treatment: Confronting Racial and Ethnic Disparities in Health Care* National Academies Press: 528–551.
- Jones, D.S., (2006). The persistence of American Indian health disparities. *American Journal of Public Health* 96: 2122–2134.
- Kakol, M., Upson, D., and Sood, A. (2020). Susceptibility of southwestern American Indian tribes to coronavirus disease 2019 (COVID-19). *Journal of Rural Health* 37(1): 197-199.
- McGirt v. Oklahoma, (2020). 18 U.S. 9526 [electronic resource] https://www.supremecourt.gov/opinions/19pdf/18-9526_9okb.pdf
- McLaren, J. (2020). Racial disparity in COVID-19 deaths: seeking economic roots with census data. National Bureau of Economic Research (Working Paper 27407).
- Multistate (2020). COVID-19 state reopening guide. [electronic resource] <https://www.multistate.us/issues/covid-19-state-reopening-guide>.
- Navajo Nation Department of Health, (n.d.) COVID-19 [electronic resource] <https://www.ndoh.navajo-nsn.gov/covid-19>.
- Norris, T., Vines, P. L., and Hoeffel, E.M. (2012). The American Indian and Alaska Native population: 2010. U.S. Census Bureau. [electronic resource] <https://www.census.gov/history/pdf/c2010br-10.pdf>
- O*NET, n.d. O*NET OnLine [electronic resource] <https://www.onetonline.org/>.
- Pacheco, C.M., Daley, S.M., Brown, T., Filippi, M., Greiner, K.A., and Daley, C.M. (2013). Moving forward: breaking the cycle of mistrust between American Indians and researchers. *American Journal of Public Health* 103(12): 2152–2159.
- Patel, J.A., Nielsen, F.B.H., Badiani, A.A., Assi, S., Unadkat, V.A., Patel, B., Ravindrane, R., and Wardle, H. (2020). Poverty, inequality and COVID-19: the forgotten vulnerable. *Public Health* 183: 110–111.
- Rodriguez-Lonebear, D., Barceló, N.E., Akee, R., and Carroll, S.R. (2020). American Indian reservations and COVID-19: correlates of early infection rates in the pandemic. *Journal of Public Health Management and Practice* 26: 371–377.
- Sarche, M., and Spicer, P. (2008). Poverty and health disparities for American Indian and Alaska Native children: current knowledge and future prospects. *Annals of New York Academy of Sciences* 1136: 126–136.
- Schultz, H. (2020). Native American communities and COVID-19: how foundations can help. *Health Affairs* [electronic resource] doi: 10.1377/hblog20200331.659944.
- Selden TM, and Berdahl TA. (2020). COVID-19 and racial/ethnic disparities in health risk, employment, and household composition. *Health Affairs* 39: 1624–1632.
- Sequist, T. D., Cullen, T., Bernard, K., Shaykevich, S., Orav, E. J., and Ayanian, J. Z. (2011). Trends in quality of care and barriers to improvement in the Indian Health Service. *Journal of General Internal Medicine* 26(5): 480–486.
- Shen, M., Peng, Z., Guo, Y., Rong, L., Li, Y., Xiao, Y., Zhuang, G., and Zhang, L. (2020). Assessing the effects of metropolitan-wide quarantine on the spread of COVID-19 in public space and households. *International Journal of Infectious Diseases* 96: 503–505.
- Thakur, N., Lovinsky-Desir, S., Bime, C., Wisnivesky, J.P., and Celedón, J.C. (2020). The structural and social determinants of the racial/ethnic disparities in the U.S. COVID-19

- pandemic. What's our role? *American Journal of Respiratory and Critical Care Medicine* 202: 943–949.
- U.S. Census Bureau, (n.d.). My tribal area [electronic resource]<https://www.census.gov/tribal/?st=04&aianihh=2430>.
- U.S. Census Bureau, (2019). 2019 population estimates by age, sex, race and hispanic origin [electronic resource] <https://www.census.gov/newsroom/press-kits/2020/population-estimates-detailed.html>.
- U.S. Census Bureau (2020). Annual county resident population estimates by age, sex, race, and Hispanic origin: April 1, 2010 to July 1, 2019 [electronic resource] <https://www.census.gov/newsroom/press-kits/2020/population-estimates-detailed.html>.
- U.S. Department of Health and Human Services; Indian Health Service (2014). *Trends in Indian Health 2014 Edition*. [electronic resource]. https://www.ihs.gov/sites/dps/themes/responsive2017/display_objects/documents/Trends2014Book508.pdf
- U.S. Department of Health and Human Services. (2018). *Profile: American Indian/Alaska Native*. [electronic resource] <https://minorityhealth.hhs.gov/omh/browse.aspx?lvl=3&lvlid=62>.
- U.S. Government Accountability Office. (2018). *Indian Health Service agency faces ongoing challenges filling provider vacancies*. [electronic resource]. Washington, D.C. <https://www.gao.gov/assets/700/693940.pdf>.
- Vardavas, C.I., and Nikitara, K. (2020). COVID-19 and smoking: a systematic review of the evidence. *Tobacco Induced Diseases* 18: 2-4.
- Warne, D., and Frizzell, L.B. (2014). American Indian health Policy: historical trends and contemporary issues. *American Journal of Public Health* 103 Supplement 3: S263–S267.
- Williamson, E.J., Walker, A.J., Bhaskaran, K., Bacon, S., Bates, C., Morton, C.E., Curtis, H.J., Mehrkar, A., Evans, D., Inglesby, P., Cockburn, J., McDonald, H.I., MacKenna, B., Tomlinson, L., Douglas, I.J., Rentsch, C.T., Mathur, R., Wong, A.Y.S., Grieve, R., Harrison, D., Forbes, H., Schultze, A., Croker, R., Parry, J., Hester, F., Harper, S., Perera, R., Evans, S.J.W., Smeeth, L., and Goldacre, B. (2020). Factors associated with COVID-19-related death. *Nature* 584: 430–436.
- Wilson, N., Corbett, S., and Tovey, E. (2020). Airborne transmission of COVID-19. *British Medical Journal* 370:m3206.
- Xu, J.J., Chen, J.T., Belin, T.R., Brookmeyer, R.S., Suchard, M.A., and Ramirez, C.M. (2021). Racial and ethnic disparities in years of potential life lost attributable to COVID-19 in the United States: An analysis of 45 states and the District of Columbia. *medRxiv*. doi:10.1101/2021.01.28.21249411.
- Yellow Horse, A.J., Deschine Parkhurst, N.A., and Huyser, K.R. (2020). COVID-19 in New Mexico tribal lands: Understanding the role of social vulnerabilities and historical racisms. *Frontiers in Sociology* 5:610355.
- Yellow Horse, A.J., Yang, T.-C., and Huyser, K.R. (2021). Structural inequalities established the architecture for COVID-19 pandemic among Native Americans in Arizona: a geographically weighted regression perspective. *Journal of Racial and Ethnic Health Disparities*.
- Zhou, Y., Chi, J., Lv, W., and Wang, Y. (2020). Obesity and diabetes as high-risk factors for severe coronavirus disease 2019 (Covid-19). *Diabetes/Metabolism Research and Reviews* 37(2): e3377.

Table 1. Number of Native Americans, number of COVID-19 deaths, and percent of Native Americans on reservations, 10 states^a

State	# of Native Americans	# of COVID-19 deaths	% Native Americans on reservations
Arizona	309,580	1,014	51.8
California	296,475	119	6.0
Oklahoma	294,676	397	NA ^b
New Mexico	199,896	739	50.2
Washington	95,048	90	28.3
New York	79,644	38	9.3
South Dakota	75,331	165	62.4
Minnesota	58,982	74	31.6
Utah	32,657	63	26.7
Mississippi	13,689	90	51.4

^a All numbers are based on Native Americans recorded as single-race.

^b Until the Supreme Court ruling on July 9th, 2020, there were legal uncertainties as to what land in Oklahoma should be classified as reservation land.

Source: ACS 5-year estimates and CDC Weekly Updates by Select Demographic and Geographic Characteristics (as of Nov 4, 2020). Number of deaths are updated as of January 21, 2021.

Table 2. Risk factors for COVID-19 infection and severity by race and ethnicity in the U.S.: American Community Survey (2014-2018) and the Behavioral Risk Factor Surveillance System (2011-2018)

<u>American Community Survey^a</u>					
	Native Americans (single race)	Native Americans (multiple race)	Whites	Latinos	Blacks
Income-poverty ratio	2.4	2.8	3.5	2.6	2.6
% 3+ generation HH ^b	6.8	4.3	2.7	7.6	5.3
% crowded HH ^b	9.0	4.3	2.0	13.4	4.4
% health insurance (exc. IHS)	72.7	83.1	90.5	72.8	83.2
% health insurance (inc. IHS)	91.2	86.1	90.5	72.8	83.2
% frontline workers ^c	66.1	63.6	53.1	70.6	64.7
Number of households	35,497	34,284	2,724,237	517,360	396,491
Number of persons	86,691	76,066	5,932,772	1,377,777	939,811
<u>Behavioral Risk Factor Surveillance System^d</u>					
	Native Americans (single race)	Native Americans (multiple race)	Whites	Latinos	Blacks
% smokers	33.6	-	22.2	14.4	21.2
% asthma	21.5	-	15.0	11.9	17.0
% COPD	8.4	-	4.7	2.9	5.2
% kidney disease	2.9	-	1.6	2.2	2.1
% cancer (excl. skin)	5.0	-	3.6	2.4	2.5
% heart disease	3.0	-	1.7	1.9	2.1
% diabetes	11.9	-	6.0	10.3	10.0
% obese (BMI≥30)	36.2	-	27.2	32.3	38.3
Mean # of health conditions ^e	0.87	-	0.59	0.64	0.77
Min number of persons	26,157	-	990,131	142,728	123,203
Max number of persons	38,339	-	1,430,443	221,843	182,662

Note: All BRFSS estimates are age standardized using 2019 population estimates for the US from the US Census Bureau (<https://www.census.gov/newsroom/press-kits/2020/population-estimates-detailed.html>). The BRFSS does not provide race-specific information on multiple races.

Estimated risk factors from the ACS and BRFSS are weighted; numbers of persons are unweighted.

^a Based on respondents ages 18-60 (except for estimates of frontline workers).

^b This estimate is based on information at the household-level; the remaining ACS variables are based on individual information.

^c Based on persons of all ages reporting jobs in the past five years from the single-year 2018 ACS.

^d Based on respondents ages 18-59. The number of persons providing responses in the BRFSS varies across health conditions. The minimum sample size pertains to persons with responses to all of the health condition variables.

^e The mean number of health conditions per person is the sum of affirmative responses to the following conditions: asthma, COPD, kidney disease, cancer, diabetes, heart disease, and obesity (BMI \geq 30).

Table 3. State openness scores, 10 states, 2020

	Average openness score	Minimum openness score	Maximum openness score
Arizona	78.9	29	91
California	25.6	10	33
Minnesota	55.9	25	67
Mississippi	65.2	41	72
New Mexico	27.8	13	36
New York	38.6	9	49
Oklahoma	89.9	80	93
South Dakota	93.8	92	96
Utah	83.9	62	86
Washington	36.5	13	46

Note: These scores were obtained from MultiState:

(<https://www.multistate.us/pages/covid-19-state-reopening-guide>)

These estimates cover the period beginning May 4th, 2020 until December 30th, 2020 and were assessed five days per week. The score evaluates the degree of state ‘openness,’ ranging from 0 (full lockdown) to 100 (fully open). Eleven factors are evaluated to create the score; see the text.




Fig 1. Crude Death Rates and Standardized Mortality Ratios (relative to Whites) for COVID-19 in the U.S.

Source: CDC Weekly Updates by Select Demographic and Geographic Characteristics (as of Jan 21, 2021)




Fig 2. Standardized Mortality Ratio (relative to Whites) for COVID-19, 10 states.

Source: CDC Weekly Updates by Select Demographic and Geographic Characteristics (as of Jan. 21, 2021)




Fig 3. Correlation (and regression line) between the percent of Native Americans on reservations and the Standardized Mortality Ratio among Native Americans across states.

Source: ACS 5-year estimates and CDC Weekly Updates by Select Demographic and Geographic Characteristics (as of Jan 21, 2021)




Fig 4. Correlations (and regression lines) between selected risk factors for COVID-19 infection and severity and the Standardized Mortality Ratio among Native Americans across states.

Source: ACS 5-year estimates, ACS 1-year estimates, BRFSS, and CDC Weekly Updates by Select Demographic and Geographic Characteristics (as of Jan. 21, 2021)




Fig 5. Heatmap of correlations between risk factors for COVID-19 infection and severity and the Standardized Mortality Ratio among Native Americans across states.