

Risk of reinfection after seroconversion to SARS-CoV-2: A population-based propensity-score matched cohort study

Online-only material

Antonio Leidi^{1*} MD, Flora Koegler^{1*} MD, Roxane Dumont², Richard Dubos², María-Eugenia Zaballa² PhD, Giovanni Piumatti^{2,6} PhD, Matteo Coen¹ PhD, Amandine Berner¹ MD, Pauline Darbellay Farhoumand¹ MD, Pauline Vetter MD⁴, Prof Nicolas Vuilleumier³ MD, Prof Laurent Kaiser⁴ MD, Prof Delphine Courvoisier⁵ PhD, Andrew S Azman² PhD, Prof Idris Guessous^{2*} MD, Prof Silvia Stringhini^{2*} PhD, SEROCO-V-POP study group


eFigure 1. Diagnostic plot for the propensity score matching

Description of reinfected individuals

eTable 1. Demographic, clinical and laboratory characteristics of individuals with suspected SARS-CoV-2 reinfection

eTable 2. Sensitivity analyses

eFigure 1. Diagnostic plot for the propensity score matching



eFigure 1. Standardized mean difference before (orange dots) and after (black dots) propensity score matching. A difference of less than 0.1 considered to be acceptable.

Description of reinfected individuals

Regarding the five *likely* cases in our main analysis, three of them did not report COVID-like symptoms before serologic testing, whereas reinfections were characterized by classical symptoms. One other seropositive individual was symptomatic at the beginning of the pandemic but was not tested due to restricted testing. She developed light symptoms 120 days after the first infection, with a positive RT-PCR test without informed Ct value. We had no information for the last patient except a 142-day interval between the serologic assay and the positive RT-PCR test. None of these individuals had a RT-PCR testing performed before the serologic assay, and none of them developed a severe form of the disease nor were hospitalized. Regarding the two *unlikely* cases, one individual had a first symptomatic episode confirmed by RT-PCR, without informed Ct value. One-hundred and twenty days later she had a positive RT-PCR test performed for asymptomatic screening, with a Ct value of 34. The other *unlikely* individual had a positive RT-PCR test performed for screening 3 days after the serologic assay. Characteristics of individuals with suspected reinfection included in the main or the sensitivity analyses are reported eTable1.

eTable 1. Demographic, clinical and laboratory characteristics of individuals with suspected SARS-CoV-2 reinfection

Patients	Baseline serologic assay		No of days between episodes	No of days between serologic assay and reinfection	Clinical characteristics	Timing of PCT, Ct value	Reinfection? ^a		
	EI	rIFA					OA1	OA2	OA3
Patient 1: female 42 yrs old	0.84	Positive	unknown	185	1st episode: asymptomatic 2nd episode: symptomatic (mild, COVID-19 - like symptoms: fever, headache, sore throat, dry cough, fatigue, dysgeusia, anosmia)	1st episode: not done (asymptomatic) 2nd episode: PCR (Cobas 6800 assay), Ct 17.9	likely	likely	N/A
Patient 2: female 33 yrs old	0.58	Positive	unknown	178	1st episode: asymptomatic 2nd episode: symptomatic (mild, COVID-19 - like symptoms: headache, dysgeusia, anosmia, dyspnea)	1st episode: not done (asymptomatic) 2nd episode: PCR, no Ct value	likely	likely	N/A
Patient 3: female 77 yrs old	0.87	Positive	unknown	169	1st episode: asymptomatic 2nd episode: symptomatic (mild, COVID-19 - like symptoms: fever, headache, fatigue)	1st episode: not done (asymptomatic) 2nd episode: PCR, no Ct value	likely	likely	N/A
Patient 4: female 64 yrs old	2.0	Positive	unknown	142	1st episode: no information 2nd episode: asymptomatic (screening)	no information	likely	likely	N/A
Patient 5: female 23 yrs old	2.83	Positive	120	49	1st episode: symptomatic (mild, COVID-19 - like symptoms) 2nd episode: asymptomatic (screening)	1st episode: PCR, no Ct value 2nd episode: PCR (Cobas 6800), Ct 34	unlikely	likely	unlikely
Patient 6: male 27 yrs old	7.6	Positive	unknown	3	1st episode: no information 2nd episode: asymptomatic (screening)	no information	unlikely	unlikely	N/A
Patient 7: female 68 yrs old	1.42	Positive	120	34	1st episode: symptomatic (mild, COVID-19 - like symptoms: dysgeusia, anosmia, fatigue, strong headaches), positive contact with COVID-19. 2nd episode: symptomatic (light, fatigue)	1st episode: not done (due to testing restriction) 2nd episode: PCR, no Ct value	likely	likely	N/A
Patient 8: female 50 years old	2.42	Negative	unknown	132	1st episode: asymptomatic 2nd episode: symptomatic (light, COVID-19 - like symptoms: dysgeusia, anosmia, headache)	1st episode: not done (asymptomatic) 2nd episode: PCR, no Ct value	likely	likely	N/A
Patient 9: female 42 years old	1.28	Negative	unknown	103	1st episode: asymptomatic 2nd episode: symptomatic (light, COVID-19 - like symptoms: fever, dysgeusia, anosmia, headache)	1st episode: not done (asymptomatic) 2nd episode: PCR, Ct 18.5	likely	likely	N/A
Patient 10: male 34 years old	10.43	Negative	unknown	119	1st episode: asymptomatic 2nd episode: symptomatic (light, COVID-19 - like symptoms: dysgeusia, anosmia, headache)	1st episode: not done (asymptomatic) 2nd episode: PCR, no Ct value	likely	likely	N/A
Patient 11: male 19 years old	1.38	Negative	unknown	150	1st episode: no information 2nd episode: symptomatic (light, COVID-19 - like symptoms: fever, muscle pain, fatigue)	1st episode: not done (asymptomatic) 2nd episode: PCR, Ct 19	likely	likely	N/A
Patient 12: male 49 years old	1.27	Negative	unknown	142	1st episode: asymptomatic 2nd episode: symptomatic (light, COVID-19 - like symptoms: fever, sore throat, headache, muscle pain)	1st episode: not done (asymptomatic) 2nd episode: PCR, Ct 16.6	likely	likely	N/A
Patient 13: male 38 years old	1.6	Negative	unknown	160	1st episode: asymptomatic 2nd episode: symptomatic (light, COVID-19 - like symptoms: fever, headache, dysgeusia, anosmia)	1st episode: not done (asymptomatic) 2nd episode: PCR, Ct 17.9	likely	likely	N/A

Patient 14: male 29 years old	1.19	Negative	unknown	151	1st episode: asymptomatic 2nd episode: symptomatic (light, COVID-19 - like symptoms: rhinitis, headache, dyspnea)	1st episode: not done (asymptomatic) 2nd episode: PCR, Ct 20.6	likely	likely	N/A
Patient 15: male 16 years old	1.62	Negative	no information	153	no information	1st episode: no information 2nd episode: PCR, no Ct value	likely	likely	N/A

^aTwo adjudicators evaluated independently the reinfection probability of each case. Conflicts have been solved by a third adjudicator.

N/A means not applicable, EI means Euroimmun assay IgG ratio; rIFA means recombinant Immunofluorescence assay

eTable 2. Sensitivity analyses

Number of individuals and infections	n=7618, events= 618 reinfections=5	n= 7618, events= 618 reinfections=5	n= 996, events = 52 reinfections=5	n= 1494, events= 159 reinfections=5	n= 1494, events= 161 reinfections=7	n=1470 events=165 reinfections=5
Cox model (no imputation on missing data)	Unmatched univariable	Unmatched multivariable	Matched univariable (1 to 1 ratio)	Matched univariable (2 to 1 ratio)	Matched univariable (2 to 1 ratio)	Matched 2 ^a univariable (2 to 1 ratio)
Hazard ratio (95%CI)	0.11 (0.05, 0.26)	0.10 (0.04, 0.25)	0.10 (0.04, 0.25)	0.06 (0.02, 0.14)	0.08 (0.04, 0.18)	0.06 (0.02, 0.14)
P value	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001

^aNeighbourhood socioeconomic deprivation replace Education level in propensity score matching