

# Parent and child agreement on reported wheeze and its association with FeNO and lung function

Rebeca Mozun<sup>1,2</sup>, Cristina Ardura-Garcia<sup>1</sup>, Eva S. L. Pedersen<sup>1</sup>, Myrofora Goutaki<sup>1,3</sup>, Jakob Usemann<sup>4,5</sup>, Florian Singer<sup>3,6</sup>, Philipp Latzin<sup>3</sup>, Alexander Moeller<sup>4\*</sup>, Claudia E. Kuehni<sup>1,3\*</sup>

\*These authors contributed equally

1. Institute of Social and Preventive Medicine, University of Bern, Switzerland
2. Graduate School for Cellular and Biomedical Sciences, University of Bern, Switzerland
3. Paediatric Respiratory Medicine, Children's University Hospital of Bern, University of Bern, Switzerland
4. Division of Respiratory Medicine, University Children's Hospital Zurich and Childhood Research center, University of Zurich, Switzerland
5. University Children's Hospital Basel (UKBB), Basel, Switzerland
6. PEDNET, paediatric clinical trial unit, Children's University Hospital of Bern, University of Bern, Switzerland

## Online supplementary material

### Table of contents:

<b>Table S1:</b> Original questions in German (A) and English (B) language from the parental questionnaire and children's questionnaire used in the LuftiBus in the school (LUIS) study.....	2
<b>Table S2:</b> Prevalence of parent- and child-reported wheeze and proportion of children with wheeze by groups of agreement in answers from parents and children.....	3
<b>Table S3:</b> Median difference in FeNO and FEV1/FVC and mean difference in FEV1 and FEF25-75 in schoolchildren with wheeze when reported by (A) children, (B) parents, and (C) parents and children, compared to children with no reported wheeze.....	4
<b>Table S4:</b> Difference in median FeNO and FEV1/FVC and mean FEV1 and FEF25-75 in schoolchildren with wheeze between scenario A (child-reported) and B (parent-reported), scenario A and C (parent- and child-reported), and scenario B and C.....	5
<b>Table S5:</b> Difference in median FeNO and FEV1/FVC in schoolchildren with wheeze when reported by children compared to reported by parents stratified by age groups.....	6
<b>Figure S1:</b> Characteristics of parent-child pairs who agreed for yes (1) and for no (2), compared to parent-child pairs who disagreed in their answers.....	7

**Table S1:** Original questions in German (A) and English (B) language from the parental questionnaire and children’s questionnaire used in the LuftiBus in the school (LUIS) study.

	Parental questionnaire	Children’s questionnaire
<b>A) German</b>		
Explanation	Die Fragen im nächsten Teil beziehen sich auf pfeifende oder keuchende Atmung. Darunter verstehen wir Geräusche, die beim Atmen (vor allem beim Ausatmen) aus dem Brustkorb kommen, nicht aus der Nase.	Jetzt fragen wir dich nach der Atmung in den letzten 12 Monaten. Es gibt Kinder, bei denen es manchmal beim Atmen pfeifende oder keuchende Geräusche im Brustkorb gibt.
Wheeze	Hatte Ihr Kind <u>in den letzten 12 Monaten</u> beim Atmen pfeifende oder keuchende Geräusche im Brustkorb? <i>Ja; Nein</i>	Hattest du in den letzten 12 Monaten beim Atmen manchmal pfeifende oder keuchende Geräusche im Brustkorb? <i>Nein; Ja</i>
Exertional wheeze	Hatte Ihr Kind <u>in den letzten 12 Monaten</u> pfeifende oder keuchende Geräusche im Brustkorb während oder nach körperlicher Anstrengung? <i>Ja; Nein</i>	Hattest du in den letzten 12 Monaten jemals pfeifende oder keuchende Geräusche im Brustkorb während oder nach körperlicher Anstrengung (Rennen, Velofahren)? <i>Nein; Ja</i>
Wheeze triggers	Welche der folgenden Situationen hat bei Ihrem Kind in den letzten 12 Monaten Husten oder pfeifende/keuchende Atmung ausgelöst? (Bitte kreuzen Sie alle zutreffenden Antworten an.) <u>Pfeifende/keuchende Atmung</u> Anstrengung (Rennen, Velofahren, etc.) <i>Ja; Nein; Weiss nicht</i> Hausstaub <i>Ja; Nein; Weiss nicht</i> Blüütenstaub (Gras, Heu, Blumen, Bäume) <i>Ja; Nein; Weiss nicht</i> Kontakt mit Tieren <i>Ja; Nein; Weiss nicht</i> Kalte Luft oder Nebel <i>Ja; Nein; Weiss nicht</i> Lautes Lachen <i>Ja; Nein; Weiss nicht</i> Erkältungen <i>Ja; Nein; Weiss nicht</i>	Welche der folgenden Situationen hat bei dir in den letzten 12 Monaten Husten oder pfeifende/keuchende Atmung oder beides ausgelöst?  <u>Pfeifende/keuchende Atmung /Atemnot</u> Anstrengung (Rennen, Velo, etc.) <i>Ja; Nein; Weiss nicht</i> Hausstaub <i>Ja; Nein; Weiss nicht</i> Blüütenstaub (Gras, Heu, Blumen, Bäume) <i>Ja; Nein; Weiss nicht</i> Kontakt mit Tieren <i>Ja; Nein; Weiss nicht</i> Kalte Luft oder Nebel <i>Ja; Nein; Weiss nicht</i> Lautes Lachen <i>Ja; Nein; Weiss nicht</i> Erkältungen <i>Ja; Nein; Weiss nicht</i>
<b>B) English</b>		
Explanation	Questions on wheezing. By “wheeze” we mean breathing that makes a high-pitched sound whistling or squeaking sound from the chest not the throat.	By “wheezing” we mean breathing that makes a high-pitched whistling or squeaking sound from the chest, not the throat.
Wheeze	Has your child had wheezing or whistling in the chest in the last 12 months? <i>Yes; No</i>	Have you had wheezing or whistling in the chest in the last 12 months? <i>No; Yes</i>
Exertional wheeze	In the past 12 months, has your child’s chest sounded wheezy during or after exercise? <i>Yes; No</i>	In the past 12 months, has your chest sounded wheezy during or after exercise? (racing, cycling)? <i>No; Yes</i>
Wheeze triggers	In the past 12 months, did the following things cause coughing or wheezing in your child? (Please tick all applicable answers) <u>Wheeze</u> Exercise (running, biking, etc.) <i>Yes; No; Don’t know</i> House dust <i>Yes; No; Don’t know</i> Pollen (grass, hay, trees) <i>Yes; No; Don’t know</i> Contact with animals <i>Yes; No; Don’t know</i> Cold temperatures or fog <i>Yes; No; Don’t know</i> Intense laughing <i>Yes; No; Don’t know</i> Colds <i>Yes; No; Don’t know</i>	In the past 12 months, did the following things cause you coughing or wheezing? (Please tick all applicable answers) <u>Wheeze</u> Exercise (running, biking, etc.) <i>Yes; No; Don’t know</i> House dust <i>Yes; No; Don’t know</i> Pollen (grass, hay, trees) <i>Yes; No; Don’t know</i> Contact with animals <i>Yes; No; Don’t know</i> Cold temperatures or fog <i>Yes; No; Don’t know</i> Intense laughing <i>Yes; No; Don’t know</i> Colds <i>Yes; No; Don’t know</i>

**Table S2:** Prevalence of parent- and child-reported wheeze and proportion of children with wheeze by groups of agreement in answers from parents and children.

	N	(A)		(B)		(C)							
		Children		Parents		Parents & children agreed for yes		Parents only		Children only		Parents & children agreed for no	
		n	% [95% CI]	n	% [95% CI]	n	%	n	%	n	%	n	%
Wheeze	2982	273	9% [8-10]	236	8% [7-9]	108	4%	128	4%	165	6%	2581	87%
Exertional wheeze	3037	369	12% [11-13]	234	8% [7-9]	127	4%	107	4%	242	8%	2561	84%
Factors that trigger wheeze													
Exercise	2519	275	11% [10-12]	278	11% [10-12]	101	4%	177	7%	174	7%	2067	82%
Colds	2555	126	5% [4-6]	335	13% [12-14]	42	2%	293	11%	84	3%	2136	84%
Pollen	2517	58	2% [2-3]	135	5% [5-6]	26	1%	109	4%	32	1%	2350	93%
Cold air	2482	69	3% [2-4]	65	3% [2-3]	16	1%	49	2%	53	2%	2364	95%
Pets	2494	40	2% [1-2]	58	2% [2-3]	20	1%	38	2%	20	1%	2416	97%
Laughter	2481	61	2% [1-2]	38	2% [2-3]	10	0%	28	1%	51	2%	2392	96%
House dust	2501	54	2% [2-3]	55	2% [2-3]	14	1%	41	2%	40	2%	2406	96%

Wheeze, exertional wheeze and triggers of wheeze refer to the past 12 months. CI: confidence interval.

**Table S3:** Median difference in FeNO and FEV1/FVC and mean difference in FEV1 and FEF25-75 in schoolchildren with wheeze when reported by (A) children, (B) parents, and (C) parents and children, compared to children with no reported wheeze.

	FeNO, ppb [95% CI]			FEV1/FVC, % [95% CI]			FEV1, z-scores [95% CI]			FEF25-75, z-scores [95% CI]		
	Median	Crude diff.#	Adjusted diff.##	Median	Crude diff.#	Adjusted diff.##	Mean	Crude diff.+	Adjusted diff.**	Mean	Crude diff.+	Adjusted diff.**
<b>Wheeze</b>												
(A) Children no	11.6	Ref.	Ref.	86.9	Ref.	Ref.	-0.527	Ref.	Ref.	-0.554	Ref.	Ref.
	19.3	7.7	6.0	83.9	-3.1	-3.2	-0.649	-0.122	-0.101	-0.947	-0.393	-0.385
Children yes		[5.9, 9.5]	[4.4, 7.6]		[-4.3, -2.0]	[-4.4, -2.1]		[-0.254, 0.011]	[-0.232, 0.031]		[-0.537, -0.250]	[-0.528, -0.242]
		***	***		***	***		*			***	***
(B) Parents no	11.6	Ref.	Ref.	86.9	Ref.	Ref.	-0.521	Ref.	Ref.	-0.557	Ref.	Ref.
	22.4	10.8	11.2	83.9	-3.0	-3.1	-0.745	-0.224	-0.200	-0.984	-0.427	-0.417
Parents yes		[8.8, 12.8]	[9.4, 13.0]		[-4.2, -1.8]	[-4.3, -1.9]		[-0.368, -0.080]	[-0.342, -0.057]		[-0.582, -0.273]	[-0.571, -0.263]
		***	***		***	***		**	**		***	***
(C) Parents & children agreed in no	11.4	Ref.	Ref.	87.0	Ref.	Ref.	-0.519	Ref.	Ref.	-0.540	Ref.	Ref.
	16.6	5.3	4.1	85.2	-1.8	-2.0	-0.610	-0.090	-0.084	-0.826	-0.287	-0.284
Parents & children disagreed		[3.5, 7.1]	[2.6, 5.6]		[-2.9, -0.7]	[-3.1, -0.9]		[-0.222, 0.042]	[-0.215, 0.047]		[-0.429, -0.144]	[-0.426, -0.142]
		***	***		**	***					***	***
Parents & children agreed in yes	31.2	20.5	20.7	82.4	-4.6	-4.5	-0.795	-0.275	-0.232	-1.134	-0.594	-0.579
		[17.7, 23.3]	[18.2, 23.1]		[-6.3, -2.9]	[-6.2, -2.8]		[-0.476, -0.075]	[-0.432, -0.033]		[-0.810, -0.378]	[-0.792, -0.361]
		***	***		***	***		**	**		***	***
N		2674			2147			2479			2147	
<b>Exertional wheeze</b>												
(A) Children no	11.6	Ref.	Ref.	86.9	Ref.	Ref.	-0.509	Ref.	Ref.	-0.549	Ref.	Ref.
	16.4	4.8	3.2	85.3	-1.6	-1.8	-0.718	-0.208	-0.162	-0.883	-0.334	-0.311
Children yes		[3.3, 6.3]	[1.9, 4.5]		[-2.6, -0.6]	[-2.8, -0.8]		[-0.324, -0.093]	[-0.278, -0.047]		[-0.461, -0.207]	[-0.439, -0.184]
		***	***		**	***		***	**		***	***
(B) Parents no	11.6	Ref.	Ref.	86.8	Ref.	Ref.	-0.515	Ref.	Ref.	-0.560	Ref.	Ref.
	21.8	10.2	10.4	84.2	-2.6	-3.1	-0.776	-0.261	-0.201	-0.944	-0.383	-0.356
Parents yes		[8.2, 12.2]	[8.7, 12.1]		[-3.8, -1.4]	[-4.3, -1.8]		[-0.403, -0.120]	[-0.342, -0.059]		[-0.539, -0.228]	[-0.512, -0.200]
		***	***		***	***		***	**		***	***
(C) Parents & children agreed in no	11.5	Ref.	Ref.	86.9	Ref.	Ref.	-0.502	Ref.	Ref.	-0.542	Ref.	Ref.
	15.2	3.7	2.0	86.1	-0.8	-1.0	-0.662	-0.160	-0.125	-0.752	-0.211	-0.195
Parents & children disagreed		[2.0, 5.4]	[0.6, 3.5]		[-1.8, 0.3]	[-2.1, 0.0]		[-0.281, -0.039]	[-0.245, -0.004]		[-0.345, -0.077]	[-0.320, -0.061]
		***	**		**	**		**	**		**	**
Parents & children agreed in yes	24.8	13.4	13.3	83.6	-3.3	-3.7	-0.838	-0.336	-0.262	-1.090	-0.549	-0.515
		[10.7, 16.1]	[11.0, 15.6]		[-4.9, -1.8]	[-5.2, -2.1]		[-0.520, -0.153]	[-0.446, -0.078]		[-0.748, -0.349]	[-0.715, -0.314]
		***	***		***	***		***	**		***	***
N		2726			2189			2519			2189	

Wheeze and exertional wheeze refer to the past 12 months. #Unadjusted and ##adjusted by age and sex quantile regression models. +Unadjusted and \*\* adjusted by age and sex linear regression models. P values \* $<0.1$ , \*\*  $<0.05$ , \*\*\* $<0.001$ . CI: confidence interval. FeNO: Fractional exhaled nitric oxide. FEV1: forced expiratory volume in 1 second. FVC: forced vital capacity. FEV1/FVC: forced expiratory volume in 1 second over forced vital capacity. FEF25-75: forced expiratory flow between 25 and 75% of the forced vital capacity. Z-scores for FEV1 and FEF25-75 based on Global Lung Function Initiative references. Z-scores for FEV1 and FEF25-75 based on Global Lung Function Initiative references.

**Table S4:** Difference in median FeNO and FEV1/FVC and mean FEV1 and FEF25-75 in schoolchildren with wheeze between scenario A (child-reported) and B (parent-reported), scenario A and C (parent- and child-reported), and scenario B and C.

		FeNO, ppb [95% CI]			FEV1/FVC, % [95% CI]			FEV1, z-scores [95% CI]			FEF25-75, z-scores[95% CI]		
		Median	Crude diff. <sup>¶</sup>	Adjusted diff. <sup>¶¶</sup>	Median	Crude diff. <sup>¶</sup>	Adjusted diff. <sup>¶¶</sup>	Mean	Crude diff. <sup>‡</sup>	Adjusted diff. <sup>‡‡</sup>	Mean	Crude diff. <sup>‡</sup>	Adjusted diff. <sup>‡‡</sup>
<b>Wheeze</b>													
(A - B)	Children yes	19.3	-3.1 [-8.7, 2.5]	-5.2 [-11.2, 0.7]	83.9	0.0 [-1.5, -1.5]	-0.1 [-1.7, 1.4]	-0.649	0.096 [-0.046, 0.239]	0.099 [-0.065, 0.263]	-0.947	0.037 [-0.127, 0.201]	0.032 [-0.157, 0.221]
	Parents yes	22.4		*	83.9			-0.745			-0.984		
(A - C)	Children yes	19.3	-11.9 [-20.0, -3.8]	-14.7 [-21.7, -7.7]	83.9	1.5 [0.0, 2.9]	1.3 [-0.3, 2.8]	-0.649	0.146 [0.007, 0.284]	0.132 [-0.005, 0.269]	-0.947	0.186 [0.011, 0.362]	0.192 [-0.005, 0.379]
	Parents & children yes	31.2	**	***	82.4	**		-0.795	**	*	-1.134	**	**
(B - C)	Parents yes	22.4	-8.8 [-16.5, -1.1]	-9.5 [-16.1, -2.9]	83.9	1.5 [0.1, 2.9]	1.4 [-0.2, 2.9]	-0.745	0.049 [-0.088, 0.187]	0.033 [-0.108, 0.173]	-0.984	0.149 [-0.016, 0.314]	0.160 [-0.015, 0.201]
	Parents & children yes	31.2	**	**	82.4	**	*	-0.795			-1.134	*	*
<b>Exertional wheeze</b>													
(A - B)	Children yes	16.4	-5.4 [-9.8, -0.9]	-7.2 [-12.4, -2.0]	85.3	1.1 [-0.5, 2.7]	1.2 [-0.3, 2.8]	-0.718	0.058 [-0.070, 0.187]	0.038 [-0.099, 0.175]	-0.883	0.061 [-0.085, 0.207]	0.045 [-0.112, 0.202]
	Parents yes	21.8	**	**	84.2			-0.776			-0.944		
(A - C)	Children yes	16.4	-8.4 [-14.6, -2.2]	-10.1 [-17.4, -2.9]	85.3	1.7 [0.1, 3.4]	1.9 [0.3, 3.4]	-0.718	0.120 [-0.028, 0.269]	0.100 [-0.043, 0.242]	-0.883	0.207 [0.040, 0.375]	0.203 [0.044, 0.363]
	Parents & children yes	24.8	**	**	83.6	**	**	-0.838			-1.090	**	**
(B - C)	Parents yes	21.8	-3.1 [-8.3, 2.2]	-2.9 [-8.6, 2.8]	84.2	0.6 [-0.8, 2.1]	0.6 [-0.8, 2.1]	-0.776	0.062 [-0.059, 0.183]	0.061 [-0.053, 0.176]	-0.944	0.146 [0.009, 0.284]	0.159 [0.019, 0.298]
	Parents & children yes	24.8			83.6			-0.838			-1.090	**	**


Wheeze and exertional wheeze refer to the past 12 months. <sup>¶</sup>Unadjusted difference in medians and 95% confidence intervals (CI) calculated using the percentile bootstrap method with 500 repetitions. <sup>¶¶</sup>Difference in medians adjusted for age and sex with quantile regression models and 95% bootstrap CI. <sup>‡</sup>Unadjusted difference in means and 95% bootstrap CI. <sup>‡‡</sup> Difference in means adjusted for age and sex with linear regression models and 95% bootstrap CI. P values \* <0.1, \*\* <0.05, \*\*\* <0.001. FeNO: Fractional exhaled nitric oxide. FEV1: forced expiratory volume in 1 second. FVC: forced vital capacity. FEV1/FVC: forced expiratory volume in 1 second over forced vital capacity. FEF25-75: forced expiratory flow between 25 and 75% of the forced vital capacity. Z-scores for FEV1 and FEF25-75 based on Global Lung Function Initiative references.

**Table S5:** Difference in median FeNO and FEV1/FVC in schoolchildren with wheeze when reported by children compared to reported by parents stratified by age groups.


		FeNO, ppb [95% CI]			FEV1/FVC, % [95% CI]			FEV1 z-scores [95% CI]			FEV1 z-scores [95% CI]		
		Median	Crude diff. <sup>¶</sup>	Adjusted diff. <sup>¶¶</sup>	Median	Crude diff. <sup>¶</sup>	Adjusted diff. <sup>¶¶</sup>	Mean	Crude diff. <sup>‡</sup>	Adjusted diff. <sup>‡‡</sup>	Mean	Crude diff. <sup>‡</sup>	Adjusted diff. <sup>‡‡</sup>
<b>Wheeze</b>		(A - B)											
<10 years	Children yes	12.6	0.4 [-5.7, 6.4]	-0.3 [-7.3, 6.7]	83.8	0.3 [-4.2, 4.9]	0.5 [-3.7, 4.7]	-0.514	0.146 [-0.187, 0.479]	0.149 [-0.190, 0.489]	-0.789	0.094 [-0.271, 0.458]	0.102 [-0.284, 0.488]
	Parents yes	12.2			83.5			-0.684			-0.882		
≥ 10 years	Children yes	21.0	-7.7 [-14.6, -0.8]	-7.2 [-14.0, -0.4]	83.9	0.0 [-1.7, 1.7]	-0.1 [-1.8, 1.7]	-0.684	0.082 [-0.086, 0.250]	0.084 [-0.079, 0.247]	-0.993	0.019 [-0.170, 0.208]	0.016 [-0.198, 0.230]
	Parents yes	28.7	**	**	83.9			-0.766			-1.012		
<b>Exertional wheeze</b>		(A - B)											
<10 years	Children yes	12.1	-5.8 [-15.0, 3.4]	-5.5 [-15.1, 4.1]	85.8	3.6 [-1.4, 8.6]	0.8 [-4.0, 5.7]	-0.407	0.084 [-0.246, 0.414]	0.093 [-0.246, 0.433]	-0.715	0.141 [-0.227, 0.510]	0.126 [-0.266, 0.518]
	Parents yes	17.9			82.2			-0.491			-0.857		
≥ 10 years	Children yes	16.6	-6.8 [-12.6, -0.9]	-6.9 [-13.1, -0.73]	85.2	0.8 [-1.0, 2.6]	0.8 [-0.8, 2.5]	-0.763	0.055 [-0.085, 0.194]	0.037 [-0.120, 0.194]	-0.911	0.048 [-0.124, 0.219]	0.041 [-0.142, 0.224]
	Parents yes	23.4	**	**	84.4			-0.818			-0.959		

Wheeze and exertional wheeze refer to the past 12 months. <sup>¶</sup>Unadjusted difference in medians and 95% confidence intervals (CI) calculated using the percentile bootstrap method with 500 repetitions. <sup>¶¶</sup>Difference in medians adjusted for age and sex with quantile regression models and 95% bootstrap CI. <sup>‡</sup>Unadjusted difference in means and 95% bootstrap CI. <sup>‡‡</sup>Difference in means adjusted for age and sex with linear regression models and 95% bootstrap CI. P values \* < 0.1, \*\* < 0.05, \*\*\* < 0.001. FeNO: Fractional exhaled nitric oxide. FEV1: forced expiratory volume in 1 second. FVC: forced vital capacity. FEV1/FVC: forced expiratory volume in 1 second over forced vital capacity. FEF25-75: forced expiratory flow between 25 and 75% of the forced vital capacity. Z-scores for FEV1 and FEF25-75 based on Global Lung Function Initiative references.

**Figure S1:** Comparison of characteristics between parent-child agreement groups using multinomial logistic regression adjusted for all factors in the table.



Interpretation of relative risk ratio:



Parent-child agreement outcome groups:

		Parent-reported wheeze	
		yes	no
Child-reported wheeze	yes	(1) Agreed for yes	(ref) Disagreed
	no	(ref) Disagreed	(2) Agreed for no

Wheeze and exertional wheeze refer to the past 12 months. Abbreviations: Wh.=Wheeze; char.=characteristics; CH: Switzerland; quest.: questionnaire; Swiss SEP: Swiss socioeconomic position index. The Swiss SEP is an area-based measure developed by the Swiss Nation cohort that ranges from 0 to 100 and incorporates information on rent per square meter, education and occupation of household heads, and number of persons per household room. The urbanization degree was categorized according to the classification of the Swiss federal office of statistics classification. Graphs created using the STATA command coefplot [1].

Reference:

- Jann B. Plotting regression coefficients and other estimates. *Stata Journal* 2014; 14(4): 708-737.