

Supplemental Material (2). Respiratory parameters of cigarette smoking

Study authors & reference	Inhalation/Exhalation Volume (T_V in mL)	Puff time/volume PT seconds, PV mL	Comments
Bernstein (2004)	591 (mean), 560 (median) Range 413 – 918	PT 1.8 (mean) PT Range 1.6-2.4 PV 43 (mean) PV Range 21-66	Summary of 32 studies before 1988 US Surgeon Report 1988
Tobin et al. (1982b)	841 ± 517* Natural 748 ± 323** Natural 878 ± 431* Cig Holder 815 ± 376** Cig Holder Range 270-1990 mL		10 subjects Non invasive RIP
Tobin et al. (1982a)	790 ± 450 Group Average 460 ± 130 Rest Tidal Vol Range 270-1970 mL	PT 4.5 ± 1.3 Includes Breath Hold	19 subjects Non invasive RIP
Nil et al. (1986)	500 ± 300* Men 600 ± 500** Men 400 ± 300* Women 400 ± 300** Women	PV 42.3 ± 14.5 PV 50.2 ± 16.8 PV 41.4 ± 13.3 PV 47.0 ± 15.8	67 men, 48 women
Woodman et al. (1986)	192-644 Total Inh Smoke 315-919 Total Inh Vol	PT 1.2-2.9	Inert Krypton gas as smoke tracer
Robinson et al. (1992)	828 ± 126 Low Nicotine 845 ± 105 Normal Nicotine		
Charles et al. (2009)	833 ± 279 Inhaled Vol 897 ± 308 Exhaled Vol 500 ± 148 Rest Tidal Vol	1.82 ± 1.16 Inh Time 2.28 ± 0.87 Exh Time	74 subjects Non invasive RIP without Cig Holder
Marian et al. (2009)	702 ± 437* Inh Vol 636 ± 138** Inh Vol 577 ± 329* Exh Vol 655 ± 195** Exh Vol	1.19 ± 0.29*, Inh Time 1.22 ± 0.37**, Inh Time 2.01 ± 0.76*, Exh Time 2.89 ± 0.72*, Exh Time 0.45 ± 0.48*, Breath Hold 0.45 ± 0.57**, Breath Hold PV 44.9 ± 12.3* PV 44.5 ± 10.9**	BAT study 1986 Table 2

Table SM(2). The table lists various inhaled/exhaled volumes and associated puff times and volumes. The term “puff time” (PT) denotes the time taken to draw smoke from the cigarette (puffing) with “puff volume” (PV) denoting the drawn volume before it mixes with air. Volumes in the second column refer to the inhaled mixture of smoke and air unless it is explicitly specified that it refers to the exhaled mixture. The symbols \pm , * and ** respectively denote standard deviation, high and low TAR yields. RIP refers to Respiratory Inductive Plethysmograph, BAT is British American Tobacco.

Supplemental Material (2). Respiratory parameters of cigarette smoking

References

Bernstein, D. M., 2004.

A review of the influence of particle size, puff volume and inhalation pattern on the deposition of cigarette smoke particles in the respiratory tract. *Inhalation toxicology* 16 (10), 675–689.

Charles, Krautter, and Mariner] Charles, F., Krautter, G. R., Mariner, D. C., 2009.

Post-puff respiration measures on smokers of different tar yield cigarettes. *In-halation toxicology* 21 (8), 712–718.

Marian, C., O'Connor, R. J., Djordjevic, M. V., Rees, V. W., Hatsukami, D. K., Shields, P. G., 2009.

Reconciling human smoking behavior and machine smoking patterns: implications for understanding smoking behavior and the impact on laboratory studies. *Cancer Epidemiology and Prevention Biomarkers* 18 (12), 3305–3320.

Nil, Woodson, and Bättig] Nil, R., Woodson, P. P., Bättig, K., 1986.

Smoking behaviour and personality patterns of smokers with low and high co absorption. *Clinical science* 71 (5), 595–603.

Robinson, Pritchard, and Davis] Robinson, J. H., Pritchard, W. S., Davis, R. A., 1992.

Psychopharmacological effects of smoking a cigarette with typical "tar" and carbon monoxide yields but minimal nicotine. *Psychopharmacology* 108 (4), 466–472.

[Tobin, M. J., Jenouri, G., Sackner, M. A., 1982a.

Subjective and objective measurement of cigarette smoke inhalation. *Chest* 82 (6), 696–700.

Tobin, M. J., Schneider, A. W., Sackner, M. A., 1982b.

Breathing pattern during and after smoking cigarettes. *Clinical Science* 63 (5), 473–483.

Woodman, G., Newman, S., Pavia, D., Clarke, S., 1986.

Inhaled smoke volume, puffing indices and carbon monoxide uptake in asymptomatic cigarette smokers. *Clinical Science* 71 (4), 421–427.