

Title of manuscript:

Primary Care for Homeless Veterans: A Systematic Review of the Homeless Patient
Aligned Care Team (HPACT)

Authors:

David Rosenthal MD^{1 2}, Benjamin A. Howell MD, MPH, MHS³, Erin Johnson BA⁴,
Katherine Stemmer Frumento MLS, MBA⁵, Jack Tsai PhD^{6 7}

Running Title:

Systemic Review of Homeless PACTs

Corresponding author:

David Rosenthal, M.D., Yale School of Medicine, 333 Cedar St, New Haven CT 06510,
david.i.rosenthal@yale.edu

Abstract Word Count: 239

Manuscript Word Count (exclusive of abstract, acknowledgements, figures, tables, and references): 3235

Number of tables and figures: 1 table, 1 supplementary appendix

Keywords: homelessness, primary care, Veterans

Number of references: 32

Number of pages consecutively: 41 (including Table, excluding supplementary appendix)

¹ Yale School of Medicine, Department of General Internal Medicine, New Haven CT

² 4catalyzer Inc., Guilford CT

³ Yale School of Medicine, National Clinician Scholars Program, New Haven, CT

⁴ VA National Center on Homelessness Among Veterans, Veterans Health Administration, Office of Homeless Programs, Washington D.C.

⁵ Yale School of Medicine, Harvey Cushing/John Hay Whitney Medical Library, New Haven, CT

⁶ University of Texas Health Science Center at Houston, School of Public Health, San Antonio, TX

⁷ VA National Center on Homelessness Among Veterans, Veterans Health Administration, Office of Homeless Programs, Washington D.C.

Primary Care for Homeless Veterans: A Systematic Review of the Homeless Patient Aligned Care Team (HPACT)

Abstract (*Word Count 239*)

Background: In 2011, the Veterans Health Administration (VHA) implemented homeless-tailored primary care medical home models, called the Homeless Patient Aligned Care Teams (HPACT) to improve care for homeless Veterans. The aim of this study was to describe the existing peer-reviewed literature on HPACTs by systematically reviewing studies published since 2011 to date.

Methods: We conducted a systematic review of peer-reviewed studies published from 2011 to June 2019 to evaluate the literature since the inception of the VHA's Homeless PACT program implementation. We included original research articles evaluating the Homeless PACT and excluded those that did not contain original data.

Results: Of 379 studies screened, 20 studies met our inclusion criteria and were included for analysis. Given wide variability in research designs and outcome measures, a narrative review was conducted. The 20 included studies were categorized into 3 groups: Early HPACT pilot implementations; Association of HPACT clinics with quality and utilization; and Specialized programs within HPACTs. Observational findings suggest reductions in emergency department utilization, improvements in primary care treatment utilization, engagement, and patient experience; but limited rigorous studies exist beyond single site pilots and a few large observational cohort studies.

Discussion: The HPACT model has been successfully implemented in VHA medical centers throughout the country with multiple studies showing increased primary care engagement and improved patient experience; however, further studies are needed about quality, utilization and whether the model can be implemented outside the VHA system.

Registration Number/Funding source: none

Introduction

Homelessness is associated with premature morbidity and mortality at much higher rates compared to those who are housed. (1, 2) Health care for high-need, socially complex patients such as those persons experiencing homelessness is often inadequate, with high rates of acute healthcare utilization in the emergency department (3) decreased utilization of primary care, (4) and poor quality outcomes. (5, 6) The care provided to persons experiencing homelessness is often fragmented and reactive to acute presentations. Primary care, with its longitudinal, comprehensive, relationship-based model, in theory provides a platform for improved care and outcomes among those experiencing homelessness. Primary care has been described as “integrated, accessible health care services by clinicians who are accountable for addressing a large majority of personal health care needs, and practicing in the context of family and community.” (7) However, traditionally structured primary care models often do not meet the needs of socially vulnerable patients experiencing homelessness due to issues related to 1) trust, 2) stigma, and 3) care processes. (8)

In 2009, Secretary of Veterans Affairs (VA) Eric Shinseki unveiled the department’s comprehensive plan to end homelessness among Veterans. (9) Numerous Veterans Health Administration (VHA) programs were developed or expanded in response to this federal initiative. One of those programs was a population-tailored patient-centered medical home (known in VHA as a Patient-Aligned Care Team) for Veterans experiencing homelessness called the Homeless Patient Aligned

Care Team (HPACT), which was implemented in select VA medical centers in 2011. Building on existing VHA primary care services, the intent of the HPACT model is to adapt the standard care delivery model to further integrate and coordinate health and social services care for homeless Veterans, focusing on the highest-risk, highest-need Veterans unable or unwilling to access traditional health care. (10) The HPACT model provides low-barrier, multidisciplinary care featuring open-access, wrap-around services and supports. As designed, the HPACT model distinguishes itself from traditional primary care settings by offering five core elements: 1) enhanced access, 2) integrated services addressing social determinants of health, 3) intensive care management to coordinated with community agencies, 4) specific training for providers on homeless care skills, and 5) data-driven accountable care processes using specific data reports to inform team performance on quality and utilization. Enhanced care entails adopting both an open-access, walk-in care model, reducing traditional barriers to receiving care, and outreach into the community to engage veterans disconnected with VA services. Service integration allows for one-stop, wrap around services to connect patients to mental health and homeless services, which are often co-located, given the often multiple social needs of veterans experiencing homelessness. In addition, primary care staff are able to provide food, clothing assistance, hygiene items, shower and laundry facilities which all serve to meet the continuum of veteran needs.(11)

The program was formally launched in 2011, starting with 32 sites and subsequently expanded to over 60 medical facilities and caring for approximately over 18,000 veterans annually nationwide, and many others who have since “graduated” to traditional VA primary care team—namely for those Veterans who have become stably

housed and were able to navigate traditional VA primary care settings. Demographic data from the first 14,088 patients enrolled in HPACTs shows that the average participant was 53.4 years of age; 8.8% served in the military after September 11, 2001, 4.0% were women, and 11.0% were 65 years or older. (12) In parallel with the implementation and growth of the HPACT model, there have been many studies investigating various aspects of the HPACT model; however, no systematic review of this research in this area has yet been conducted. There is much to be learned from a review of published evaluations of the HPACT program. Specifically, a review would provide a summary and broader understanding about what is known about the HPACT Program, and to identify gaps in our understanding about the program. It may also provide insights and lessons learned for health systems caring for vulnerable populations outside the VA. This is a systematic review of the peer-reviewed literature to date.

Methods

Literature Search

This systematic review was conducted according to Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. (13) We conducted a search using Medline, Embase, PsycINFO, Scopus, Web of Science, and Cochrane databases using the search strategy outlined in the accompanying appendix. Our search was limited to studies published in peer-reviewed literature.

Study Selection

Articles were screened by two reviewers (D.R, B.H.) and included if they (1) evaluated an HPACT, (2) contained original data, and (3) were published after the implementation start of the HPACT program in 2011. As per figure 1 in the accompanying appendix, 379 studies were imported for screening, after removal of duplications, 371 studies were screened by two reviewers (D.R, B.H.) with 20 full text studies included for analysis. Each full-text article was reviewed by a pair of investigators (D.R, B.H). Disagreements were resolved by consensus.

Data Extraction and Narrative Synthesis

For studies that met our inclusion criteria we extracted information on study objective, methodology, intervention, population, and relevant findings. Given the small number of studies and heterogenous methods, we conducted a narrative synthesis to describe our findings. Each article was reviewed in terms of research design, methods, and results. In attempt to pull out common themes across studies, we grouped studies into categories based on a combination of research design and results. These categories were determined by consensus of team members after reviewing the content of the included studies.

Results

Ultimately, 20 studies were included in the review. Given the small number of studies, and the variability in research design and outcomes, we conducted a narrative systematic review of the included studies. We categorized the 20 included studies into

three groups based on research design and content area. These groups were: early HPACT pilot implementations (five studies), association of HPACT Clinics with quality and utilization (nine studies), and specialized programs within HPACTs (six studies).

See Table 1 (Attached)

(1) Early HPACT pilot implementations

Five papers were included that reported early implementation data. Four of these papers present original data from the pilot HPACT (known initially as Homeless-Oriented Primary Care Clinic) from the Providence VA Healthcare System. These Providence studies helped build the foundational evidence base for the creation and expansion of specialized PACT models, and specifically the HPACT model. In one study comparing the HPACT model to other specialized PACT models, O'Toole and colleagues used a quasi-experimental 6 month pre- and post-enrollment study tailoring patient-centered medical home interventions to four different high-risk groups and compared engagement and utilization – one of which was tailored to 71 homeless Veterans. This study demonstrated that a specific, homeless-oriented primary care model was feasible within the VA, and that pre-post comparison found a significant increase in primary care use although it was also accompanied by greater use of emergency department visits as well. (14) Furthermore, a larger randomized controlled trial (RCT), the only RCT to date about HPACTs with published data, by Providence VA researchers suggests that 185 out-of-treatment homeless Veterans could be effectively engaged in primary care, measured as any clinical visit in primary care during the study

period, through a tailored outreach process through a Personal Health Assessment/Brief Intervention Arm (PHA/BI), a Clinic Orientation arm (HPACT), or both (PHA/BI+CO). (15) Both of these interventions significantly increased percentages of Veterans accessing primary care compared to a control arm of usual care, but had small numbers (around 40 in each intervention arm) of participants. O'Toole and colleagues in the Providence VA further observed that 79 Veterans experiencing homelessness had increased utilization of primary, mental health, substance abuse and ED care during the first 6 months of HPACT engagement as compared to 98 Veterans experiencing homelessness seen in traditional primary care teams. (16) They found by tailoring primary care to homeless Veterans there were modest, statistically significant improvements in chronic disease management metrics (blood pressure, diabetes care, hyperlipidemia). (17) Some limitations of these original pilot studies are that they were small (in aggregate there were a total of about 441 non-unique Veterans experiencing homelessness in the four papers), at a single-site (Providence VA Healthcare System), and utilized different methodologies (1 quasi-experimental pre-post study, 1 retrospective cohort, 1 case-control nested cohort, and 1 prospective RCT). Despite these limitations, the data provided a foundational base for the national expansion of the HPACT model.

As the program moved from a single site to multiple sites, ability to assess fidelity to the original model became important. To better understand fidelity, O'Toole and colleagues in 2014 conducted an observational study of the first 33 VHA facilities with HPACTs with over 14,000 patients enrolled at the time. (12) Of the 33 sites studied, 82% provided

hygiene care (on-site showers, hygiene kits, and laundry), 76% provided transportation, 55% had an on-site clothes pantry, and 42% had a food pantry and provided on-site meals or food assistance. Six-month patterns of acute-care use pre-enrollment and post-enrollment for 3,543 consecutively enrolled patients showed a 19.0% reduction in emergency department use and a 34.7% reduction in hospitalizations. Higher performance, as defined by the researchers as having greater reductions in ED and hospitalization use, was significantly associated with 1) higher staffing ratios than other sites, 2) integration of social supports and social services into clinical care, and 3) outreach to and integration with community agencies.

(2) Association of HPACT Clinics with quality and utilization

A plurality of studies (nine) we included studied the association of HPACT clinics with quality of care and utilization of care by homeless Veterans. Four studies were surveys, three were retrospective observational studies, one a prospective quasi-experimental trial, and one a mixed-methodology qualitative interviews with descriptive data. None were prospective randomized control trials. Multiple studies focused on patient experience and engagement. Three studies suggest that care in an HPACT was associated with improved patient experience for homeless Veterans compared to traditional VA care settings (18, 19), especially for those Veterans with severe psychiatric symptoms. (20) Kertesz and colleagues in surveying about 600 homeless-experienced patients in 3 Veteran's Affairs (VA) mainstream primary care settings, a Homeless-PACT VA clinic in California, and a highly tailored non-VA Health Care for the

Homeless Program (HCHP) in Massachusetts reported that highly tailored primary care services were associated with decreased odds of unfavorable experiences in the domains of patient-clinician relationship, cooperation, and access or coordination compared to mainstream VA primary care sites, whereas the VA HPACT site attained intermediate results. (21)

Between 2012 and 2014, O'Toole and colleagues studied 2 VHA medical centers to assess health services use, cost, and satisfaction over 12 months for a cohort of 266 homeless Veterans. (22) They found that compared with PACT patients, HPACT patients had more social work visits (4.6 vs 2.7 visits) and fewer emergency department (ED) visits for ambulatory care-sensitive conditions (0 vs 0.2 visits); a significantly smaller percentage of Veterans in HPACT were hospitalized (23.1% vs 35.4%) or had mental health-related ED visits (34.1% vs 47.6%). They also found significant differences in primary care provider-specific visits (HPACT, 5.1 vs PACT, 3.6 visits), mental health care visits (HPACT, 8.8 vs PACT, 13.4 visits), 30-day prescription drug fills (HPACT, 40.5 vs PACT, 58.8 fills), and use of group therapy (HPACT, 40.1% vs PACT, 53.7%). Annual costs per patient were significantly higher in the PACT group than the HPACT group (\$37,415 vs \$28,036), with higher primary care costs in the HPACT group but less cost from hospitalizations, based on managerial cost accounting estimates.

Regarding the effect of the HPACT model on utilization of care, in addition to the fidelity assessment performed by O'Toole and colleagues, several other studies have been performed. Some studies suggest that increased primary care engagement decreases ED and hospital utilization. Gundlapalli and colleagues showed that primary

care treatment engagement can reduce ED visits, especially in the highest ED utilizers. (23) Jones and colleagues showed at a single HPACT site (Pittsburgh VAMC) that Veterans were more likely to have increased primary care visits and less likely to have ED visits following HPACT engagement. (24) Trivedi and colleagues observed that Veterans enrollment in HPACTs with significant dual use of Medicare and VA outpatient care was strongly associated with acute hospitalizations financed by Medicare and high fragmentation of care. (25) Other studies showed mixed results. Patel provided an early case study in 2013 at the VA greater Los Angeles Healthcare System of 47 Veterans who were assigned to the West LA HPACT for at least six months without any statistically significant change in acute ED utilization after six months despite significant increases in HPACT visits and SW encounters. (26) While limited by different methodologies and their observational nature, overall these studies seem to suggest that HPACTs increase engagement and utilization of primary care, and in the highest of utilizers, may be associated with a reduction in acute ED use and hospitalization.

(3) Specialized programs within HPACTs

As the HPACT model has now been implemented in over 60 sites nationally, there have been innovative pilots and model variation described in the literature. Six studies were included in this category for analysis although all were of lower methodologic rigor. Several studies describe innovative staffing models. Pauly et al describe the integration of a mental health clinical pharmacy specialist in the South Texas HPACT that showed reduction in wait-time for psychiatric pharmacotherapy (27) whereas Yoon describes

the use of Peer Mentors for Veterans experiencing Homelessness within HPACT and Non-HPACT clinics without significant impacts on Veteran utilization or cost. (28) Gabrielian and colleagues discuss the feasibility of a variation on the original HPACT model, describing a co-located HPACT clinic within the Emergency Department of the Greater Los Angeles VA Medical Center rather than in an ambulatory clinic. (29) Other innovations studied and described in the literature include implementation of food insecurity screening initiatives, (30) photo elicitation strategies for recovery, (31) and the MISSION-Vet program within an HPACT. (32) Each of these studies were included in the analysis because they contain original data; showing variation and ongoing innovation within the HPACT clinics.

Discussion

The HPACT model has been implemented in VHA medical centers throughout the country and has great potential to engage Veterans experiencing homelessness in specialized access and services within primary care. Despite this rapid, widespread implementation, limited data exists about the program in the peer-reviewed literature. Our systematic review identified only 20 studies to date that provide an evidentiary base for evaluation of the program. The strongest evidence, obtained through the only randomized controlled trial of the HPACT model showed that it can increase patient engagement and increase primary care utilization. (14) In addition, there is modest observational data demonstrating an association between HPACT engagement and reduction in use of emergency care and acute hospitalization in certain high utilizing

populations. Similarly, one observational study showed an association with HPACT program engagement and annual cost decreases from aforementioned decrease reduction in acute care. Importantly, HPACTs were not necessarily designed to improve housing outcomes per se so there is limited evidence in that respect, although several studies have shown that HPACTs can improve primary care engagement, which in theory could enhance health and housing outcomes. The impact of HPACTs on long-term housing instability has yet to be examined rigorously.

Implication for Primary Care Settings and Health Care Policy

Overall, the HPACT model is a successful example of how an innovative new program can scale nationally within the VA. Within a few years from its single site pilot, over 60 sites nationally employ an HPACT model of care for Veterans experiencing homelessness. The rapid spread of the HPACT program does highlight the shift of framework from a “one-size-fits-all” primary care medical home model to a specialized model where segmentation and specialization of services may provide a slightly different model of care for vulnerable populations. Perhaps most striking from the majority of the studies, HPACTs are associated with an improved patient experience for Veterans experiencing homelessness compared to non-tailored primary care clinics. The implication for this is that better customizing and tailoring care to meet the specialized needs of this population is likely to result in better experience, higher engagement in treatment, and likely further retention in primary care. Additional research is clearly needed to better understand the effect of HPACT within VA, but also to better understand whether such a model could be implemented outside of a VA

system, where other healthcare organizations now have an increased attention on improving social determinants of health.

Limitations and Strengths

This study has significant limitations. First, there may have been bias in our search and study selection strategy. Our search was limited to peer-reviewed publications and therefore excluded non-peer reviewed literature about HPACTs and internal, non-published quality-improvement data about HPACTs that exist such as within the National VA VISN Support Services Center. We attempted to mitigate this bias by conducting our search according to Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines and standard accepted databases, but it is likely that there exists more evidence about the HPACT program outside of the published literature. Furthermore, regarding potential selection bias, the two reviewers of studies both work clinically in HPACT clinics and may have implicitly excluded or interpreted studies towards a positive associations. Ten studies were excluded from analysis because they did not evaluate the HPACT program but evaluated them in combination with other VHA homeless programs such as HUDVASH and supportive housing interventions. It is possible that these exclusions removed important data from the review; however, we limited the scope of this review to interventions specific to HPACTs.

Second, one of the largest challenges with conducting this study was the issue of fidelity to the HPACT model. In almost all of the published studies, is not clear and consistent to the authors what the HPACT team is – i.e. whether there are consistent staffing

ratios, consistent training between sites, and how geographic factors may play an outsized role in accessibility to services. Perhaps most importantly, its not clear that there are predetermined standard performance metrics and outcomes that the HPACTs are seeking to achieve and whether HPACTs consistently use data reports to manage towards these specific outcomes. Due to the significant internal variability of these programs, generalizability of the HPACT model is significantly impaired. Regarding costs, studies suggesting cost benefit of HPACT do not clarify whether they include HPACT staffing costs, and should be noted that the data used for non-VA use and cost are only estimates based on Fee Basis files.

Lastly, the methodology and quality of the studies makes comparison of studies challenging. In particular, numerous observational trials were subject to multiple confounders, in particular, the significant growth of homeless services in VA at the same time as the implementation of the HPACT programs, that may increase patient engagement to VA services independently of the HPACT program.

Conclusion

People experiencing homelessness are high utilizers of high cost services in our health care system and unfortunately also have high levels of morbidity compared to the non-homeless population. The VHA should be applauded for implementing a specialized primary care model, the HPACT, to address these issues for veterans experiencing homelessness. Our review shows that the HPACT model has been successfully implemented in VHA medical centers throughout the country with multiple studies

showing increased primary care engagement and improved patient experience.

However, to date, published studies evaluating the HPACT model are limited and further studies are needed to refine our understanding of the impact that HPACTs have on quality, utilization and whether the model can be implemented outside the VHA system.

Table 1. HPACT papers included in analysis

Category	Title	Author	Year	Journal	Objective	Method	Results	Conclusions
Early HPACT pilot implementations	Building care systems to improve access for high-risk and vulnerable veteran populations.	O'Toole TP, et al.	2011	Journal of General Internal Medicine	Examine health outcomes by tailoring medical home model for high-risk patients, accessing primary care.	A quasi-experimental pre-post study examining a vulnerable population-based application of the patient-centered medical home applied to four high-risk groups: (1) homeless veterans (2) cognitively impaired elderly (3) women veterans and (4) patients with serious mental illness.	Overall 457 patients were studied, assessing care use and outcomes for the last 6 months in each study year. Compared with 2006, in 2008 there was a significant increase in primary care use ($p < 0.001$) and improvement in chronic disease monitoring and diabetes control (2006 HBA1C: 8.5 vs. 2008 HBA1C 6.9) in all four groups. However, there was also an increase in both emergency department use and hospitalizations. Most of the increased utilization was driven by a small proportion of patients in each group.	Tailoring the medical home model to the specific needs and challenges facing high-risk populations can increase primary care utilization and improve chronic disease monitoring and diabetes management.

<p>Early HPACT pilot implementations</p>	<p>Applying the chronic care model to homeless veterans: effect of a population approach to primary care on utilization and clinical outcomes.</p>	<p>O'Toole TP, et al.</p>	<p>2010</p>	<p>American Journal of Public Health</p>	<p>Compare a population-tailored approach to primary care for homeless veterans with a usual care approach.</p>	<p>A retrospective prospective cohort study of homeless veterans enrolled in a population-tailored primary care clinic matched to a historical sample in general internal medicine clinics. Overall, 177 patients were enrolled: 79 in the Homeless-Oriented Primary Care Clinic and 98 in general internal medicine primary care.</p>	<p>Homeless-oriented primary care-enrolled patients had greater improvements in hypertension, diabetes, and lipid control, and primary care use was higher during the first 6 months (5.96 visits per person vs. 1.63 for general internal medicine) but stabilized to comparable rates during the second 6 months (2.01 vs. 1.31, respectively). Emergency department (ED) use was also higher (2.59 vs. 1.89 visits), although with 40% lower odds for non-acute ED visits than for the general internal medicine group (95% confidence interval = 0.2, 0.8). Excluding substance abuse and mental health admissions, hospitalizations were reduced among the homeless veterans between the 2 periods (28.6% vs. 10.8%; $P < .01$) compared with the general internal medicine group (48.2% vs. 44.4%; $P = .6$; difference of differences, $P < .01$).</p>	<p>Tailoring primary care to homeless veterans can decrease unnecessary ED use and medical admissions and improve chronic disease management.</p>
--	--	---------------------------	-------------	--	---	--	--	---

<p>Early HPAC T pilot implementations</p>	<p>New to care: demands on a health system when homeless veterans are enrolled in a medical home model</p>	<p>O'Toole, TP, et al.</p>	<p>2013</p>	<p>American Journal of Public Health</p>	<p>Compare service use among homeless and nonhomeless veterans newly enrolled in a medical home model and identified patterns of ED use.</p>	<p>Case-control matching with a nested cohort analysis to measure 6-month health services use, new diagnoses, and care use patterns in veterans at the Providence, Rhode Island, Veterans Affairs Medical Center from 2008 to 2011. Followed 127 homeless and 106 nonhomeless veterans.</p>	<p>Homeless veterans used significantly more primary, mental health, substance abuse, and ED care during the first 6 months. Homeless veterans who accessed primary care at higher rates (relative risk ratio [RRR] = 1.46; 95% confidence interval [CI] = 1.11, 1.92) or who used specialty and primary care (RRR = 10.95; 95% CI = 1.58, 75.78) had reduced ED usage. Homeless veterans in transitional housing or doubled-up at baseline (RRR = 3.41; 95% CI = 1.24, 9.42) had similar reductions in ED usage.</p>	<p>Homeless adults had substantial health needs when presenting for care. High-intensity primary care and access to specialty care services could reduce ED use.</p>
<p>Early HPAC T pilot implementations</p>	<p>Tailoring Outreach Efforts to Increase Primary Care Use Among Homeless Veterans: Results of a Randomized Controlled Trial.</p>	<p>O'Toole, TP, et al.</p>	<p>2015</p>	<p>Journal of General Internal Medicine</p>	<p>Increase health-seeking behavior and receipt of health care among homeless Veterans</p>	<p>Multi-center, prospective, community-based, two-by-two randomized controlled trial of homeless Veterans. An outreach intervention that included a personal health assessment and brief intervention (PHA/BI), and/or a clinic orientation (CO) was implemented. Overall, 185 homeless Veterans were enrolled.</p>	<p>At one month, 77.3% of the PHA/BI plus CO group accessed primary care and by 6 months, 88.7% had been seen in primary care. This was followed by the CO-only group, 50.0% of whom accessed care in the first 4 weeks, the PHI/BI-only arm at 41.0% and the Usual Care arm at 30.6%. Chi-squared tests by group were significant ($p < 0.001$) at both 4 weeks and 6 months. There was no difference in attitudes about care at baseline and 6 months or in use patterns once enrolled in care.</p>	<p>Treatment-resistant/avoidant homeless Veterans can be effectively engaged in primary and other clinical care services through a relatively low intensity, targeted and tailored outreach effort.</p>

<p>Early HPACT pilot implementations</p>	<p>Tailoring Care to Vulnerable Populations by Incorporating Social Determinants of Health: the Veterans Health Administration's "Homeless Patient Aligned Care Team" Program.</p>	<p>O'Toole, TP, et al.</p>	<p>2016</p>	<p>Preventing Chronic Disease</p>	<p>Describe the national implementation of HPACT initiative in VHA and correlate patient health outcomes with characteristics of high-performing sites.</p>	<p>An observational study of 33 VHA facilities with homeless medical homes and patient-aligned care teams that served more than 14,000 patients. Site-specific health care performance data for the 3,543 homeless veterans enrolled in the program from October 2013 through March 2014, 6 months prior to enrollment and 6 months post-enrollment with corresponding survey data on the Homeless Patient Aligned Care Team (HPACT) program implementation.</p>	<p>Of the 33 sites studied, 82% provided hygiene care (on-site showers, hygiene kits, and laundry), 76% provided transportation, and 55% had an on-site clothes pantry; 42% had a food pantry and provided on-site meals or other food assistance. Six-month patterns of acute-care use pre-enrollment and post-enrollment for 3,543 consecutively enrolled patients showed a 19.0% reduction in emergency department use and a 34.7% reduction in hospitalizations. Three features were significantly associated with high performance: 1) higher staffing ratios than other sites, 2) integration of social supports and social services into clinical care, and 3) outreach to and integration with community agencies.</p>	<p>Integrating social determinants of health into clinical care can be effective for high-risk homeless veterans.</p>
--	--	----------------------------	-------------	-----------------------------------	---	--	--	---

<p>Association of HPACT Clinics with quality and utilization</p>	<p>Comparing homeless persons' care experiences in tailored versus nontailored primary care programs.</p>	<p>Kertesz SG, et al.</p>	<p>2013</p>	<p>American Journal of Public Health</p>	<p>Compare homeless patients' experiences of care in health care organizations that differed in their degree of primary care design service tailoring.</p>	<p>A multi-site survey of homeless-experienced patients (either recently or currently homeless) at 3 Veterans Affairs (VA) mainstream primary care settings in Pennsylvania and Alabama, a homeless-tailored VA clinic in California, and a highly tailored non-VA Health Care for the Homeless Program in Massachusetts (January 2011-March 2012).</p>	<p>Mean scores at the tailored non-VA site were superior to those from the 3 mainstream VA sites ($P < .001$). Adjusting for patient characteristics, these differences remained significant for subscales assessing the patient-clinician relationship ($P < .001$) and perceptions of cooperation among providers ($P = .004$). There were 1.5- to 3-fold increased odds of an unfavorable experience in the domains of the patient-clinician relationship, cooperation, and access or coordination for the mainstream VA sites compared with the tailored non-VA site; the tailored VA site attained intermediate results.</p>	<p>Tailored primary care service design was associated with a superior service experience for patients who experienced homelessness.</p>
--	---	---------------------------	-------------	--	--	---	--	--

<p>Association of HPACT Clinics with quality and utilization</p>	<p>Experience of Primary Care among Homeless Individuals with Mental Health Conditions</p>	<p>Chrystal JG, et al</p>	<p>2015</p>	<p>PLoS One</p>	<p>Study predictors of favorable primary care experience among homeless persons with mental health conditions treated tailoring.</p>	<p>A multi-site, survey-based comparison of primary care experiences at three mainstream primary care clinics of the Veterans Administration (VA), one homeless-tailored VA clinic, and one tailored non-VA healthcare program. Used Primary Care Quality-Homeless (PCQ-H) questionnaire, a validated survey instrument.</p>	<p>Significant predictors of a positive experience included a site offering tailored service design, perceived choice among providers, and currently domiciled status. For persons with severe psychiatric symptoms, a homeless-tailored service design was significantly associated with a more favorable primary care experience. For persons without severe psychiatric symptoms, this difference was not significant.</p>	<p>This study supports the importance of tailored healthcare delivery designed for homeless persons' needs, with such services potentially holding special relevance for persons with mental health conditions. To improve patient experience among the homeless, organizations may want to deliver services that are tailored to homelessness and offer a choice of providers.</p>
--	--	---------------------------	-------------	-----------------	--	--	---	---

<p>Association of HPACT Clinics with quality and utilization</p>	<p>Providing Positive Primary Care Experiences for Homeless Veterans Through Tailored Medical Homes</p>	<p>Jones, A, et al.</p>	<p>2019</p>	<p>Medical Care</p>	<p>Determine whether HPACT offers a better patient experience than standard VHA primary care.</p>	<p>A National survey. Multivariable logistic regressions to estimate differences in the probability of reporting positive primary care experiences. Homeless-experienced survey respondents enrolled in HPACT (n=251) or standard primary care in facilities with HPACT available (n=1527) and facilities without HPACT (n=10,079).</p>	<p>HPACT patients were more likely than standard primary care patients in the same facilities to report positive experiences with access [adjusted risk difference (RD)=17.4], communication (RD=13.9), office staff (RD=13.1), provider ratings (RD=11.0), and comprehensiveness (RD=9.3). Standard primary care patients in facilities with HPACT available were more likely than those from facilities without HPACT to report positive experiences with communication (RD=4.7) and self-management support (RD=4.6).</p>	<p>Patient-centered medical homes designed to address the social determinants of health offer a better care experience for homeless patients, when compared with standard primary care approaches.</p>
--	---	-------------------------	-------------	---------------------	---	---	--	--

<p>Association of HPACT Clinics with quality and utilization</p>	<p>Complex care management to decrease emergency department utilization: A case study of the homeless patient aligned care team demonstration project at VA greater Los Angeles healthcare system</p>	<p>Patel, BI</p>	<p>2014</p>	<p>Dissertation Abstracts International : Section B: The Sciences and Engineering</p>	<p>Undertake a local needs assessment to determine the organizational context for the WLA HPACT program and how best to evaluate HPACT's objective to reduce ED utilization rates.</p>	<p>A quality improvement (QI) dissertation case study of the homeless-oriented Patient Centered Medical Home demonstration program, referred to as the Homeless Patient Aligned Care Team (HPACT) at West Los Angeles VA Medical Center (WLA).</p>	<p>The implementation phase of this QI dissertation employed a mixed-method progress-focused formative evaluation to examine patient characteristics, intervention processes, ED utilization patterns, staff perceptions on perceived barriers and interventions to address gaps in care to decrease ED visits for current HPACT patients. The evaluation used qualitative key informant interviews and quantitative descriptive data analysis.</p>	<p>Recommendations include complex care management interventions for patients with pain needs, substance abuse conditions, as well as modifications to organizational features of the program such as hours of operations, and team composition.</p>
--	---	------------------	-------------	---	--	--	---	--

<p>Association of HPACT Clinics with quality and utilization</p>	<p>Patient-aligned Care Team Engagement to Connect Veterans Experiencing Homelessness With Appropriate Health Care</p>	<p>Gundlapalli AV, et al</p>	<p>2017</p>	<p>Medical Care</p>	<p>Examine the effect of HPACT on frequency and type of ED visits in Veterans Health Administration (VHA) medical facilities.</p>	<p>An observational study. During a 12-month period, ED visits for 3981 homeless Veterans enrolled in (1) HPACT at 20 VHA medical centers (enrolled) were compared with those of (2) 24,363 homeless Veterans not enrolled in HPACT at the same sites (nonenrolled), and (3) 23,542 homeless Veterans at 12 non-HPACT sites (usual care) using a difference-in-differences approach.</p>	<p>In comparing HPACT enrollees with usual care, there was a significant decrease in ED usage among the highest ED utilizers (difference-in-differences, 4.43; $P < 0.001$). The decrease in ED visits were significant though less intense for HPACT enrollees versus nonenrolled (0.29, $P < 0.001$). HPACT enrollees demonstrated a significant increase in the proportion of ED care visits that were not preventable/avoidable in the 6 months after enrollment, but had stable rates of primary care, mental health, social work, and substance abuse visits over the 12 months.</p>	<p>Primary care treatment engagement can reduce ED visits and increase appropriate use of ED services in VHA for Veterans experiencing homelessness, especially in the highest ED utilizers.</p>
--	--	------------------------------	-------------	---------------------	---	--	--	--

<p>Association of HPACT Clinics with quality and utilization</p>	<p>Patient predictors and utilization of health services within a medical home for homeless persons</p>	<p>Jones AL, et al.</p>	<p>2018</p>	<p>Substance Abuse</p>	<p>Compare the number and type of health care visits in the 12 months before and after enrollment in HPACT at one VHA facility.</p>	<p>Chart reviews of VHA medical records were conducted for all patients enrolled in an HPACT in Pittsburgh, Pennsylvania, between May 2012 and December 2013 (N = 179).</p>	<p>Compared with 0-6 months prior to HPACT, patients were more likely to visit primary care in the 0-6 months (adjusted odds ratio [aOR] = 4.91, 95% confidence interval [CI] = 2.94-8.20) and 7-12 months (aOR = 2.30, 95% CI = 1.42-3.72) following HPACT. Patients were less likely to visit the emergency department (ED) or to be hospitalized in the 0-6 months (aOR = 0.57, 95% CI = 0.34-0.94; and aOR = 0.55, 95% CI = 0.25-0.76) and 7-12 months (aOR = 0.43, 95% CI = 0.33-0.91; and aOR = 0.45, 95% CI = 0.26-0.80) following HPACT. Patients were less likely to visit mental health (aOR = 0.35, 95% CI = 0.20-0.60) and addiction specialists (aOR = 0.39, 95% CI = 0.18-0.84) in the 7-12 months following HPACT. Overall, 59% of patients had increases in primary care visits following HPACT.</p>	<p>An integrated HPACT model was successful in engaging homeless veterans in primary care for 1 year, potentially contributing to reductions in ED use. More tailored approaches may be needed for vulnerable populations experiencing homelessness, including homeless women.</p>
--	---	-------------------------	-------------	------------------------	---	---	--	--

<p>Association of HPACT Clinics with quality and utilization</p>	<p>Dual Use and Hospital Admissions among Veterans Enrolled in the VA's Homeless Patient Aligned Care Team</p>	<p>Trivedi AN, et al</p>	<p>2018</p>	<p>Health Services Research</p>	<p>Examine the association between reliance on VA outpatient care and hospital admissions among Medicare-eligible Veterans enrolled in the Homeless Patient Aligned Care Team (HPACT)</p>	<p>National registry of HPACT enrollees linked to VA and Medicare utilization data for 2013.</p>	<p>Compared with higher reliance Veterans, lower reliance Veterans had an equivalent number of annual VA hospitalizations (0.63 vs. 0.50; $p = .14$) but substantially greater Medicare hospitalizations (0.85 vs. 0.08; $p < .001$). Among Veterans in the highest tertile of outpatient visits, we observed statistically similar rates of VA hospital use but over 10-fold greater rates of Medicare-financed hospitalizations (1.31 for lower reliance vs. 0.15 for high reliance; $p < .001$).</p>	<p>Among Veterans receiving integrated care in VA's HPACT, dual use of Medicare and VA outpatient care is strongly associated with acute hospitalizations financed by Medicare. Linking VA and non-VA data may identify a subset of homeless Veterans with fragmented outpatient care who are at increased risk of poor outcomes.</p>
--	--	--------------------------	-------------	---------------------------------	---	--	--	---

<p>Association of HPACT Clinics with quality and utilization</p>	<p>Differences in Experiences With Care Between Homeless and Nonhomeless Patients in Veterans Affairs Facilities With Tailored and Nontailored Primary Care Teams</p>	<p>Jones, A, et al.</p>	<p>2018</p>	<p>Medical Care</p>	<p>Assess differences in primary care experiences between homeless and nonhomeless Veterans receiving care in HPACT facilities and non-HPACT facilities.</p>	<p>National VHA survey. Compared the homeless versus nonhomeless risk differences (RDs) in reporting negative or positive experiences in 25 HPACT facilities versus 485 non-HPACT facilities. Survey respondents from non-HPACT facilities (homeless: n=10,148; nonhomeless: n=309,779) and HPACT facilities (homeless: n=2022; nonhomeless: n=20,941).</p>	<p>In non-HPACT facilities, homeless patients reported more negative and fewer positive experiences than nonhomeless patients. However, these patterns of homeless versus nonhomeless differences were reversed in HPACT facilities for the domains of communication (positive experience RDs in non-HPACT versus HPACT facilities =-2.0 and 2.0, respectively); comprehensiveness (negative RDs=2.1 and -2.3), shared decision-making (negative RDs=1.2 and -1.8), and self-management support (negative RDs=0.1 and -4.5; positive RDs =0.5 and 8.0).</p>	<p>VHA facilities with HPACT programs appear to offer a better primary care experience for homeless versus nonhomeless Veterans, reversing the pattern of relatively poor primary care experiences often associated with homelessness.</p>
--	---	-------------------------	-------------	---------------------	--	---	---	--

<p>Association of HPACT Clinics with quality and utilization</p>	<p>Population-Tailored Care for Homeless Veterans and Acute Care Use, Cost, and Satisfaction: A Prospective Quasi-Experimental Trial</p>	<p>O'Toole TP, et al.</p>	<p>2018</p>	<p>Preventing Chronic Disease (CDC)</p>	<p>Examine the impact of enrollment in HPACT on health services use, cost, and satisfaction compared with enrollment in a traditional, nontailored PCMH.</p>	<p>A prospective, multicenter, quasi-experimental, single-blinded study at 2 VHA medical centers to assess health services use, cost, and satisfaction during 12 months among 2 groups of homeless veterans: 1) veterans receiving HPACT and 2) veterans receiving traditional primary care services (PACT). A cohort of 266 homeless veterans enrolled from June 2012 through January 2014.</p>	<p>Compared with PACT patients, HPACT patients had more social work visits (4.6 vs. 2.7 visits) and fewer emergency department (ED) visits for ambulatory care-sensitive conditions (0 vs. 0.2 visits); a significantly smaller percentage of veterans in HPACT were hospitalized (23.1% vs. 35.4%) or had mental health-related ED visits (34.1% vs. 47.6%). Significant differences in primary care provider-specific visits (HPACT, 5.1 vs. PACT, 3.6 visits), mental health care visits (HPACT, 8.8 vs. PACT, 13.4 visits), 30-day prescription drug fills (HPACT, 40.5 vs. PACT, 58.8 fills), and use of group therapy (HPACT, 40.1% vs. PACT, 53.7%). Annual costs per patient were significantly higher in the PACT group than the HPACT group (\$37,415 vs. \$28,036).</p>	<p>Homeless veterans enrolled in HPACT used less acute care and costs were lower.</p>
--	--	---------------------------	-------------	---	--	--	--	---

Specialized programs within HPACTs	Feasibility and Acceptability of a Colocated Homeless-Tailored Primary Care Clinic and Emergency Department.	Gabieli an, S, et al.	2017	Journal of Primary Care & Community Health	Study feasibility and acceptability of a pilot initiative to colocate a homeless-tailored PCMH with an ED.	Single-site survey of patients and ED clinicians (physicians, nurses, social workers); from May to September 2012.	281 patients were screened in ED; 172 (61.2%) screened positive for homelessness; Acceptability data were obtained from 56% (n = 32) of surveyed clinicians. Most (54.6%) clinicians agreed with the homelessness screening procedures. Nearly all (90.3%) clinicians supported expansion of the homeless-tailored clinic; a minority (42.0%) agreed that ED colocation worked well.	Suggest feasibility of recruiting patients to a homeless-tailored primary care clinic colocated with the ED; however, the clinic's acceptability was mixed.
------------------------------------	--	-----------------------	------	--	--	--	--	---

<p>Specialized programs within HPACTs</p>	<p>Screening for Food Insecurity in Six Veterans Administration Clinics for the Homeless, June–December 2015</p>	<p>O'Toole TP, et al.</p>	<p>2017</p>	<p>Preventing Chronic Disease (CDC)</p>	<p>Assess findings from a food-insecurity screening of a national sample of Veterans Administration clinics for homeless and formerly homeless veterans.</p>	<p>Multisite screenings. Results from initial screenings administered at 6 Veterans Administration primary care clinics for the homeless and responses from clinic staff members interviewed about the screening program.</p>	<p>A total of 270 patients were screened. Screening showed a high prevalence of food insecurity. Of the 270, 48.5% reported they experienced food insecurity in the previous 3 months, 55.0% reported averaging 2 meals a day, and 27.3% averaged 1 meal a day. Eighty-seven percent prepared their own meals, relying on food they bought (54.2%), help from friends and family (19.1%), and soup kitchens and food pantries (22%); 47.3% received Supplemental Nutrition Assistance Program benefits (food stamps). Additionally, of those who screened positive for food insecurity 19.8% had diabetes or prediabetes, and 43.5% reported hypoglycemia symptoms when without food. Clinic staff members responded positively to the screening program and described it as a good rapport builder with patients</p>	<p>Integrating screening for food insecurity among patients in clinical settings was well received by both patients and health care providers. Addressing these positive findings of food insecurity requires a multidisciplinary health care approach.</p>
---	--	---------------------------	-------------	---	--	---	---	---

Specialized programs within HPACTs	Homeless Veterans' Use of Peer Mentors and Effects on Costs and Utilization in VA Clinics.	Yoon J, et al.	2017	Psychological Services	Compare health care utilization and costs among homeless veterans randomly assigned to peer mentors or usual care and described contacts with peer mentors.	Homeless patients at four Department of Veterans Affairs clinics were randomly assigned to a peer mentor (N=195) or to usual care (N=180). Administrative data on utilization and costs over a six-month follow-up were combined with peer mentors' reports of patient contacts.	Most patients (87%) in the peer mentor group had at least one peer contact. Patients in this group spent the largest proportions of time discussing housing and health issues with peer mentors and had more outpatient encounters than those in usual care, although differences were not significant. No other between-group differences were found in utilization or costs.	Although significant impacts of peer mentors on health care patterns or costs were not detected, some patients had frequent contact with peer mentors.
Specialized programs within HPACTs	Integrating a mental health clinical pharmacy specialist into the Homeless Patient Aligned Care Teams.	Pauly JB, et al.	2018	Mental Health Clinician	Assess the impact of a clinical pharmacy specialist in mental health (MH) embedded within an HPACT.	Single site South Texas. An HPACT MH pharmacy resident clinic was created and managed by a postgraduate year-2 psychiatric pharmacy resident.	Over the 6-month clinic time frame, there were a total of 40 patient encounters in which 21 veterans had MH medication evaluation on at least 1 occasion. The average wait time for Veterans previously followed by the HPACT psychiatrist was approximately 8 weeks. The HPACT MH pharmacy resident clinic enabled veterans to be evaluated every 4 to 6 weeks. Estimated cost savings from clinic interventions totaled \$33 613.67.	The HPACT MH pharmacy resident clinic allowed for an improvement in wait time for psychiatric pharmacotherapy follow-up for homeless veterans, with interventions that were associated with significant estimated cost savings.

Specialized programs within HPACTs	Homeless veterans' experiences with substance use, recovery, and treatment through photo elicitation.	Sestito, S. F., et al. (2017)	2017	Substance Abuse	Describe HPACT veterans' experiences with substance use (SU), substance use recovery (SUR), and substance use treatment (SUT) using photo elicitation.	Homeless veterans were recruited from a single site veteran primary care medical home clinic between September 2014 and March 2015. Twenty veterans were given digital cameras and prompts for taking photographs about their health and health care and participated in 2 photo elicitation interviews.	The majority of participants (75%, n = 15) discussed SU, SUR, and/or SUT in regards to their health and health care utilization. SU themes centered on disclosure of addiction or dependency; substances used; repercussions of SU; SU as a coping mechanism; and association of SU with military service. SUR themes included disclosure of length of sobriety; perceived facilitators of SUR in health, beliefs, social, environmental, financial, and creative pursuit domains; and perceived barriers to SUR in beliefs, social, and environmental domains. SUT themes focused on perceived facilitators of SUT in access to Department of Veterans Affairs (VA) and non-VA services and social domains and perceived barriers to SUT in the social domain.	Providers seeking to elicit addiction-related clinical history and facilitate SUR and SUT might look to the current findings for guidance. Provider training in motivational interviewing may be warranted, which allows for an exploration of health-related consequences of SU and supports patients' self-efficacy.
------------------------------------	---	-------------------------------	------	-----------------	--	--	---	--

<p>Specialized programs within HPACTs</p>	<p>A Hybrid III stepped wedge cluster randomized trial testing an implementation strategy to facilitate the use of an evidence-based practice in VA Homeless Primary Care Treatment Programs.</p>	<p>Simmons, MM, et al.</p>	<p>2017</p>	<p>Implementing Independence and Sobriety through Systems Integration, Outreach, and Networking-Veterans Edition) within HPACT.</p>	<p>Identify an appropriate implementation strategy for MISSION-Vet (Maintaining Independence and Sobriety through Systems Integration, Outreach, and Networking-Veterans Edition) within HPACT.</p> <p>Type III Hybrid modified stepped wedge implementation comparing IU to Facilitation across seven HPACT teams in three sites in the greater Los Angeles VA system. This is a cluster-randomized trial.</p>	<p>An implementation science study for potential roll out of MISSION-Vet within HPACTs. No results reported.</p>	<p>Integrating MISSION-Vet within HPACT has the potential to improve the health of thousands of veterans, but it is crucial to implement the intervention appropriately in order for it to succeed. No results reported.</p>
---	---	----------------------------	-------------	---	---	--	--

- (1) Hwang SW, Orav J, O'Connell JJ, et al. Causes of death in homeless adults in Boston. *Ann Intern Med.* 1997;126(8):625---628.
- (2) Baggett TP, Hwang SW, O'Connell JJ, et al. Mortality among homeless adults in Boston: shifts in causes of death over a 15-year period. *JAMA Intern Med.* 2013; 173(3):189---195.
- (3) Kushel MB, Perry S, Bangsberg D, et al. Emergency department use among the homeless and marginally housed: results from a community-based study. *Am J Public Health.* 2002;92(5):778---784
- (4) Kushel MB, Vittinghoff E, Haas JS. Factors associated with the health care utilization of homeless persons. *JAMA.* 2001;285(2):200–6
- (5) Institute of Medicine/Committee on Health Care for Homeless People. *Homelessness, Health, and Human Needs.* Washington, DC: National Academy Press; 1988
- (6) O'Toole TP, Gibbon JL, Hanusa BH, et al. Utilization of health care services among subgroups of urban homeless. *J Health Polit Policy Law.* 1999;24(1): 91–114

(7) Donaldson MS, Vanselow NA, KD Yordy, editors. Institute of Medicine Committee on the Future of Primary Care. Defining primary care: an interim report. Washington, DC: National Academies Press; 1994

(8) O'Toole TP, Johnson EE, Redihan S, et al. Needing primary care but not getting it: the role of trust, stigma and organizational obstacles reported by homeless Veterans. J Health Care Poor Underserved. 2015;26(3):1019–1031.

(9) US Department of Veterans Affairs, Office of Public and Intergovernmental Affairs. Secretary Shinseki details plan to end homelessness for Veterans.
<http://www.va.gov/opa/pressrel/pressrelease.cfm?id=1807> Accessed 1/7/2020

(10) O'Toole TP, Buckel L, Bourgault C, et al. Applying the chronic care model to homeless Veterans: effect of a population approach to primary care on utilization and clinical outcomes. American journal of public health 2010;100:2493-9.

(11) O'Toole TP, Johnson EE, Aiello R, et al. Tailoring Care to Vulnerable Populations by Incorporating Social Determinants of Health: the Veterans Health Administration's "Homeless Patient Aligned Care Team" Program. Prev Chronic Dis 2015;30(7):886–98.

(12) O'Toole TP, Johnson EE, Aiello R, et al. Tailoring care to vulnerable populations by incorporating social determinants of health: the Veterans health administration's "Homeless Patient Aligned Care Team" program. *Prev Chronic Dis.* 2016;13:150567.

(13) Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009; 339: b2535.; *Annals of Internal Medicine.* 151. 264. 10.7326/0003-4819-151-4-200908180-00135.

(14) O'Toole TP, Pirraglia PA, Dosa D, et al. Building care systems to improve access for high-risk and vulnerable Veteran populations. *J Gen Intern Med.* 2011;26 Suppl 2(Suppl 2):683–688. doi:10.1007/s11606-011-1818-2

(15) O'Toole TP, Johnson EE, Borgia ML, et al. Tailoring outreach efforts to increase primary care use among homeless Veterans: results of a randomized controlled trial. *J Gen Intern Med.* 2015;30:886–898.

(16) O'Toole TP, Bourgault C, Johnson EE, et al. New to care: demands on a health system when homeless Veterans are enrolled in a medical home model. *Am J Public Health.* 2013;103 Suppl 2(Suppl 2):S374–S379. doi:10.2105/AJPH.2013.301632

(17) O'Toole TP, Buckel L, Bougault C, et al. Applying the chronic care model to homeless Veterans: effect of a population approach to primary care on utilization and clinical outcomes. *Am J Public Health.* 2010;100(12):2493–9.

(18) Jones AL, et al. Providing positive primary care experiences for homeless Veterans through tailored medical homes: the Veterans health administration's homeless patient aligned care teams. *Med Care*. 2019;57(4):270–8.

(19) Jones, AL, Hausmann, LRM, Kertesz, SG, et al. (2018). Differences in experiences with care between homeless and nonhomeless patients in Veterans affairs facilities with tailored and nontailored primary care teams. *Medical Care*, 56, 610–618.

(20) Chrystal, JG., Glover, DL, Young, AS, et al. (2015). Experience of primary care among homeless individuals with mental health conditions. *PLoS One*, 10(2), e0117395

(21) Kertesz SG, Holt CL, Steward JL, et al. Comparing homeless persons' care experiences in tailored versus nontailored primary care programs. *Am J Public Health*, 103 (suppl 2) (2013), pp. S331-S339

(22) O'Toole TP, Johnson EE, Borgia M, et al. Population-tailored care for homeless Veterans and acute care use, cost, and satisfaction: A prospective quasi-experimental trial. *Preventing chronic disease*. 2018;15:E23.

(23) Gundlapalli AV, Redd A, Bolton D, et al. Patient-aligned care team engagement to connect Veterans experiencing homelessness with appropriate health care. *Med Care*. 2017;55:S104-10.

(24) Jones AL, Thomas R, Hedayati DO, et al. Patient predictors and utilization of health services within a medical home for homeless persons. Substance abuse.

2018;39(3):354-360. PMID: 29412071.

(25) Trivedi AN , Jiang, L, Johnson EE, et al. Dual Use and Hospital Admissions Among Veterans Enrolled in the VA's Homeless Patient Aligned Care Team. Health Services

Research 2018; 53 (6 Pt 2): 5219- 37.

(26) Patel, BI, Complex Care Management to Decrease Emergency Department Utilization: A Case Study of the Homeless Patient Aligned Care Team Demonstration Project at VA Greater Los Angeles Healthcare—Escholarship. UCLA, 2013. Available

online: <https://escholarship.org/uc/item/1681c184>

(27) Pauly JB, Moore TA, Shishko I. Integrating a mental health clinical pharmacy specialist into the Homeless Patient Aligned Care Teams. Ment Health Clin

[Internet]. 2018; 8 (4): 169- 74.

(28) Yoon J, Lo J, Gehlert E, et al.: Homeless Veterans' use of peer mentors and effects on costs and utilization in VA clinics. Psychiatric Services 2017; 68:628–631.

(29) Gabrielian S., Chen JC, Minhaj BP, et al. Feasibility and acceptability of a co-located homeless-tailored primary care clinic and emergency department, J Prim Care Commun Health, 8 (4) (2017), pp. 338-344

(30) O'Toole TP, Roberts CB, Johnson EE. Screening for food insecurity in six Veterans administration clinics for the homeless, June-December 2015. *Prev Chronic Dis.* 2017;14:4.

(31) Sestito SF, Rodriguez KL, Saba SK, et al. Homeless Veterans' experiences with substance use, recovery, and treatment through photo elicitation. *Subst Abus.* 2017;38(4):422–31.

(32) Simmons MM, Gabrielian S, Byrne T, et al. A Hybrid III stepped wedge cluster randomized trial testing an implementation strategy to facilitate the use of an evidence-based practice in VA Homeless Primary Care Treatment Programs. *Implement Sci.* 2017;12:46.