Introduction

The NZED COVID-19 Preparedness Survey focuses on your safety and well-being during the COVID-19 pandemic.

The questions are peer-reviewed and created with the help of emergency specialists and infectious disease experts focused on health care worker safety and well-being.

We hope the results from this research will enable dialogue as to best infection control measures, and assist with standardising local protocols with the aim of minimising risk of nosocomial COVID-19 infections in the Emergency Department (ED).

ASMS understands that your commitment to the welfare of your patients and colleagues is predicated on your ability to focus on providing the best medical care in the safest possible work environment. We understand this requires practice and preparation.

Please take this opportunity to relate the current and proposed practices in your ED and share your opinions and thoughts.


If you have any questions regarding this research, please do not hesitate to contact Dr Charlotte Chambers at ASMS: CC@asms.nz

Thank you for your time.

	here is your primary place of work?
2. D e	o you have any negative pressure <u>beds</u> in your ED and if so, how many?
	he purposes of this survey, a negative pressure or negative flow bed is defined as any bed in singl
multi	ple rooms with minimum of 6 air changes per hour with or without an ante room.
	We don't have any negative pressure beds in our ED
	1-4
	5-9
	10-14
	15-19
	20 or more
	Other (please specify)
,	
3. H	ow many beds in your ED at the time of this survey are in shared rooms for cohorted patien
For t	he purposes of this survey shared rooms are defined as large rooms with multiple curtained beds
	1-4
	5-9
	10-14
	15 or more
	I'm not sure
	None
	Other (places appoint)
0	Other (please specify)
0	Other (piease specify)

No						
Ont applicable						
I'm not sure						
Other (please	specify)					
I will be treating	a segregated ED g COVID patients only g Non-COVID patients alternating between the	s only	d			
Comment:	to most minimum	n nhysical	distancing rogu	iromonto in tl	ao non clinio	al staff areas
Do you feel able your ED?			Neither agree		Strongly	
Do you feel able	to meet minimur Strongly agree	m physical Agree		irements in the		
Do you feel able your ED? Offices Work stations on ED			Neither agree		Strongly	
Do you feel able your ED? Offices Vork stations on ED			Neither agree		Strongly	
Do you feel able your ED?			Neither agree		Strongly	
Do you feel able your ED? Offices Vork stations on ED oor Break rooms			Neither agree		Strongly	
Do you feel able your ED? Offices Work stations on ED door Break rooms Bathrooms			Neither agree		Strongly	
Do you feel able your ED? Offices Vork stations on ED loor Break rooms Bathrooms Changing rooms			Neither agree		Strongly	
Do you feel able your ED? Offices Vork stations on ED loor Break rooms Bathrooms Changing rooms Canteens/Cafeteria			Neither agree		Strongly	
Do you feel able your ED? Offices Vork stations on ED loor Break rooms Bathrooms Changing rooms Canteens/Cafeteria			Neither agree		Strongly	Not applicable

1. What type of training have	you had in the	use of PPE a	at your curren	t place of emp	loyment?
Please select all that apply from	m the following:				
None - I have not completed ar	· ·	employer required	d/directed training		
Self-taught using online training	g (video, reading ma	nterial)			
In-person group demonstration	in which I only wate	ched			
In-person group session in whi	ch I practiced puttino	g PPE on and rem	noving it properly		
In-person individual demonstra	tion in which I only v	vatched			
In-person individual session in	which I was observe	ed putting PPE on	and removing it p	properly	
Other (please specify)					
. Approximately, how frequent	ly are you train	ed in any of t	he following a	activities?	
	Upon request	Never	Once	Annually	I'm not sure
		/ \		()	()
Training for donning and doffing PPE	0				
Training for donning and doffing PPE Simulation sessions on intubating COVID patients	0	0	0	0	0
Simulation sessions on intubating	0	0	0	0	0
Simulation sessions on intubating COVID patients Simulation session for non-invasive	0	0	0	0	0
Simulation sessions on intubating COVID patients Simulation session for non-invasive ventilation (NIV: CPAP, HFNC) Simulation sessions on awake/self			0	0	0
Simulation sessions on intubating COVID patients Simulation session for non-invasive ventilation (NIV: CPAP, HFNC) Simulation sessions on awake/self prone positioning of COVID patients Simulation sessions on transporting				0	
Simulation sessions on intubating COVID patients Simulation session for non-invasive ventilation (NIV: CPAP, HFNC) Simulation sessions on awake/self prone positioning of COVID patients Simulation sessions on transporting COVID patients					
Simulation sessions on intubating COVID patients Simulation session for non-invasive ventilation (NIV: CPAP, HFNC) Simulation sessions on awake/self prone positioning of COVID patients Simulation sessions on transporting COVID patients ther (please specify)					
Simulation sessions on intubating COVID patients Simulation session for non-invasive ventilation (NIV: CPAP, HFNC) Simulation sessions on awake/self prone positioning of COVID patients Simulation sessions on transporting COVID patients ther (please specify)	vith a N-95 mask	/respirator with	nin the last 12	months, what m	nethod was use
Simulation sessions on intubating COVID patients Simulation session for non-invasive ventilation (NIV: CPAP, HFNC) Simulation sessions on awake/self prone positioning of COVID patients Simulation sessions on transporting COVID patients ther (please specify)		•	nin the last 12	months, what m	nethod was use
Simulation sessions on intubating COVID patients Simulation session for non-invasive ventilation (NIV: CPAP, HFNC) Simulation sessions on awake/self prone positioning of COVID patients Simulation sessions on transporting COVID patients ther (please specify) 3. If you have been fit tested we determine fit?	n fit tested within the	•	nin the last 12	months, what m	nethod was use
Simulation sessions on intubating COVID patients Simulation session for non-invasive ventilation (NIV: CPAP, HFNC) Simulation sessions on awake/self prone positioning of COVID patients Simulation sessions on transporting COVID patients ther (please specify) 3. If you have been fit tested we determine fit? Not applicable; I have not been	n fit tested within the	•	nin the last 12	months, what m	nethod was use
Simulation sessions on intubating COVID patients Simulation session for non-invasive ventilation (NIV: CPAP, HFNC) Simulation sessions on awake/self prone positioning of COVID patients Simulation sessions on transporting COVID patients ther (please specify) 3. If you have been fit tested we determine fit? Not applicable; I have not been Qualitative (odour or taste determine)	n fit tested within the	•	onin the last 12	months, what m	nethod was use



PE Scenarios						
. What personal pr	ecautions are y	ou currently	using in the f	ollowing settir	ngs?	
Please select all that	apply from the f	following:				
	When in your ED but not providing patient care (charting, making telephone calls)	When in your Break/Tea Room (eating, conversing with colleagues)	When in your ED providing care for a non- COVID-19 low index of suspicion (LIS) patient	When in your ED providing care for COVID-19	When within 2m of an aerosol- generating procedure for a confirmed or HIS COVID-19 case	When using the bathroom facilities
Standard precautions (handwashing, distancing from others)						
Face shield						
Safety glasses/goggles						
Surgical masks						
Reusable fabric masks						
N-95 masks/respirators						
Elastomeric respirators						
Powered air-purifying respirator systems (PAPR CAPR)						
Disposable surgical hat						
Reusable surgical hat						
Standard disposable isolation gown						
Full-body impermeable suit						
Gloves						
Double gloves						
Foot coverings						
other (please specify):						


as we have show	2. Does	your ED have a formal policy on	showering/s	shampooing a	after caring f	or COVID-19	patients?
episode Only if PPE was breached At the end of the shift After reaching home Omment: 3. In your ED, is PPE donning monitored by an observer prior to care for COVID-19 Yes, all the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure Other (please specify) 4. In your ED, is PPE doffing (removal) monitored by an observer to identify breaks technique after care for COVID-19 patients? Yes, all of the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure			Mandatory	Not mandatory	as we have	We don't have shower facilities	Not aware of a formal policy
At the end of the shift After reaching home 3. In your ED, is PPE donning monitored by an observer prior to care for COVID-19 Yes, all the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure Other (please specify) 4. In your ED, is PPE doffing (removal) monitored by an observer to identify breaks technique after care for COVID-19 patients? Yes, all of the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure		tely after every single patient-contact	\circ		\circ	\circ	
After reaching home 3. In your ED, is PPE donning monitored by an observer prior to care for COVID-19 Yes, all the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure Other (please specify) 4. In your ED, is PPE doffing (removal) monitored by an observer to identify breaks technique after care for COVID-19 patients? Yes, all of the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure	Only if P	PE was breached		\bigcirc		\bigcirc	\bigcirc
3. In your ED, is PPE donning monitored by an observer prior to care for COVID-19 Yes, all the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure Other (please specify) 4. In your ED, is PPE doffing (removal) monitored by an observer to identify breaks technique after care for COVID-19 patients? Yes, all of the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure	At the en	nd of the shift					
3. In your ED, is PPE donning monitored by an observer prior to care for COVID-19 Yes, all the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure Other (please specify) 4. In your ED, is PPE doffing (removal) monitored by an observer to identify breaks technique after care for COVID-19 patients? Yes, all of the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure	After rea	ching home		\bigcirc		\bigcirc	\bigcirc
Yes, all the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure Other (please specify) 4. In your ED, is PPE doffing (removal) monitored by an observer to identify breaks technique after care for COVID-19 patients? Yes, all of the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure	Comment:						
Yes, all the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure Other (please specify) 4. In your ED, is PPE doffing (removal) monitored by an observer to identify breaks technique after care for COVID-19 patients? Yes, all of the time (mandatory) Yes, some of the time (ad-hoc) No I'm not sure							
☐ I'm not sure	4. In techn	No I'm not sure Other (please specify) your ED, is PPE doffing (removanique after care for COVID-19 pates) Yes, all of the time (mandatory)		by an observ	ver to identify	v breaks in d	offing
\odot		No					
Other (please specify)							
	Г	Other (please specify)					

loose coloct all that apply from the following	na	
lease select all that apply from the follow	Low Index of Suspicion for COVID-19	High Index of Suspicion/Confirmed
Negative pressure bed (bed with minimum of 6 air changes per hour)		
Video laryngoscopy		
Intubation barrier protection (e.g., intubating boxes intubating bags, etc.)	,	
Intubation barrier protection with integrated suction (e.g., intubating boxes connected to suction; negative pressure hood)		
Intubation response teams (with dedicated staff)		
Intubation through a supraglottic device (e.g., intubating LMA, etc.)		
None of these		
ther (please specify)		
2. Does your ED have a dedicated into Yes, consisting of ICU/Anaesthesia responsible Yes, consisting of ICU/ED formalised Yes, ED only No I'm not sure		uspected COVID-19 patients?

will be treated with non-invasive positive pressure ventilation (NIV, including CPAP or BiPAP), if needed? Yes, in any area in the ED Yes, only in a negative flow/pressure room in ED Yes, only with in-line or expiration viral filter in negative flow/pressure room in ED Yes, only after transfer to the ICU if appropriate No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (includ CPAP or BiPAP) be used in your ED?		Yes
We don't have a formal protocol Comment: 4. For ED patients with confirmed or suspected COVID-19, is your ED practice/protocol that patievill be treated with non-invasive positive pressure ventilation (NIV, including CPAP or BiPAP), if needed? Yes, in any area in the ED Yes, only in a negative flow/pressure room in ED Yes, only with in-line or expiration viral filter in negative flow/pressure room in ED Yes, only after transfer to the ICU if appropriate No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (included CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not intubate" or a "Do Not Resuscitate" order		No
4. For ED patients with confirmed or suspected COVID-19, is your ED practice/protocol that patients will be treated with non-invasive positive pressure ventilation (NIV, including CPAP or BiPAP), if needed? Yes, in any area in the ED Yes, only in a negative flow/pressure room in ED Yes, only with in-line or expiration viral filter in negative flow/pressure room in ED Yes, only after transfer to the ICU if appropriate No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (included CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BIPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order		I'm not sure
4. For ED patients with confirmed or suspected COVID-19, is your ED practice/protocol that patievill be treated with non-invasive positive pressure ventilation (NIV, including CPAP or BiPAP), if needed? Yes, in any area in the ED Yes, only in a negative flow/pressure room in ED Yes, only with in-line or expiration viral filter in negative flow/pressure room in ED Yes, only after transfer to the ICU if appropriate No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (included CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order		We don't have a formal protocol
will be treated with non-invasive positive pressure ventilation (NIV, including CPAP or BiPAP), if needed? Yes, in any area in the ED Yes, only in a negative flow/pressure room in ED Yes, only with in-line or expiration viral filter in negative flow/pressure room in ED Yes, only after transfer to the ICU if appropriate No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (included CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		Comment:
will be treated with non-invasive positive pressure ventilation (NIV, including CPAP or BiPAP), if needed? Yes, in any area in the ED Yes, only in a negative flow/pressure room in ED Yes, only with in-line or expiration viral filter in negative flow/pressure room in ED Yes, only after transfer to the ICU if appropriate No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (included CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		
Yes, only in a negative flow/pressure room in ED Yes, only with in-line or expiration viral filter in negative flow/pressure room in ED Yes, only after transfer to the ICU if appropriate No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (includ CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		
Yes, in any area in the ED Yes, only in a negative flow/pressure room in ED Yes, only with in-line or expiration viral filter in negative flow/pressure room in ED Yes, only after transfer to the ICU if appropriate No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (includ CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		
Yes, only in a negative flow/pressure room in ED Yes, only with in-line or expiration viral filter in negative flow/pressure room in ED Yes, only after transfer to the ICU if appropriate No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (includ CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce	need	ded?
Yes, only with in-line or expiration viral filter in negative flow/pressure room in ED Yes, only after transfer to the ICU if appropriate No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (include CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce	\bigcirc	Yes, in any area in the ED
Yes, only after transfer to the ICU if appropriate No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (included CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce	\bigcirc	Yes, only in a negative flow/pressure room in ED
No We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (includ CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		Yes, only with in-line or expiration viral filter in negative flow/pressure room in ED
We don't have a formal protocol Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (included CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		Yes, only after transfer to the ICU if appropriate
Other (please specify) 5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (included CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		No
5. For confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (included CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		We don't have a formal protocol
CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		Other (please specify)
CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		
CPAP or BiPAP) be used in your ED? Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		
Please select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		
Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce	5. F c	or confirmed or suspect COVID-19 ED patients, under what circumstances might NIV (includ
Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce		
Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce	CPA	P or BiPAP) be used in your ED?
Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce	CPA	P or BiPAP) be used in your ED? se select from the following:
Only when mechanical ventilators are scarce	CPA	P or BiPAP) be used in your ED? se select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP)
	CPA	P or BiPAP) be used in your ED? se select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA)
	CPA	P or BiPAP) be used in your ED? se select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order
	CPA	P or BiPAP) be used in your ED? se select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce
	CPA	P or BiPAP) be used in your ED? se select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce
	CPA	P or BiPAP) be used in your ED? se select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce
	CPA	P or BiPAP) be used in your ED? se select from the following: Any patient with respiratory failure that I think will benefit from NIV if indicated (NIV: CPAP/BiPAP) Only patients that have other co-morbidities known to benefit from NIV (eg. COPD, CHF, OSA) Only patients who have a "Do Not Intubate" or a "Do Not Resuscitate" order Only when mechanical ventilators are scarce

Department	Breach	Not a breach	Not aware of formal polic
Inadequate face protection			
Inadequate eye protection			
Improper donning/doffing			\circ
Exposure of skin due to a glove or gown tear	\circ	\circ	\circ
Inadequate shoe cover	\circ		
Direct contact of skin to any secretion	\bigcirc	\bigcirc	\bigcirc
Needle stick			
INCCUIC SUCK			
Poor mask fit Other (please specify) 2. What measures have been advised by			Not aware of a formal
Poor mask fit Other (please specify) 2. What measures have been advised by dentified?	your hospital/ED ad	Iministration when	
Poor mask fit Other (please specify) 2. What measures have been advised by dentified? Shower immediately			Not aware of a formal
Poor mask fit Other (please specify) 2. What measures have been advised by dentified? Shower immediately Report to ID/designated authorities			Not aware of a formal
Poor mask fit Other (please specify) 2. What measures have been advised by dentified? Shower immediately Report to ID/designated authorities Retraining given for donning or doffing			Not aware of a formal
Poor mask fit Other (please specify) 2. What measures have been advised by dentified? Shower immediately Report to ID/designated authorities			Not aware of a formal
Poor mask fit Other (please specify) 2. What measures have been advised by dentified? Shower immediately Report to ID/designated authorities Retraining given for donning or doffing			Not aware of a formal
Poor mask fit Other (please specify) 2. What measures have been advised by dentified? Shower immediately Report to ID/designated authorities Retraining given for donning or doffing Quarantine with testing protocol			Not aware of a formal
Poor mask fit Other (please specify) 2. What measures have been advised by dentified? Shower immediately Report to ID/designated authorities Retraining given for donning or doffing Quarantine with testing protocol			Not aware of a formal
Poor mask fit Other (please specify) 2. What measures have been advised by dentified? Shower immediately Report to ID/designated authorities Retraining given for donning or doffing Quarantine with testing protocol			Not aware of a formal

PPE supply and re-use			
1. At the time of this survey, please select unavailable for clinical use in your ED:	t from the followir	ng any PPE that is	out of stock or other
Reusable face shields			
Disposable face-shields (single use)			
Safety glasses/goggles			
Surgical masks			
Reusable fabric masks			
N-95 masks/respirators			
Elastomeric respirators			
Powered air- purifying respirator systems (PAPR	R, CAPR, etc.)		
Disposable surgical hat			
Reusable surgical hat			
Standard disposable isolation gown			
Full-body impermeable suit			
Gloves			
Foot coverings			
Other (please specify)			
. Have you ever re-used PPE equipment acc	cording to any of	the following scena	arios?
	Yes	No	I'm not sure
We re-use N-95 masks without sterilization			
We re-use N-95 masks after sterilization			
We re-use face shields without washing with cleaning solution	0	0	0
We re-use face shields after washing with cleaning solution provided			
	0	0	0
solution provided We re-use goggles without washing with cleaning	0	0	0



I am/was confident that our PPE protocol will keep	uring Level 4 Loc	Kaown	At the t	ime of this	survey
me completely safe					
I think my ED's protocol put me at risk and that I should have better PPE than is available, or use PPE more often than required by protocol					
I think my ED's PPE protocol is too restrictive, and I feel that I can safely practice without wearing PPE every time that it is required by protocol					
I am/was unsure about the safety of our PPE protocol and feel neither safe or unsafe					
ther (please specify) . At the time of the initial COVID-19 outbreak in	n New Zealand	l (during l	evel 4 lockde	own), ple	ase
I am/was not aware of a PPE protocol in my ED other (please specify) . At the time of the initial COVID-19 outbreak in onsider how you felt about the following: I worried that family members or other close contacts were at	Never	I (during le	evel 4 lockdo	own), ple	ase Always
ther (please specify) . At the time of the initial COVID-19 outbreak in onsider how you felt about the following: I worried that family members or other close contacts were at of exposure to COVID-19 because of my work	Never risk				
ther (please specify) . At the time of the initial COVID-19 outbreak in onsider how you felt about the following: I worried that family members or other close contacts were at	Never risk				
Ther (please specify) At the time of the initial COVID-19 outbreak in onsider how you felt about the following: I worried that family members or other close contacts were at of exposure to COVID-19 because of my work I worried that I was at risk of exposure to COVID-19 because	Never risk of				
ther (please specify) . At the time of the initial COVID-19 outbreak in onsider how you felt about the following: I worried that family members or other close contacts were at of exposure to COVID-19 because of my work I worried that I was at risk of exposure to COVID-19 because my work	Never risk of of o				
ther (please specify) At the time of the initial COVID-19 outbreak in onsider how you felt about the following: I worried that family members or other close contacts were at of exposure to COVID-19 because of my work I worried that I was at risk of exposure to COVID-19 because my work I worried about the level of preparedness of my hospital and E I worried about the supply of adequate and appropriate PPE i	Never risk of of o				
Ther (please specify) At the time of the initial COVID-19 outbreak in onsider how you felt about the following: I worried that family members or other close contacts were at of exposure to COVID-19 because of my work I worried that I was at risk of exposure to COVID-19 because my work I worried about the level of preparedness of my hospital and E worried about the supply of adequate and appropriate PPE i ED	Never risk of of o				
ther (please specify) At the time of the initial COVID-19 outbreak in onsider how you felt about the following: I worried that family members or other close contacts were at of exposure to COVID-19 because of my work I worried that I was at risk of exposure to COVID-19 because my work I worried about the level of preparedness of my hospital and E worried about the supply of adequate and appropriate PPE in ED I felt anxious and stressed because of COVID-19	Never risk of of o				

B. Please answer the follow	ing questions rega	arding your v	icws on now prep	-	
there was another wave of (COVID-19 in New 2	Zealand			
	Strongly agree	Agree	Neither agree nor disagree	Disagree	Strongly disagree
Our ED is prepared and ready for another wave of COVID-19					
Our ED would have adequate staff levels if there was another wave of COVID-19					
Our ED would have adequate supporting of appropriate PPE if there was another wave of COVID-19	olies				
We would be able to rapidly test ar a timely manner diagnose possible cases of COVID-19					
Other (please specify)					
•	Ν	leither agree nor			
4. Do you feel you are at ris Strongly agree	Agree	leither agree nor disagree	Disagree	5	Strongly disagree
				ξ	Strongly disagree
				\$	Strongly disagree
Strongly agree				\$	Strongly disagree
Strongly agree				Ç	Strongly disagree
Strongly agree				Ş	Strongly disagree
Strongly agree				S	Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree
Strongly agree					Strongly disagree

thir	u feel you are at risk of COVID-19 infection AT WO nk would put you at risk of exposure to COVID-19	
ase :	select 1 for the most likely reason through to 8 as	the least likely.
ot app	plicable- I do not fear risk of COVID-19 exposure at work	
ot bei	ing able to access adequate PPE	
oarin	ng inadequate PPE for patient(s) not suspected of COVID-19 infe	rtion
Calli	ig madequate FFE for patient(s) not suspected of COVID-19 lines	AUGUT
earin	ng inadequate PPE for patient(s) suspected of COVID-19 infection	
ontrad	acting it from a fellow staff member in the ED	
\neg		
ccide	ental PPE doffing exposure	
eane	ers have been provided inadequate training and/or inadequate PP	E
	what a most fit tooting for staff	
adeqi	quate mask fit testing for staff	
∕e w	vould be grateful to hear your thoughts on any other a	spects regarding the level of preparedness o
r ED	or the impact of COVID-19	