

# Supplementary Information

## Improved screening of COVID-19 cases through a Bayesian network symptoms model and psychophysical olfactory test

Susana Eyheramendy<sup>a,b,\*</sup>, Pedro A. Saa<sup>c,\*</sup>, Eduardo A. Undurraga<sup>d,e,f</sup>, Carlos Valencia<sup>g</sup>, Carolina López<sup>g</sup>, Luis Méndez<sup>h,i</sup>, Javier Pizarro-Berdichevsky<sup>j,k</sup>, Andrés Finkelstein-Kulka<sup>l,m</sup>, Sandra Solari<sup>n</sup>, Nicolás Salas<sup>b</sup>, Pedro Bahamondes<sup>b</sup>, Martín Ugarte<sup>b</sup>, Pablo Barceló<sup>b,o</sup>, Marcelo Arenas<sup>b,p,\*\*</sup>, and Eduardo Agosin<sup>c,g,\*\*</sup>

<sup>a</sup>Faculty of Engineering and Science, Universidad Adolfo Ibáñez, Chile

<sup>b</sup>Millennium Institute for Foundational Research on Data (IMFD), Chile

<sup>c</sup>Department of Chemical and Bioprocess Engineering, School of Engineering, Pontificia Universidad Católica de Chile, Chile

<sup>d</sup>School of Government, Pontificia Universidad Católica de Chile, Chile

<sup>e</sup>Millennium Initiative for Collaborative Research in Bacterial Resistance (MICROB-R), Chile

<sup>f</sup>Research Center for Integrated Disaster Risk Management (CIGIDEN), Chile

<sup>g</sup>Center for Aromas and Flavors, DICTUC SA., Chile

<sup>h</sup>Endoscopy Unit, Hospital Padre Hurtado, Chile

<sup>i</sup>Department of Gastroenterology, Clinica Alemana de Santiago, Chile

<sup>j</sup>Center for Innovation in Pelvic Floor, Hospital Sótero del Río, Chile

<sup>k</sup>Department of Obstetrics and Gynecology, Faculty of Medicine, Pontificia Universidad Católica de Chile, Chile

<sup>l</sup>Department of Otolaryngology, Clinica Alemana de Santiago, Chile

<sup>m</sup>Faculty of Medicine, Clínica Alemana Universidad del Desarrollo, Chile

<sup>n</sup>Department of Clinical Laboratory, Faculty of Medicine, Pontificia Universidad Católica de Chile, Chile

<sup>o</sup>Institute for Mathematical and Computational Engineering, Pontificia Universidad Católica de Chile, Chile

<sup>p</sup>Department of Computer Science, School of Engineering, Pontificia Universidad Católica de Chile, Chile

\*S.E. and P.A.S. contributed equally to this work

\*\*To whom correspondence should be addressed. E-mail: marenas@ing.puc.cl; agosin@ing.puc.cl

This manuscript was compiled on April 16, 2021

# Supplementary Text

## S1. KOR test protocol

To carry out an olfactory test, an evaluator must first enter the information of the individual to undergo the olfactory test. All this information is requested and stored in the KOR platform:

- Personal information: national identification number (or passport number), name, date of birth, nationality and gender.
- Information about previous SARS-CoV-2 test: the individual is asked whether he/she has undergone the SARS-CoV-2 RT-PCR test, and if the answer is yes, then he/she is asked the date of the test and whether the result was either negative or positive or still pending.
- Information on comorbidities: the individual is asked about his/her comorbidities. There is a predefined list that the evaluator can refer to, which includes: loss of smell, respiratory allergies, hypertension, rhinitis or sinusitis, Parkinson's disease, Alzheimer's disease, and diabetes. In addition, the platform provides an open question where the evaluator can enter any additional comorbidities.
- Information on smoking: the individual must indicate if he/she smokes.

This information is only requested once; the next time the individual undergoes the olfactory test, he/she only needs to provide the identification number in the platform for his/her information to be displayed.

As a second step, the evaluator must ask about the individual's symptoms and discomforts that may be associated with SARS-CoV-2. In particular, there is a check list that the evaluator has to use, which includes cough, cold or flu, fatigue, headache, and loss of appetite. If any symptom or discomfort is entered, then the number of days with this condition is asked. This data is requested every time an individual undergoes the olfactory test.

Next, the evaluator proceeds to ask six questions to measure partial or total loss of smell in the individual being tested. For each of these questions, the platform first tells the evaluator which odor (scent) to present to the individual, and then shows that person four pictures of odors asking him/her to identify the one he/she is smelling (see next section), and also giving a fifth alternative to indicate that he/she cannot identify it. It is important to notice here that the platform chooses at random the order in which the odors are presented, and also the distractors that will be used for each question.

Finally, according to the responses of the individual being tested, the platform produces a result in real time, which indicates if the individual has a reduction in his/her olfactory capacity, and therefore is a SARS-CoV-2 suspect.

### Scents labels

Each scent of the KOR test is coded by a letter from "A" to "F", as follows: (A) Banana; (B) Caramel; (C) Mint; (D) Orange; (E) Pineapple; (F) Vanilla. Below, the available options for each multiple-choice odor recognition question are shown.

A: Banana



1 Flowers



2 Vanilla



3 Garlic



4 Banana

B: Caramel



1 Butter



2 Grass



3 Caramel



4 Banana

C: Mint



1 Mint



2 Tangerine



3 Chili pepper



4 Caramel

D: Orange



1 Watermelon



2 Oregano



3 Orange



4 Mint

E: Pineapple



1 Onion



2 Pineapple



3 Lavander



4 Banana

F: Vanilla



1 Pepper



2 Honey



3 Vanilla



4 Orange

## Supplementary Tables

Table 1: Proportion of presence of symptoms and variables among PCR positive and PCR negative individuals.

Variables	PCR negative	PCR positive
Cough	0.27	0.51
Fever	0.10	0.31
Muscular pain	0.32	0.48
Breathing difficulty	0.12	0.18
I1	0.19	0.53
I4	0.06	0.23
sDias	0.44	0.68
Anosmia	0.02	0.16
Ageusia	0.00	0.08
I7	0.16	0.07
I8	0.00	0.09
Gender	0.51	0.53

Table 2: KOR test odor recognition statistics in the pilot study sample by different criteria

Criteria	Score							Total
Genre	0	1	2	3	4	5	6	
Male	4	15	23	58	169	520	1009	1798
Female	0	1	4	17	44	146	279	491
Country of origin								
Chile	4	15	19	52	158	501	1083	1832
Haití	0	1	6	4	13	25	23	72
Perú	0	0	0	6	7	23	25	61
Venezuela	0	0	2	7	26	81	123	239
Colombia	0	0	0	4	5	14	17	40
Other	0	0	0	2	4	22	17	45
Age range								
16-20	1	0	0	0	5	12	24	42
21-25	2	1	4	12	29	84	194	326
26-30	0	6	8	16	38	143	286	497
31-35	1	3	3	11	29	110	239	396
36-40	0	0	5	13	22	101	192	333
41-45	0	1	1	8	24	65	115	214
46-50	0	2	3	6	22	66	117	216
51-55	0	2	2	1	20	45	63	133
56-60	0	1	1	4	15	22	31	74
61-65	0	0	0	4	7	7	8	26
66-70	0	0	0	0	0	3	0	3

Table 3: Chemical composition of KOR test scents

Scent	Provider code*	Solvent	Main odorant compounds	Concentration (g/L)
Banana	PTMEZ382401	Propylene glycol	Isoamyl acetate	11.49
			Vanillin	4.57
			2,3-Dihydroxypropyl acetate	3.85
			Phenethyl isovalerate	3.50
			Isoamyl butanoate	3.00
Caramel	PTMEZ382701	Propylene glycol	Vanillin	7.20
			Ethyl maltol	6.08
			Nonanoic acid	2.45
			Furaneol	1.57
			Octanoic acid	0.60
Mint	PTMEZ381801	Propylene glycol	Menthyl acetate	14.40
			p-Menthan-3-one	3.80
			D-Menthol	2.10
			D-Carvone	2.00
Orange	PTMEZ381901	Tributyl acetylcitrate	D-Limonene	2.60
			Linalool	1.70
			8-p-Menthadien-1,2-diol	0.45
			Decanal	0.31
Pineapple	PTMEZ382101	Propylene glycol	Allyl hexanoate	7.05
			Isoamyl butanoate	2.99
			Ethyl hexanoate	2.97
			Ethyl octanoate	1.40
Vanilla	PTMEZ383177	Propylene glycol	Ethyl isovanilline	20.66
			Benzyl alcohol	4.67
			Piperonal	3.90
			Vanillin	0.63

\* Alpha Group, Santiago, Chile.

## Supplementary Figures




Figure 1: Weights contributions of the recognized odors to the risk of a positive PCR result for COVID-19. Red dots represent the weights when the individual fail to recognize an odor, whereas black dots represent the opposite. X1=Banana, X2=Caramel, X3=Mint, X4=Orange, X5=Pineapple, X6=Vanilla




Figure 2: Weights contribution of the symptoms to the risk of a positive PCR result for COVID-19. Red dots represent the weights when there is absence of the symptom or condition, whereas black dots represent the opposite, i.e., presence of the symptom or condition. Cough ( $X_7$ ), fever ( $X_8$ ), muscular pain ( $X_9$ ), breathing difficulty ( $X_{10}$ ), self-reported anosmia ( $X_{14}$ ), ageusia ( $X_{15}$ ), five indicator variables ( $X_{11}, X_{12}, X_{13}, X_{16}, X_{17}$ ) and gender  $X_{18}$






Figure 3: Recognition level of 7 odors in a pilot study for the KOR test. All odors were highly recognized in sample population with the exception of peach. The latter odor was excluded from the odor panel of the KOR test.





Figure 4: Conditional probabilities for the perception of smells given the flu and RT-PCR status: 0: without flu and without COVID-19; 1:with flu and without COVID-19; 2:without flu and with COVID-19; 3:with flu and with COVID-19.



*KOR test intrinsic characteristics for a score of 4.*

Cutoff $k \leq 4$		
Statistic	Value	95% CI
Sensitivity	53%	48% to 59%
Specificity	81%	78% to 84%
Positive Likelihood Ratio	2.8	2.3 to 3.4
Negative Likelihood Ratio	58%	0.5 to 0.7
Disease Prevalence	33%	
Positive Predictive Value	58%	54% to 63%
Negative Predictive Value	78%	75% to 80%
Accuracy	72%	

Figure 5: COVID-19 prediction performance of the olfactory test assuming equal weights for the odors. The best classification performance is obtained with a score of four or less correctly identified odors (COVID-19 positive). The UC-Christus dataset was employed to evaluate the performance of this model.