

EFFECT OF LINEAR GROWTH RATE AND CHANGE IN BODY MASS INDEX IN CHILDHOOD AND ADOLESCENCE ON BLOOD PRESSURE IN AFRO-CARIBBEAN YOUTH: THE VULNERABLE WINDOWS COHORT STUDY

Trevor S. Ferguson¹, Tamika Y. Royal-Thomas², Lisa Chin-Harty³, Minerva M. Thame⁴,
Terrence E. Forrester⁵, Clive Osmond⁶, Michael S. Boyne³, Rainford. J. Wilks¹

¹*Caribbean Institute for Health Research, The University of the West Indies, Mona, Kingston 7, Jamaica*

²*Department of Mathematics, Faculty of Science and Technology, The University of the West Indies, Mona, Kingston 7, Jamaica*

³*Department of Medicine, Faculty of Medical Sciences, The University of the West Indies, Mona, Kingston 7, Jamaica*

⁴*Department of Child and Adolescent Health, Faculty of Medical Sciences, The University of the West Indies, Mona, Kingston 7, Jamaica*

⁵*UWI Solutions for Developing Countries, The University of the West Indies, Mona, Kingston 7, Jamaica*

⁶*Faculty of Medicine, University of Southampton, England*

Supplementary Tables and Figures

Table S1: Correlation coefficients for linear growth and rate of change BMI together at given time periods

Variable	cH 0-6 months	cH 6 months – 2 years	cH 2-8 years	cH 8-15 years	cBMI 0-6 months	cBMI 6 months – 2 years	cBMI 2-8 years	cBMI 8-15 years
cH 0-6 months	1.0							
cH 6 months – 2 years	0.076	1.0						
cH 2-8 years	0.095	0.040	1.0					
cH 8-15 years	0.268 ***	-0.177 *	-0.227 ***	1.0				
cBMI 0-6 months	0.235 ***	0.147 *	-0.003	0.119	1.0			
cBMI 6 months – 2 years	0.058	0.155 **	0.007	-0.193 **	-0.049	1.0		
cBMI 2-8 years	0.025	0.199 **	0.330 ***	-0.229 ***	-0.105	0.144 *	1.0	
cBMI 8-15 years	-0.024	-0.069	0.035	-0.111	-0.022	0.014	-0.018	1.0

Coefficients are derived from Person's correlation. All variables were entered as z-score; cH = change in height; cBMI = change in body mass index.

*p<0.05; **p<0.01; ***p<0.001

Table S2: Number of missing and non-missing values for variables used in the analysis

Variable	Number of missing observations	Number of non-missing observations	% Missing
Birth Weight	28	1879	1.5
Birth length	45	1862	2.4
Birth BMI	45	1862	2.4
Gestational age at birth	55	1852	2.9
Socioeconomic status	1	1906	0.1
Change in BMI 0-6 month	291	1616	15.3
Change in BMI 6 months – 2 years	396	1511	20.8
Change in BMI 2 years – 8 years	492	1415	25.8
Change in BMI 8 years – 15 years	468	1439	24.5
Linear growth rate 0-6 months	291	1616	15.3
Linear growth rate 6 months – 2 years	396	1511	20.8
Linear growth rate 2 years – 8 years	492	1415	25.8
Linear growth rate 8 years – 15 years	465	1442	24.4

Analyses were limited to participants with available blood pressure measurements after age 15 years. Estimates based on 1907 estimates used for diastolic BP models. Systolic blood pressure used 1909 observations. There were no missing values for age or sex among participants included in the analysis.

Table S3: Bivariate models for systolic blood pressure from age 15-21 years as outcome variable

Variable	Regression Coefficient	95% Confidence Interval	P-value
Birth length z-score	0.66	-0.31, 1.64	0.182
Linear growth rate 0-6 month	2.06	1.28, 2.84	<0.001
Linear growth rate 6 months – 2 years	0.72	-0.23, 1.68	0.138
Linear growth rate 2 years – 8 years	0.65	-0.11, 1.42	0.091
Linear growth rate 8 years – 15 years	1.54	0.74, 2.34	<0.001
Age (years)	-1.10	-1.40, -0.80	<0.001
Sex (male vs. female)	7.43	5.68, 9.18	<0.001
Height at age 15	0.08	0.02, 0.14	0.009
BMI at age 15	0.32	0.13, 0.52	0.001
Birth weight	0.90	-0.07, 1.87	0.069
Birth length	0.66	-0.31, 1.64	0.182
Gestational age (days)	-0.04	-0.12, 0.03	0.247
Socioeconomic status at birth	0.09	-0.12, 0.31	0.390

Table S4: Bivariate models for diastolic blood pressure from age 15-21 years as outcome variable

Variable	Regression Coefficient	95% Confidence Interval	P-value
Birth length z-score	0.63	0.01, 1.26	0.047
Linear growth rate 0-6 month	0.54	0.01, 1.06	0.045
Linear growth rate 6 months – 2 years	0.49	-0.08, 1.06	0.092
Linear growth rate 2 – 8 years	0.59	0.02, 1.17	0.044
Linear growth rate 8 – 15 years	-0.19	-0.77, 0.39	0.514
Age (years)	0.72	0.45, 0.99	<0.001
Sex (male vs. female)	0.43	-0.78, 1.65	0.485
Height at age 15	0.01	-0.03, 0.05	0.705
BMI at age 15	0.22	0.09, 0.36	0.001
Birth weight	0.53	-0.07, 1.12	0.082
Birth length	0.63	0.01, 1.26	0.047
Gestational age (days)	-0.01	-0.06, 0.04	0.646
Socioeconomic status at birth	0.04	-0.09, 0.18	0.517

Table S5: Stratified models for postnatal linear growth and systolic and diastolic blood pressure by birth weight category

Variable	Low Birth Weight		Normal Birth Weight	
	Regression Coefficient	P-value	Regression Coefficient	P-value
Linear Growth with SBP				
Birth length z-score	1.00	0.578	0.56	0.323
Linear growth rate 0-6 months	1.52	0.264	1.09	0.015
Linear growth rate 6 months – 2 years	1.09	0.288	0.70	0.116
Linear growth rate 2 years – 8 years	-0.04	0.975	0.58	0.167
Linear growth rate 8 years – 15 years	0.71	0.652	-0.07	0.916
Gestational age at birth (days)	0.02	0.826	-0.11	0.013
Age (years)	-0.42	0.435	-1.13	<0.001
Sex (male vs. female)	7.66	0.020	7.24	<0.001
BMI at age 15 (kg/m ²)	0.48	0.091	0.33	0.001
Linear Growth with DBP				
Birth length z-score	1.24	0.355	1.04	0.009
Linear growth rate 0-6 months	0.76	0.560	0.36	0.261
Linear growth rate 6 months – 2 years	0.05	0.956	0.45	0.150
Linear growth rate 2 years – 8 years	-0.36	0.759	0.57	0.082
Linear growth rate 8 years – 15 years	-0.65	0.704	-0.65	0.213
Gestational age at birth (days)	-0.02	0.694	-0.06	0.060
Age (years)	1.59	0.001	0.66	<0.001
Sex (male vs. female)	1.13	0.689	1.10	0.253
BMI at age 15 (kg/m ²)	0.23	0.346	0.15	0.037

BP = blood pressure; SBP = systolic BP; DBP = diastolic BP; BMI = body mass index

Table S6: Stratified models for rate of change in body mass index on systolic and diastolic blood pressure by birth weight category

Variable	Low Birth Weight		Normal Birth Weight	
	Regression Coefficient	P-value	Regression Coefficient	P-value
Change in BMI and SBP				
BMI at birth z-score	-4.28	0.005	0.42	0.429
Change in BMI 0-6 months	1.11	0.356	0.67	0.148
Change in BMI 6 months – 2 years	2.08	0.009	1.51	<0.001
Change in BMI 2 years – 8 years	1.50	0.314	1.17	0.002
Change in BMI 8 years – 15 years	1.51	0.165	0.66	0.129
Gestational age at birth (days)	0.08	0.161	-0.10	0.028
Socioeconomic status score at birth	0.72	0.003	-0.02	0.869
Age (years)	-0.45	0.404	-1.11	<0.001
Sex (male vs. female)	8.59	0.002	6.84	<0.001
Height at age 15	-0.001	0.977	0.07	0.056
Change in BMI and DBP				
BMI at birth z-score	-2.63	0.013	0.05	0.888
Change in BMI 0-6 months	-0.71	0.334	0.49	0.116
Change in BMI 6 months – 2 years	0.42	0.535	0.65	0.044
Change in BMI 2 years – 8 years	0.33	0.761	0.64	0.025
Change in BMI 8 years – 15 years	2.06	0.022	0.57	0.102
Gestational age at birth (days)	0.02	0.527	-0.04	0.262
Socioeconomic status score at birth	0.32	0.052	0.05	0.517
Age (years)	1.46	<0.001	0.65	<0.001
Sex (male vs. female)	1.53	0.452	0.19	0.784
Height at age 15	-0.02	0.525	0.02	0.489

BP = blood pressure; SBP = systolic BP; DBP = diastolic BP; BMI = body mass index




Figure S1: Annualized mean linear growth rates in age periods during childhood






Figure S2: Mean change in body mass index in age periods during childhood



Panel A: Systolic Blood Pressure (mm Hg)



Panel B: Diastolic Blood Pressure (mm Hg)

Figure S3: Spaghetti plot showing participants' systolic blood pressure (SBP, mm Hg) [Panel A] and diastolic blood pressure (DBP) [Panel B] from age 15-21 years vs age (years) for male and female participants.

Each line represents the change in the SBP for each participant in the study from 15-21 years. The centre red line represents the mean SBP for all participants together over the period.