STROBE Statement—Checklist of items that should be included in reports of *cohort studies*

	Item No	Recommendation	Page No
Title and abstract	1	(<i>a</i>) Indicate the study's design with a commonly used term in the title or the abstract	1 and 2
		(b) Provide in the abstract an informative and balanced summary of what was	2
		done and what was found	
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	3-4
Objectives	3	State specific objectives, including any prespecified hypotheses	4
Methods			
Study design	4	Present key elements of study design early in the paper	4
Setting	5	Describe the setting, locations, and relevant dates, including periods of	4
		recruitment, exposure, follow-up, and data collection	
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of	4
		participants. Describe methods of follow-up	
		(b) For matched studies, give matching criteria and number of exposed and	N/A
		unexposed	4-5
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and	4-3
	0.*	effect modifiers. Give diagnostic criteria, if applicable	4-5
Data sources/	8*	For each variable of interest, give sources of data and details of methods of	4-5
measurement		assessment (measurement). Describe comparability of assessment methods if there is more than one group	
Bias	9	Describe any efforts to address potential sources of bias	4-5
Study size	10	Explain how the study size was arrived at	5
Quantitative variables	11	Explain how due study size was drived at Explain how quantitative variables were handled in the analyses. If applicable,	6
		describe which groupings were chosen and why	
Statistical methods	12	(<i>a</i>) Describe all statistical methods, including those used to control for	6
		confounding	
		(b) Describe any methods used to examine subgroups and interactions	6
		(c) Explain how missing data were addressed	6
		(d) If applicable, explain how loss to follow-up was addressed	6
		(<u>e</u>) Describe any sensitivity analyses	N/A
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially	4,7
		eligible, examined for eligibility, confirmed eligible, included in the study,	
		completing follow-up, and analysed	
		(b) Give reasons for non-participation at each stage	4
		(c) Consider use of a flow diagram	None
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social)	7
		and information on exposures and potential confounders	Table
		(b) Indicate number of participants with missing data for each variable of interest	S2 7
		(c) Summarise follow-up time (eg, average and total amount)	

Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their	7-9
		precision (eg, 95% confidence interval). Make clear which confounders were adjusted for	
		and why they were included	
		(b) Report category boundaries when continuous variables were categorized	Yes
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a	N/A
		meaningful time period	
Other analyses	17	Report other analyses done-eg analyses of subgroups and interactions, and sensitivity	
		analyses	
Discussion			
Key results	18	Summarise key results with reference to study objectives	9-10
Limitations 1	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision.	11-
		Discuss both direction and magnitude of any potential bias	12
Interpretation 2	20	Give a cautious overall interpretation of results considering objectives, limitations,	10-
		multiplicity of analyses, results from similar studies, and other relevant evidence	12
Generalisability	21	Discuss the generalisability (external validity) of the study results	12
Other informati	on		
Funding	22	Give the source of funding and the role of the funders for the present study and, if	13
		applicable, for the original study on which the present article is based	
,			

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at http://www.plosmedicine.org/, Annals of Internal Medicine at http://www.annals.org/, and Epidemiology at http://www.epidem.com/). Information on the STROBE Initiative is available at http://www.strobe-statement.org.