

S2 Table

Information on the function of genes with increased expression in prior asymptomatic versus clinical SARS-CoV-2 infection

Gene	Description and Biological Terms Related (if applicable)
DUSP8	Dual Specificity Phosphatase 8 Pathways: Signaling by GPCR and MAPK signaling pathway Terms: phosphatase activity and protein tyrosine/serine/threonine phosphatase activity
CACNA2D2	Calcium Voltage-Gated Channel Auxiliary Subunit Alpha2delta 2 Pathways: ERK Signaling and Activation of cAMP-Dependent PKA. Terms: voltage-gated calcium channel activity and calcium channel regulator activity
GCAT	Glycine C-Acetyltransferase Pathways: Glycine, serine and threonine metabolism, Viral mRNA Translation. Terms: pyridoxalphosphatebinding and glycine C-acetyltransferaseactivity.
GZMH	GranzymeH Reported to be constitutively expressed in the NK (natural killer) cells of the immune system and may play a role in the cytotoxic arm of the innate immune response by inducing target cell death and by directly cleaving substrates in pathogen-infected cells. [RefSeq, Nov 2015] Pathways are Peptide hormone metabolism and Metabolism of proteins. Terms: serine-type endopeptidase activity.
CLEC12A (MICL)	C-lectine-like receptor Pathways are Innate Immune System and C-type lectin receptor signaling pathway. Terms: includecarbohydratebinding.
CLEC1B (CLEC2)	C-Type Lectin Domain Family 1 Member B Natural killer (NK) cells express multiple calcium-dependent (C-type) lectin-like receptors, such as CD94 and NKG2D, that interact with major histocompatibility complex class I molecules and either inhibit or activate cytotoxicity and cytokine secretion. CLEC2 is a C-type lectin-like receptor expressed in myeloid cells and NK cell (Colonna et al., 2000 [PubMed 10671229]) Pathways: Response to elevated platelet cytosolic Ca ²⁺ and C-type lectin receptor signaling pathway. Terms: transmembrane signaling receptor activity and carbohydrate binding.
BEX5	Brain expressed X-linked5, Nerve Growth Factor Receptor-Associated Protein 2 [Alvarez et al. 2005]
ENC1	Ectodermal-Neural Cortex 1 Pathways are WNT Signaling. Terms: actinbinding.
CYP3A43	Cytochrome P450 Family 3 Subfamily A Member 43 Pathways are Paroxetine Pathway, Pharmacokinetics and Drug metabolism - cytochrome P450. Terms: iron ion binding and oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen