

Figure S1. SNP effects estimated by the joint models (JM-cmp, JM-mis, JM-sep) for each SNP on (a.) HbA1c ($\beta_{g(1)}$), (b.) SBP ($\beta_{g(2)}$), (c.) Time-to-DR ($\gamma_{g(1)}$) and (d.) Time-to-DN ($\gamma_{g(2)}$), assessed using $R=1000$ replicates of $N=667$ DCCT subjects simulated under the causal genetic scenario from Figure 2. For (a.) and (b.), estimates from the multivariate linear mixed models (LMM) are the same for JM-cmp, JM-mis and JM-sep. For (c.) and (d.), we also compare the estimates from the joint models with the estimates from CM-obs and from marginal time-to-event models (MM). For each longitudinal (a., b.) or time-to-event trait (c., d.), causal SNP associations are indicated by red boxes. For SNP3 which has indirect effects on both time-to-T1DC traits via the unmeasured risk factor U , all methods except JM-cmp, exhibit a biased estimate of direct effect, which captures some of the indirect SNP3 effect via U , ignored from the time-to-event sub-models. For SNP5 with direct and indirect effects on time-to-DN induced via SBP, CM-obs and MM overestimate the direct SNP5 effect due to ignoring measurement errors in SBP (CM-obs) or ignoring the indirect effect of SNP5 via SBP (MM).






Figure S2. Quantile-quantile plots of the P -values from the Wald tests for each SNP effect in longitudinal (upper) and time-to-event components (bottom) of the complete joint model (JM-cmp), assessed using $R=1000$ replicates of $N=667$ DCCT subjects simulated under the *global null genetic scenario*. For each trait, SNP association P -values obtained for all $M=5$ SNPs and all replicates are pooled, yielding $R=5000$ replicates per plot.


Figure S3. Classification rates of each SNP as direct and/or indirect association with each time-to-event trait through each longitudinal trait, assessed using $R=1000$ replicates of $N=667$ DCCT subjects simulated under the *global null genetic scenario*. Panels with classification rates for associations presented in Table 3 are indicated by red boxes. For SNP3 which has indirect effects on time-to-T1DC traits induced via U , assumed *unmeasured*, the hypothesis testing procedure detect it as a direct association. As explained in *Materials and Methods*, joint models are fitted separately for each SNP, and the SNP association is tested with each of the longitudinal traits (HbA1c, SBP) and each of the time-to-T1DC traits (time-to-DR, time-to-DN).



SNP3



SNP4



model JM-cmp JM-mis CM-obs

SNP5







Figure S4. Classification rates of each SNP as direct and/or indirect association with each time-to-event trait through each longitudinal trait, assessed using $R=1000$ replicates of $N=667$ DCCT subjects simulated under the *causal* genetic scenario from Figure 2. Panels with underlying causal associations are indicated by red boxes for each SNP; other associations represent misclassified associations. For SNP3 which has indirect effects on both time-to-T1DC traits, induced via U , assumed *unmeasured*, the hypothesis testing procedure detect it as a direct association. As explained in *Materials and Methods*, joint models are fitted separately for each SNP, and the SNP association is tested with each of the longitudinal traits (HbA1c, SBP) and each of the time-to-T1DC traits (time-to-DR, time-to-DN).



SNP3



SNP4



SNP5






Figure S5. Classification of direct and/or indirect SNP association for 307 candidate SNPs tested in $N=516$ DCCT subjects using the contemporaneous association structure and the two alternative *cumulative* association structures (updated cumulative mean and time-weighted cumulative) for HbA1c effects on time-to-T1DC traits. Scatter plots represent the P-values ($-\log_{10}$) for tests of $\beta_{g(l)}$ (X axis) and $\gamma_{g(k)}$ (Y axis) for DR/HbA1c, DN/HbA1c and DN/SBP trait pairs respectively based on: **(a.)** the joint model results and **(b.)** same results as (a.) on the X axis but substituting $P_{\gamma_{g(k)}}$ from CM-obs to assess the impact of measurement errors in longitudinal traits on the classification of direct and/or indirect associations. Significance levels $P^* = 1.7 \times 10^{-4}$ and $P^* = 0.05$ are indicated by red and dark grey lines.

(see **Figure S5** on next page)

a. Joint model results



b. CM-obs results



Time-dependent association structures for HbA1c effects on each time-to-T1DC trait

- Contemporaneous
- Time-weighted cumulative
- Updated cumulative mean