






























**Supplementary Table 3.** The structures and names of *N*-glycans identified in plasma by capillary electrophoresis

Glycan name and structure	Group	Glycan name and structure	Group
	S2 + ST + FB + LB		G2 + GT + B + FC + LB
	G2 + GT + LB		S2 + ST + B + FC + LB
	S1 + ST + LB		G0 + FC + LB
	S1 + ST + G1 + GT + LB		G1 + GT + FC + LB
	S2 + ST + LB		G1 + GT + FC + LB
	S3 + ST + FB + HB		G2 + GT + FC + LB
	S2 + ST + G1 + GT + HB		S1 + ST + G1 + GT + FC + LB
	S3 + ST + HB		S1 + ST + G1 + GT + FC + LB
	S3 + ST + G1 + GT + HB		S1 + ST + G1 + GT + FC + LB
	S4 + ST + HB		S2 + ST + FC + LB
	G0 + B + FC + LB		S3 + ST + G1 + GT + FC + HB
	G1 + GT + B + FC + LB		LB

-  Fucose
-  *N*-acetylglucosamine
-  Mannose
-  Galactose
-  *N*-acetylneuraminic acid

These 24 glycan structures can be grouped into 15 groups: bisecting GlcNAc (B group), sialic acid (non-sialylated (S0), mono-sialylated (S1), di-sialylated (S2), tri-sialylated and (S3), tetra-sialylated (S4), and total sialylated (ST)), galactose (agalactosylated (G0), mono-galactosylated (G1), di-galactosylated (G2), and total galactosylated (GT)), core fucose (FC group), branched fucose (FB group), high branched (HB group), and low branch (LB group).