




























Supplementary Table 4. The structures and names of *N*-glycans identified in isolated IgG by capillary electrophoresis

Glycan name and structure	Group	Glycan name and structure	Group
A2 	S2 + ST	G0FB 	G0 + F + B
A2F 	S2 + ST + F	G1F [6] 	G1 + GT + F
A2FB 	S2 + ST + F + B	G1F [3] 	G1 + GT + F
A2B 	S2 + ST + B	G1FB 	G1 + GT + F + B
G1FS1 	S1 + ST + F	G0 	G0
A1 	S1 + ST	G1S1 	S1 + ST
A1F 	S1 + ST + F	G2 	G2 + GT
A1FB 	S1 + ST + F	G2F 	G2 + GT + F
G0F 	G0 + F	G2FB 	G2 + GT + F + B
G1 	G1 + GT	G1B [3] 	G1 + GT + B
G0B 	G0 + B	G1B [6] 	G1 + GT + B

-  Fucose
-  *N*-acetylglucosamine
-  Mannose
-  Galactose
-  *N*-acetylneuraminic acid

These 22 glycan structures were grouped into 9 groups, depending on the presence or absence of four key monosaccharides: bisecting GlcNAc (B group), sialic acid (mon-sialylated (S1), di-sialylated (S2), and total sialylated (ST)), terminal galactose (agalactosylated (G0), mono-galactosylated (G1), di-galactosylated (G2), and total galactose (GT)), and fucose (F group).