

SUPPLEMENTARY MATERIAL

Dysbiosis and structural disruption of the respiratory microbiota in COVID-19 patients with severe and fatal outcomes

Alejandra Hernández-Terán¹, Fidencio Mejía-Nepomuceno¹, María Teresa Herrera², Omar Barreto³, Emma García³, Manuel Castillejos⁴, Celia Boukadida⁵, Margarita Matias-Florentino⁵, Alma Rincón-Rubio⁵, Santiago Avila-Rios⁵, Mario Mújica-Sánchez⁶, Ricardo Serna-Muñoz¹, Eduardo Becerril-Vargas⁶, Cristobal Guadarrama-Pérez⁷, Víctor Hugo Ahumada-Topete⁴, Sebastián Rodríguez¹, José Arturo Martínez-Orozco⁶, Jorge Salas-Hernández⁸, Rogelio Pérez-Padilla¹, Joel Armando Vázquez-Pérez^{1*}.

S1. Specifics for 16S rRNA gene amplification

PCR cycles:

95°C for 3 min, followed by 35 cycles at 95° C for 30s, 55°C for 30s, and 72°C for 30s, with a final extension at 72°C for 5 min

Primers used:

F - 5' CCTACGGGNGGCWGCAG 3'

R - 5' GACTACHVGGGTATCTAATCC 3'

Table S2. Wilcoxon paired comparisons of taxa abundance among groups.

| Taxa/ Comparison | Healthy vs Mild | Healthy vs Severe | Healthy vs Fatal | Healthy vs Non- COVID- 19-pneu. | Mild vs Severe | Mild vs Fatal | Mild vs Non- COVID-19- pneu | Severe vs Fatal | Severe vs Non- COVID- 19-pneu. | Fatal vs Non- COVID- 19-pneu |
|------------------------|--------------------|----------------------|---------------------|--|-------------------|------------------|-----------------------------------|--------------------|---|---------------------------------------|
| Phylum | | | | | | | | | | |
| Firmicutes | NS | NS | 0.03* | 0.04* | 0.05* | NS | NS | 0.02* | 0.03* | NS |
| Bacteroidetes | 0.03* | NS | 0.02* | 0.02* | NS | NS | NS | NS | NS | NS |
| Proteobacteria | 0.01* | NS | 0.04* | 0.03* | 0.001** | NS | NS | NS | NS | NS |
| Actinobacteria | NA | NA | NA | NA | NS | NS | 0.003** | NS | 0.007** | 0.004** |
| Fusobacteria | 0.004** | 0.02* | NS | 0.0003*** | NS | 0.02* | 0.0004*** | NS | 0.0001*** | 0.0004*** |
| TM7 | 0.0001*** | 0.002** | 0.006** | NA | 0.03* | 0.01* | NA | 0.008** | NA | NA |
| Synegistetes | NS | 0.01* | NS | NA | 0.004** | NS | NA | 0.002** | NA | NA |
| OD1 | NA | NA | NA | 0.0001*** | NA | NA | NA | NA | NA | NA |
| GN02 | NA | NA | NA | 0.0003*** | NA | NA | NA | NA | NA | NA |
| Plantomycetes | NA | NA | NA | 0.002** | NA | NA | NA | NA | NA | NA |
| Genus | | | | | | | | | | |
| <i>Prevotella</i> | NS | NS | NS | 0.002** | NS | NS | 0.001** | NS | 0.002** | 0.003** |
| <i>Streptococcus</i> | NS | 0.006** | NS | NS | 0.01* | NS | 0.04* | 0.02* | 0.01* | NS |
| <i>Veillonella</i> | 0.002** | 0.003** | 0.006** | 0.002** | NS | NS | 0.004** | NS | 0.003** | 0.008** |
| <i>Staphylococcus</i> | 0.0004*** | 0.003** | 0.0008*** | 0.005** | 0.0007*** | 0.0001*** | 0.006** | 0.002** | NS | 0.01* |
| <i>Moraxella</i> | 1.2e-05*** | NS | NA | 0.003** | 1.8e-07*** | NA | 2.3e-04*** | NA | 0.0003*** | NA |
| <i>Alloiococcus</i> | 3.2e-09*** | 2e-08*** | 1.6e-05*** | NA | NS | 0.0005*** | NA | 0.003** | NA | NA |
| <i>Corynebacterium</i> | 0.0002*** | 1.7e-04*** | 0.02** | 0.008** | 0.03* | 0.02* | 0.02* | 0.001** | 0.002** | NS |
| <i>Actinobacillus</i> | NS | 0.02** | 0.001** | NA | 0.03* | NS | NA | 0.05* | NA | NA |
| <i>Haemophilus</i> | NS | NS | 0.02* | 0.001** | NS | 0.05* | 0.03* | NS | 0.001** | 0.003** |
| <i>Neisseria</i> | 0.01* | 0.01* | NS | NA | NS | NS | NA | NS | NA | NA |
| <i>Magasphaera</i> | NS | NS | NS | 0.001** | NS | NS | 0.002** | NS | 0.004** | 0.001** |
| <i>Actinomyces</i> | 0.008** | 0.006** | 0.009** | 0.02* | NS | NS | 0.03* | NS | 0.03* | 0.007** |
| <i>Capnocytophaga</i> | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS |
| <i>Selenomonas</i> | 0.003** | 0.02* | 0.02* | NS | NS | NS | 0.01* | NS | 0.05* | NS |

p-values corresponds to paired Wilcoxon rank-sum test.

NS= non-significant *p*-value

NA= comparison not available due to absence of such taxa in one of the comparison group.




Figure S3. Main composition at phylum and genus level between lower and upper respiratory tract. A: median abundance of most abundant phyla in severe and fatal COVID-19. **B:** median abundance of most abundant genera in severe and fatal COVID-19. Asterisks denote statistical differences given by Wilcoxon rank-sum test test (p -values, * < 0.05, ** < 0.005, *** < 0.0005).

Table S4. Ružička metric as a proxy of dysbiosis among severity levels for COVID-19 and controls.

| Comparison | | Mean similarity | | Wilcoxon Test <i>p</i> value |
|-----------------|------------------------|-----------------|----------|------------------------------|
| Healthy | Diseased | Healthy | Diseased | |
| Healthy control | Mild COVID-19 | 0.94 | 0.89 | 0.05* |
| | Severe COVID-19 | 0.94 | 0.88 | 0.01* |
| | Fatal COVID-19 | 0.94 | 0.89 | 0.05* |
| | Non-COVID-19-pneumonia | 0.94 | 0.97 | 0.08 |

Intra-treatment similarity for healthy and diseased microbiota in terms of Ružička distance. Statistically significant differences were assessed with Wilcoxon rank-sum test. Dysbiosis was assumed when the similarities between the healthy microbiota samples were significantly higher than the similarities between the diseased microbiota samples (more dispersed).

Table S5. Topological metrics for all the calculated co-occurrence networks.

| Metric | Mild COVID-19 | Severe COVID-19 | Fatal COVID-19 |
|-----------------------------|----------------------|------------------------|-----------------------|
| Number of nodes | 148 | 84 | 74 |
| Number of edges | 4758 | 688 | 75 |
| Average number of neighbors | 65.86 | 16.35 | 2.31 |
| Diameter | 4 | 5 | 11 |
| Radius | 2 | 3 | 6 |
| Characteristic path length | 1.59 | 1.94 | 4.99 |
| Clustering | 0.788 | 0.47 | 0.155 |
| Density | 0.461 | 0.197 | 0.075 |
| Heterogeneity | 0.574 | 0.856 | 0.512 |
| Centralization | 0.391 | 0.563 | 0.127 |
| Connected components | 3 | 1 | 14 |