

SUPPLEMENTARY DIGITAL CONTENT 1

Title: Estimates of cases and hospitalizations averted by COVID-19 case investigation and contact tracing in 14 health jurisdictions in the United States

Authors: Seonghye Jeon*, Gabriel Rainisch*†, R. Ryan Lash, Patrick K. Moonan, John E. Oeltmann, Bradford Greening Jr., Bishwa B. Adhikari, Contact Tracing Impact Group**, Martin I. Meltzer

*Authors contributed equally to this work

**Contact Tracing Impact Group (alphabetically listed): Greta Anschuetz, Robert Bonacci, Brittany Byers, Joshua Clayton, Nickolas DeLuca, Matthew Donahue, Catherine Donovan, Veronica Fialkowski, Aaron Fleischauer, Heather Forbes, Clay Goddard, Heather Grome, Gibbie Harris, Susan Hayes, Blake Hendrickson, Julia Janssen, Sara Johnson, Amanda Jones, Catherine J. Knott, Reed Magleby, Stephen McCurdy, Alana McGrath, Heather McLaughlin, James Miller, Zack Moore, Michelle Morris, Jill Moses, Allison Newman, Sai Paritala, Caroline Q. Pratt, Lauren Prinzing, Pratima Raghunathan, Jonathan Steinberg, Alana Sulka, Christina G. Tan, Melanie Taylor, Puthiery Va, Kate Varela, Andee Weisbeck, Matthew Willis.

†**Corresponding Author:** Gabriel Rainisch

Author Affiliations: CDC COVID-19 Response Team

Data

We used a previously published evaluation study that assessed the COVID-19 case investigation and contact tracing efforts of 14 health departments (HDs) in 11 states and one tribal nation in the United States over 4-week asynchronous periods during June – October 2020, covering a total of 20 million population.¹ The study focused on assessing process indicators of the completeness and timeliness of case investigation and contact tracing (CICT) program implementation such as the proportions of cases interviewed, contacts notified and monitored, and the time between case identification and contact notification. We obtained the COVID-19 daily incidence data from the Centers for Disease Control and Prevention (CDC)'s COVID Data Tracker.² The tribal jurisdiction included in this data set is not represented in the CDC COVID Tracker database, and the data was directly provided to us by the jurisdiction.

Epidemiological Inputs

COVIDTracer Advanced³ is a spreadsheet-based compartmental Susceptible-Exposed-Infectious-Recovered (SEIR) epidemiological model, which illustrates the spread of a pathogen, resultant disease, and impact of interventions in a user-defined population. Readers can download the tool and enter input values of their choosing, exploring the impact of scenarios and assumptions beyond those covered in this manuscript. To model the clinical progression and transmission of disease using COVIDTracer Advanced, we used the following definitions and assumptions. A “case” was defined as a person who has been exposed, infected and subsequently becomes infectious, regardless of the presence of clinical symptoms. We assumed that for the first 3 days after exposure and infection, cases do not infect others. During days 4–5 post-infection, cases are pre-symptomatic but shed virus in amounts that may infect others.⁴⁻⁷ During days 6–14, the infected person can be symptomatic and shedding virus, albeit during days 11–14 the risk of onward transmission is relatively low but non-zero (the complete infectivity distribution is given in Table A1). We assumed that approximately 40% of cases are asymptomatic during days 6-14 yet have a risk of onward transmission equal to 75% of symptomatic cases (Table A2).

Table A1: Daily percentage risk of onward transmission by state of infectiousness and clinical symptoms.

Day Post exposure- and -infection	Daily percentage of risk of onward transmission ^a	Infectious person’s state
1	0.00%	<i>Infected, not yet infectious</i>
2	0.00%	
3	0.00%	
4	16.78%	<i>Infectious, pre-symptomatic</i>
5	18.03%	
6	17.07%	<i>Infectious, symptomatic</i>
7	14.52%	
8	11.27%	
9	8.10%	
10	5.48%	
11	3.55%	
12	2.26%	
13	1.46%	
14	1.48%	
Total	100%	

^aPercentages show when onward transmission might occur by day of infectiousness

Sources: He *et al.*^{4,5} and Ferretti *et al.*⁶. See also COVIDTracer Advanced manual.¹⁰










Table A2: Epidemiological parameters, values, and sources.

Parameter	Default Value	Source
Infected but not yet infectious period	3 days	CDC COVID-19 Pandemic Planning Scenarios ⁷
Pre-symptomatic and contagious (infectious) period	2 days	He <i>et al.</i> ^{4,5} , Ferretti <i>et al.</i> ⁶
Symptomatic and contagious (infectious) period	9 days	He <i>et al.</i> ^{4,5} , Ferretti <i>et al.</i> ⁶
New infections per case (R_0)	2.5	CDC COVID-19 Pandemic Planning Scenarios ⁷
% of cases that are asymptomatic	40%	CDC COVID-19 Pandemic Planning Scenarios ⁷
Infectiousness of asymptomatic cases (relative to symptomatic cases)	75%	CDC COVID-19 Pandemic Planning Scenarios ⁷

Table A3: Assumed proportion of cases by age group and infection-to-hospitalization rate, default values in COVIDTracer Advanced and sources.

Age group (year)	% of Total Cases	Source	% of all cases admitted to hospital care	Source
0 to 17	15%	CDC COVID Data Tracker ²	0.21	CDC COVID-19
18 to 64	55%		2.17	Response Team ⁸ ,
65+	30%		4.12	Wu <i>et al.</i> ⁹





Figure A1 (locations 1-9*). Fitted cumulative epidemic curve output from COVIDTracer Advanced and observed data for the 60-day period beginning at the time program evaluations began at each location.



* Locations 1-4 here correspond to locations 1-4 in the main text.

Figure A1 Continued (locations 10-14). Fitted cumulative epidemic curve output from COVIDTracer Advanced and observed data for the 60-day period beginning at the time program evaluations began at each location.

Location 10 not fitted due to unavailable data on days from case identification to isolation



Case Investigation and Contact Tracing Effectiveness

The effectiveness of case investigation and contact tracing is determined by the proportion of cases and their contacts that are effectively isolated and quarantined, preventing further transmission in the susceptible population. We assumed that confirmed cases are effectively isolated following case interviews. We further assumed that contacts are quarantined upon either contact notification, or through active monitoring. If infected contacts are left undetected, they will infect additional contacts. Therefore, we calculated the average proportion of cases and contacts isolated and quarantined for each location as follows:

Equation 1:

Contacts are assumed to be quarantined upon notification (hereafter named as “high effectiveness”):

$$[\% \text{ Cases interviewed} + R_0 * \% \text{ Contacts identified} * \% \text{ Contacts notified}] / (1+R_0)$$

Equation 2:

Contacts are assumed to be quarantined through active monitoring (hereafter named as “low effectiveness”):

$$[\% \text{ Cases interviewed} + (R_0 * \% \text{ Contacts identified} * \% \text{ Contacts monitored})] / (1+R_0)$$

where R_0 is the assumed number of new infections per case without any interventions and when the population is entirely susceptible to infection (Table A2). The % Cases interviewed, % Contacts notified, and % Contacts monitored were process metrics gathered in the evaluation study,¹ and % Contacts identified was an intermediate value calculated as the number of named contacts divided by the expected number of contacts per case:

$$\# \text{ Contacts named} / (\# \text{ Cases reported} * \text{Average} \# \text{ Contacts per Case in each location})$$

In general, the % of Contacts notified was higher than the % of Contacts actively monitored. Therefore, the estimated effectiveness was higher when contacts were assumed to be quarantined upon notification. Likewise, the estimated effectiveness was lower when active monitoring was assumed to be required for contact quarantine.

In addition, reducing the time from case identification to effective isolation is critical for case investigation and contact tracing to succeed. The longer the cases and contacts interact with the

susceptible population, the greater the opportunity for onward transmission. In practice, cases with no known exposure can be identified and isolated only after symptom onset, and cases with known exposures (*i.e.*, contacts that eventually become infected cases) can begin quarantine upon contact notification (even potentially prior to symptom onset). We assumed asymptomatic cases can only be identified and isolated if they are notified through case investigation and contact tracing.

For the purposes of our study we assumed the proportions of cases with no known exposure and cases with known exposures were equal (*i.e.*, 50/50 breakdown) since we didn't have data on what prompted case identification in each location. Therefore, for each location the days to effective case isolation was determined by taking the average of the days to effective isolation between case groups with known and no known exposures. The time to effective case isolation for each of the two case groups was determined as follows:

For symptomatic cases with no known exposures (*i.e.*, symptoms prompt identification):

We assumed cases experience a 5-day pre-symptomatic period (See Table A2), get tested the next day after symptom onset, wait the number of days observed by Lash *et al.*¹ to learn of their positive result, and begin effective isolation the day after learning this result. See the "Index Case" row in Figure A2 for a visual depiction of this timeline.

For cases with known exposures (*i.e.*, those who were notified they were a contact and eventually became a case):

We assumed contacts quarantine the day after being notified as a contact and that these individuals could contribute to onward transmission based on when they were exposed. Since we did not have information on when exposures actually occurred, we assumed these individuals' exposures occurred at the midpoint of their potential exposure window (in days). We identified the earliest date in this window as the first day of infectiousness among cases to which contacts were exposed. Based on our assumed 5-day pre-symptomatic period for symptomatic cases (described above), this was two days prior to the symptom onset date in cases exposing the contact. We identified the latest possible exposure date as the date the cases exposing them were notified of their positive case status (since they began isolation the next day). See both "Contacts" rows in Figure A2 for a visual depiction of this timeline.


The days between cases with known exposures becoming infected and their exposure notification can vary from what we assumed. For example, cases may take longer to become symptomatic, or get tested the same day they become symptomatic or begin their isolation on the same day as their results notification. Similarly, contacts which become cases may be exposed earlier or later than we assumed and may make up a larger or smaller share of the case pool. Table A6 shows the impact of varying our assumed time to case isolation by 1-day higher and lower prior to completing our fitting procedure for determining the share of transmission reductions attributable to NPIs and Case Investigation and Contact Tracing. The sensitivity analysis associated with 1-day changes described and shown in the main text (Figures 2 and A3) is different in that it shows the impact of varying notification speed when NPI effects are held constant (*i.e.*, given the originally derived share of transmission reductions attributable to NPIs, how speeding or slowing notification would have affected averted cases).

Figure A2. Illustrative example of the timing of case isolation and contact quarantine based on reported data from Location 1

	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Day 8	Day 9	Day 10	Days from Exposure to Isolation
Index Case	Exposed			Begin Contagious		Symptom Onset	Tested		Results Notification	Begin Isolation	9
Contacts (Earliest possible exposure)				Exposed					Exposure Notification	Begin Quarantine	6
Contacts (Latest possible exposure)									Exposed	Begin Quarantine	1

Notes: Location 1 reported 2 days from specimen collection to results notification and 2 days from specimen collection to contact notification (Table 1). The index case (symptomatic case with no known exposure) began showing symptoms on day 6 post exposure-and-infection, got tested on day 7, and was notified of the positive test results on day 9. Its contacts (cases with known exposure) were exposed between day 4 to 9 and notified of their exposure on day 9. Therefore, both the index case and its contacts began isolation and quarantine on day 10.

FIGURE A3. Effects of improvements and constraints to case investigation and contact tracing performance measures on the baseline percent cases averted by the program* when monitoring contacts is assumed necessary for effective quarantine (Low CICT effectiveness approach)



* Percent cases averted by cases investigated and contact tracing (CICT) calculated as percentage of total cases averted if only nonpharmaceutical interventions (NPIs) were implemented.

Notes. Baseline results shown in Table 2. Cases Interviewed and Contacts Monitored were capped at 100% and 0% when the baseline percentage interviewed was greater than 80% or less than 20%. Similar figure in the main text (Figure 2) is for the High CICT effectiveness scenario: when simply notifying contacts is assumed sufficient to trigger effective quarantine.

Table A4 (locations 1-7). Estimated impacts of case investigation and contact tracing (CICT), and nonpharmaceutical interventions (NPIs), by location^a over 60-day period after contact tracing evaluations initiated

	Location 1	Location 2	Location 3	Location 4	Location 5	Location 6	Location 7
Scenario Inputs							
Population	Category B	Category A	Category C	Category A	Category C	Category C	Category C
Start date of observation	7/26/2020	8/1/2020	7/30/2020	7/5/2020	8/23/2020	6/28/2020	6/15/2020
COVID-19 cases							
Total cases before the start of observation	1 396	18 488	4 817	10 724	2 231	13	453
Cases in the last 14 days	113	4 686	870	3 853	43	5	33
Trend in the last 14 days	plateaued	plateaued	plateaued	slowly increasing	slowly decreasing	slowly increasing	plateaued
CICT Effectiveness							
% of cases and contacts isolated ^b	39.5 – 86.0%	52.7 – 55.0%	26.1 – 50.5%	11.8 – 12.1%	76.6 – 81.4%	70.3 – 71.0%	21.4 – 30.2%
Days from infection to isolation ^c	6 days	8 days	9 days	10 days	6 days	6 days	7 days
Scenario Results							
Transmission Fraction							
Reduction from CICT	8.6 – 26.2%	5.0 – 5.2%	1.4 – 2.7%	0.4 – 0.4%	28.7 – 31.7%	27.8 – 28.1%	4.8 – 7.0%
Reduction from NPIs ^d	54.6 – 36.6%	57.6 – 57.3%	63.5 – 62.0%	61.0 – 61.0%	22.2 – 19.1%	18.0 – 17.7%	32.8 – 30.9%
Remaining Transmission (100% minus above values)	36.8 – 37.2%	37.4 – 37.5%	35.1 – 35.3%	38.6 – 38.6%	49.1 – 49.2%	54.2 – 54.2%	62.4 – 62.1%
Additional Cases Averted by CICT (%), 60 days^e	651 – 9 480 (67.1 – 96.8%)	12 598 – 13 568 (47.1 – 48.8%)	344 – 768 (15.4 – 28.8%)	859 – 882 (4.4 – 4.5%)	5 238 – 6 879 (96.1 – 97.0%)	4 319 – 4 450 (93.8 – 94.0%)	1 684 – 2 738 (38.3 – 50.0%)
Additional Hospitalizations Averted by CICT (%), 60 days^e	16 – 233 (67.1 – 96.8%)	310 – 333 (47.1 – 48.8%)	8 – 19 (15.4 – 28.8%)	21 – 22 (4.4 – 4.5%)	129 – 160 (96.1 – 97.0%)	106 – 109 (93.8 – 94.0%)	41 – 67 (38.3 – 50.0%)

[†] To preserve anonymity, populations are categorized by population size as follows: Category A > 1 million; Category B >500 000 to ≤ 1 million; Category C: ≤ 500 000.

^aCase investigation and contact tracing implemented per scenarios in Table 4A and effects were assumed constant over 60 days.

^bCalculated as follows using values observed at locations during case investigation and contact tracing evaluations and an assumed $R_0=2.5$:

Minimum (low) value assumes monitoring is required for effective quarantine of contacts:

$$[\% \text{ Cases interviewed} + (R_0 * \% \text{ Contacts identified} * \% \text{ Contacts actively monitored})] / (1+R_0)$$

Maximum (high) value assumes contact notification is sufficient for effective quarantine of contacts:

$$[\% \text{ Cases interviewed} + (R_0 * \% \text{ Contacts identified} * \% \text{ Contacts notified})] / (1+R_0)$$

^cThe average length of time from exposure-and-infection to isolation (including contacts which later became cases). See CICT Effectiveness in the Supplement.

^dNPI interventions including masks use, social distancing, school and restaurant closures, etc. Low NPI effectiveness values were generated with the fitting process when CICT effectiveness was high; similarly, high NPI effectiveness values were generated when CICT effectiveness was low.

^ePercent calculated as (Total Cases Averted when both CICT and NPIs implemented - Total Cases Averted when only NPIs implemented) / Total Cases Averted when both CICT and NPIs implemented). For example, for every 100 cases prevented by nonpharmaceutical interventions in locations 1-4, CICT averted between 4.4 and 96.8 additional cases.

TABLE A4 Continued (locations 8-14). Estimated impacts of case investigation and contact tracing (CICT), and nonpharmaceutical interventions (NPIs), by location^a over 60-day period after contact tracing evaluations initiated

	Location 8	Location 9	Location 10 ^f	Location 11	Location 12	Location 13	Location 14
Scenario Inputs							
Population	Category C	Category C	Category B	Category B	Category A	Category A	Category A
Start date of observation	6/15/2020	6/21/2020	7/1/2020	10/4/2020	6/1/2020	8/1/2020	8/2/2020
COVID-19 cases							
Total cases before the start of observation	931	218	9 237	23 988	4 307	26 211	184 358
Cases in the last 14 days	249	44	2 221	5 545	1 685	3 850	5 835
Trend in the last 14 days	plateaued	rapidly increasing	plateaued	slowly increasing	slowly increasing	plateaued	plateaued
CICT Effectiveness							
% of cases and contacts isolated ^b	50.6–52.7%	54.5 – 61.1	22.1 – 22.1%	21.3 – 30.8%	43.4 – 43.9%	22.2 – 52.7%	32.2 – 35.6%
Days from infection to isolation ^c	9 days	7 days	Not Available	7 days	8 days	8 days	8 days
Scenario Results							
Transmission Fraction							
Reduction from CICT	3.3 – 3.5%	11.7 – 13.4%		3.6 – 5.4%	4.7 – 4.8%	2.2 – 5.5%	3.1 – 3.5%
Reduction from NPIs ^d	54.0–53.9%	36.2 – 34.8%		49.6 – 47.9%	51.2 – 51.1%	56.7 – 53.4%	56.3 – 55.9%
Remaining Transmission (100% minus above values)	42.7–42.6%	52.1 – 51.8%		46.8 – 46.7%	44.1 – 44.1%	41.1 – 41.1%	40.6 – 40.6%
Additional Cases Averted by CICT (%), 60 days^e	733 – 768 (33.3–34.5%)	4 857–6 112 (70.0–74.7%)		28 141–45 869 (31.0–42.2%)	12 608–12 904 (42.0 – 42.4%)	5 386–17 264 (23.2–49.3%)	11 634–13 378 (32.3 – 35.3%)
Additional Hospitalizations Averted by CICT (%), 60 days^e	18 – 19 (33.3–34.5%)	119–150 (70.0–74.7%)		692–1 127 (31.0–42.2%)	310 – 317 (42.0 – 42.4%)	132–424 (23.2–49.3%)	286 – 329 (32.3 – 35.3%)

[†] To preserve anonymity, populations are categorized by population size as follows: Category A > 1 million; Category B >500 000 to ≤ 1 million; Category C: ≤ 500 000.

^aCase investigation and contact tracing implemented per scenarios in Table 4A and effects were assumed constant over 60 days.

^bCalculated as follows using values observed at locations during case investigation and contact tracing evaluations and an assumed $R_0=2.5$:

Minimum (low) value assumes monitoring is required for effective quarantine of contacts:

$$[\% \text{ Cases interviewed} + (R_0 * \% \text{ Contacts identified} * \% \text{ Contacts actively monitored})] / (1+R_0)$$

Maximum (high) value assumes contact notification is sufficient for effective quarantine of contacts:

$$[\% \text{ Cases interviewed} + (R_0 * \% \text{ Contacts identified} * \% \text{ Contacts notified})] / (1+R_0)$$

^cThe average length of time from exposure-and-infection to isolation (including contacts which later became cases). See CICT Effectiveness in the Supplement.

^dNPI interventions including masks use, social distancing, school and restaurant closures, etc. Low NPI effectiveness values were generated with the fitting process when CICT effectiveness was high; similarly, high NPI effectiveness values were generated when CICT effectiveness was low.

^ePercent calculated as (Total Cases Averted when both CICT and NPIs implemented - Total Cases Averted when only NPIs implemented) / Total Cases Averted when both CICT and NPIs implemented). For example, for every 100 cases prevented by nonpharmaceutical interventions in locations 1-4, CICT averted between 4.4 and 96.8 additional cases.

^fLocation 10 did not report the median days from case identification to isolation, thus the impact of CICT could not be estimated.

TABLE A5(a). Estimated cases averted with improving and constraining case investigation and contact tracing performance measures relative to baseline, assuming contact notification is a sufficient trigger for contacts to effectively quarantine, by location, over 60-days after program evaluations were initiated

	<u>Location 1</u>	<u>Location 2</u>	<u>Location 3</u>	<u>Location 4</u>
	Baseline Values: Cases Interviewed = 99% Contacts Notified = 95% Days to isolation = 6	Baseline Values: Cases Interviewed = 83% Contacts Notified = 61% Days to isolation = 8	Baseline Values: Cases Interviewed = 91% Contacts Notified = 85% Days to isolation = 9	Baseline Values: Cases Interviewed = 33% Contacts Notified = 54% Days to isolation = 10
Performance measure				
Cases Interviewed (%)				
Baseline +20% ^a	9 486 (96.8%)	14 431 (52.0%)	856 (32.1%)	1 286 (6.6%)
Baseline	9 480 (96.8%)	13 568 (48.8%)	768 (28.8%)	882 (4.5%)
Baseline -20%	9 386 (95.8%)	12 509 (45.0%)	692 (26.0%)	471 (2.4%)
Contacts Notified (%)				
Baseline +20% ^a	9 525 (97.2%)	15 314 (55.1%)	824 (30.9%)	953 (4.9%)
Baseline	9 480 (96.8%)	13 568 (48.8%)	768 (28.8%)	882 (4.5%)
Baseline -20%	9 245 (94.4%)	11 605 (41.8%)	691 (25.9%)	810 (4.2%)
Median days from cases being infected to their contacts isolated				
3 days faster than Baseline	-- ^b	23 479 (84.5%)	1 775 (66.6%)	3 289 (16.9%)
2 days faster than Baseline	-- ^b	21 106 (76.0%)	1 451 (54.5%)	2 232 (11.5%)
1 day faster than Baseline	9 703 (99.9%)	17 667 (63.6%)	1 096 (41.1%)	1 435 (7.4%)
Baseline	9 480 (96.8%)	13 568 (48.8%)	768 (28.8%)	882 (4.5%)
1 day slower than Baseline	8 753 (89.3%)	9 572 (34.5%)	504 (18.9%)	520 (2.7%)
2 days slower than Baseline	7 269 (74.2%)	6 281 (22.6%)	312 (11.7%)	291 (1.5%)
3 days slower than Baseline	5 343 (54.5%)	3 879 (14.0%)	181 (6.8%)	145 (0.7%)

Notes

^aThis value was capped at 100% when the baseline percentage interviewed was greater than 80%.

^bThis value not calculated since the minimum median days from case infection to contacts being isolated is six (5 days from infection to symptom onset + 1 day for testing).

TABLE A5(b). Estimated cases averted with improving and constraining case investigation and contact tracing performance measures relative to baseline, assuming active monitoring is required for effective contact quarantine, by location, over 60-days after program evaluations were initiated

Performance measure	Location 1	Location 2	Location 3	Location 4
	Baseline Values:	Baseline Values:	Baseline Values:	Baseline Values:
	Cases Interviewed = 99% Contacts Monitored = 19% Days to isolation = 6	Cases Interviewed = 83% Contacts Monitored = 56% Days to isolation = 8	Cases Interviewed = 91% Contacts Monitored = 0% Days to isolation = 9	Cases Interviewed = 33% Contacts Monitored = 48% Days to isolation = 10
Cases Interviewed (%)				
Baseline +20% ^a	654 (67.5%)	13 448 (50.3%)	375 (16.7%)	1 263 (6.5%)
Baseline	651 (67.1%)	12 598 (47.1%)	344 (15.4%)	859 (4.4%)
Baseline -20%	594 (61.3%)	11 555 (43.2%)	273 (12.2%)	448 (2.3%)
Contacts Quarantined (%)				
Baseline +20% ^a	744 (76.8%)	14 319 (53.5%)	414 (18.5%)	931 (4.8%)
Baseline	651 (67.1%)	12 598 (47.1%)	344 (15.4%)	859 (4.4%)
Baseline -20% ^b	529 (54.6%)	10 664 (39.9%)	344 (15.4%)	787 (4.1%)
Median days from cases being infected to their contacts isolated				
3 days faster than Baseline	--‡	22 207 (83.0%)	934 (41.7%)	3 209 (16.5%)
2 days faster than Baseline	--‡	19 843 (74.2%)	715 (31.9%)	2 177 (11.2%)
1 day faster than Baseline	750 (77.4%)	16 499 (61.7%)	512 (22.8%)	1 399 (7.2%)
Baseline	651 (67.1%)	12 598 (47.1%)	344 (15.4%)	859 (4.4%)
1 day slower than Baseline	522 (53.9%)	8 851 (33.1%)	220 (9.8%)	507 (2.6%)
2 days slower than Baseline	386 (39.8%)	5 794 (21.7%)	135 (6.0%)	283 (1.5%)
3 days slower than Baseline	264 (27.2%)	3 573 (13.4%)	77 (3.5%)	141 (0.7%)

Notes

^aThis value was capped at 100% when the baseline percentage interviewed was greater than 80%.

^bThis value was kept at 0% when the baseline percentage monitored was less than 20%.

TABLE A6(a). Estimated impacts of case investigation and contact tracing and nonpharmaceutical interventions in Location 1 over 60-day period after program evaluations were initiated, varying the days from infection to isolation

Days from Infection to Isolation	5	6 (default)	7
Transmission Fraction			
Reduction from CICT	12.8 – 47.5%	8.6 – 26.2%	5.6 – 15.1%
Reduction from All Other Interventions (NPIs)	50.5 – 15.2%	54.6 – 36.6%	57.6 – 47.9%
Remaining Transmission (100% minus above values)	36.7 – 37.3%	36.8 – 37.2%	36.8 – 37.0%
Cases Averted by CICT (%), 60 days	1 338 – 97 014 (80.8 – 99.7%)	651 – 9 480 (67.1 – 96.8%)	339 – 2 009 (51.4 – 86.3%)
Hospitalizations Averted by CICT (%), 60 days	33 – 2 433 (80.8 – 99.7%)	16 – 233 (67.1 – 96.8%)	8 – 49 (51.4 – 86.3%)

TABLE A6(b). Estimated impacts of case investigation and contact tracing and nonpharmaceutical interventions in Location 2 over 60-day period after program evaluations were initiated, varying the days from infection to isolation

Days from Infection to Isolation	7	8 (default)	9
Transmission Fraction			
Reduction from CICT	8.1 – 8.5%	5.0 – 5.2%	3.0 – 3.2%
Reduction from All Other Interventions (NPIs)	54.5 – 54.0%	57.6 – 57.3%	59.6 – 59.4%
Remaining Transmission (100% minus above values)	37.4 – 37.5%	37.4 – 37.5%	37.4 – 37.4%
Cases Averted by CICT (%), 60 days	25 552 – 28 102 (64.4 – 66.4%)	12 598 – 13 568 (47.1 – 48.8%)	6 663 – 7 108 (32.0 – 33.3%)
Hospitalizations Averted by CICT (%), 60 days	628 – 691 (64.4 – 66.4%)	310 – 333 (47.1 – 48.8%)	164 – 175 (32.0 – 33.3%)

TABLE A6(c). Estimated impacts of case investigation and contact tracing and nonpharmaceutical interventions in Location 3 over 60-day period after program evaluations were initiated, varying the days from infection to isolation

Days from Infection to Isolation	8	9 (default)	10
Transmission Fraction			
Reduction from CICT	2.2 – 4.5%	1.4 – 2.7%	0.8 – 1.6%
Reduction from All Other Interventions (NPIs)	62.7 – 60.2%	63.6 – 62.0%	64.1 – 63.2%
Remaining Transmission (100% minus above values)	35.1 – 35.3%	35.1 – 35.3%	35.1 – 35.2%
Cases Averted by CICT (%), 60 days	573 – 1 407 (23.3 – 42.7%)	344 – 768 (15.4 – 28.8%)	203 – 427 (9.7 – 18.4%)
Hospitalizations Averted by CICT (%), 60 days	14 – 35 (23.3 – 42.7%)	8 – 19 (15.4 – 28.8%)	5 – 10 (9.7 – 18.4%)

TABLE A6(d). Estimated impacts of case investigation and contact tracing and nonpharmaceutical interventions in Location 4 over 60-day period after program evaluations were initiated, varying the days from infection to isolation

Days from Infection to Isolation	9	10 (default)	11
Transmission Fraction			
Reduction from CICT	0.7 – 0.7%	0.4 – 0.4%	0.2 – 0.2%
Reduction from All Other Interventions (NPIs)	60.8 – 60.8%	61.0 – 61.0%	61.2 – 61.2%
Remaining Transmission (100% minus above values)	38.5 – 38.5%	38.6 – 38.6%	38.6 – 38.6%
Cases Averted by CICT (%), 60 days	1 438 – 1 476 (7.2 – 7.4%)	859 – 882 (4.4 – 4.5%)	493 – 506 (2.6 – 2.7%)
Hospitalizations Averted by CICT (%), 60 days	35 – 36 (7.2 – 7.4%)	21 – 22 (4.4 – 4.5%)	12 – 12 (2.6 – 2.7%)

Adjustment for Under-reported and Under-detected Cases

Table A7(a). Estimated impacts of case investigation and contact tracing and nonpharmaceutical interventions over 60-day period after program evaluations were initiated, with and without adjustment for under-reported asymptomatic cases; Locations 1 and 2

	<u>Location 1</u> Baseline	<u>Location 1</u> Multiplying reported cases by x2	<u>Location 2</u> Baseline	<u>Location 2</u> Multiplying reported cases by x2
Transmission Fraction				
Reduction from CICT	8.6 – 26.2%	8.7 – 26.3%	5.0 – 5.2%	5.0 – 5.3%
Reduction from All Other Interventions (NPIs)	54.6 – 36.6%	54.5 – 36.4%	57.6 – 57.3%	57.3 – 57.1%
Remaining Transmission (100% minus above values)	36.8 – 37.2%	36.8 – 37.3%	37.4 – 37.5%	37.7 – 37.6%
Cases Averted by CICT (%), 60 days	651 – 9 480 (67.1 – 96.8%)	1 296 – 18 707 (67.0 – 96.7%)	12 598 – 13 568 (47.1 – 48.8%)	25 173 – 26 719 (46.9 – 48.6%)
Hospitalizations Averted by CICT (%), 60 days	16 – 233 (67.1 – 96.8%)	32 – 460 (67.0 – 96.7%)	310 – 333 (47.1 – 48.8%)	619 – 657 (46.9 – 48.6%)

Table A7(b). Estimated impacts of case investigation and contact tracing and nonpharmaceutical interventions over 60-day period after program evaluations were initiated, with and without adjustment for under-reported asymptomatic cases; Locations 3 and 4

	<u>Location 3</u> Baseline	<u>Location 3</u> Multiplying reported cases by x2	<u>Location 4</u> Baseline	<u>Location 4</u> Multiplying reported cases by x2
Transmission Fraction				
Reduction from CICT	1.4 – 2.7%	1.4 – 2.8%	0.4 – 0.4%	0.4 – 0.4%
Reduction from All Other Interventions (NPIs)	63.5 – 62.0%	62.7 – 61.2%	61.0 – 61.0%	60.4 – 60.4%
Remaining Transmission (100% minus above values)	35.1 – 35.3%	35.9 – 36.0%	38.6 – 38.6%	39.2 – 39.2%
Cases Averted by CICT (%), 60 days	344 – 768 (15.4 – 28.8%)	678 – 1 501 (15.2 – 28.5%)	859 – 882 (4.4 – 4.5%)	1 674 – 1 718 (4.3 – 4.4%)
Hospitalizations Averted by CICT (%), 60 days	8 – 19 (15.4 – 28.8%)	17 – 37 (15.2 – 28.5%)	21 – 22 (4.4 – 4.5%)	41 – 42 (4.3 – 4.4%)

References

1. Lash RR, Moonan PK, Byers BL, et al. COVID-19 contact tracing in the United States, 2020. *JAMA Network Open*. 2021;In Press
2. CDC COVID Data Tracker. <https://covid.cdc.gov/covid-data-tracker/>
3. *COVIDTracer Advanced: A Planning Tool to Illustrate the Resources Needed to Conduct Contact Tracing and Monitoring of coronavirus disease 2019 (COVID-19) Cases and the Potential Impact of Community Interventions and Contact Tracing Efforts on the Spread of COVID-19* 2020. <https://www.cdc.gov/coronavirus/2019-ncov/php/contact-tracing/COVIDTracerTools.html>
4. He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nature Medicine*. 2020;26(5):672-675. doi:10.1038/s41591-020-0869-5
5. He X, Lau EHY, Wu P, et al. Author Correction: Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nature Medicine*. 2020;26(9):1491-1493. doi:10.1038/s41591-020-1016-z
6. Ferretti L, Wymant C, Kendall M, et al. Quantifying SARS-CoV-2 transmission suggests epidemic control with digital contact tracing. *Science*. 2020;368(6491):eabb6936. doi:10.1126/science.abb6936
7. CDC. COVID-19 Pandemic Planning Scenarios. Feb 14, 2021. Updated Sep 10, 2020. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/planning-scenarios.html>
8. CDC COVID-19 Response Team. Severe Outcomes Among Patients with Coronavirus Disease 2019 (COVID-19) — United States, February 12–March 16, 2020. *Morbidity and Mortality Weekly Report*. 2020;69(12):343-346.
9. Wu SL, Mertens AN, Crider YS, et al. Substantial underestimation of SARS-CoV-2 infection in the United States. *Nature Communications*. 2020;11(4507)doi:<https://doi.org/10.1038/s41467-020-18272-4>