

A rapid and simple protocol for concentration of SARS-CoV-2 from sewage

Diana P. Wehrendt^{a,b}, Mariana G. Massó^c, Adrián Gonzales Machuca^c, Claudia V. Vargas^d, Melina E. Barrios^{b,e}, Josefina Campos^g, Damián Costamagna^h, Luis Bruzzoneⁱ, Daniel M. Cisterna^g, Néstor Gabriel Iglesias^{b,j}, Viviana A. Mbayed^{b,e,f}, Elsa Baumeister^k, Daniela Centrón^{b,c}, María Paula Quiroga^{b,c}, Leonardo Erijman^{a,b,l*}

Supplemental Material

Table S1: Effect of storage during 7 days at 4°C of concentrated sewage samples on RT-qPCR detection of SARS-CoV-2

Sample ID	Target gene			
	N		Orf1ab	
	Before storage	After storage	Before storage	After storage
AK31	31,18	33,24	32,52	38,89
AK32	31,33	33,27	32,10	36,72
SF29	30,84	30,59	31,24	30,80
SF31	35,37	34,1	35,85	n.d.
SF32	34,52	n.d.	44,26	37,67
SP29	31,87	30,5	33,88	31,48
SP32	35,94	35,2	n.d.	n.d.
SV31	33,63	33,06	n.d.	34,23
SV32	29,76	30,3	31,05	31,91

n.d.: non- detection.