

Enrollment




Figure S1: CONSORT flow diagram of study

		Dose of carbohydrate				
		10%	15%	20%	25%	30%
		g	g	g	g	g
Breakfast	Bagels	20	35	50	60	80
	Butter	22	24	18	15	15
	Keto Pancake	125	120	105	100	90
	Walnuts	25	20	25	25	20
Lunch	White fish	80	80	80	70	60
	Rice	35	65	100	130	150
	Cauliflower	60	60	60	60	60
	Sunflower oil	22	22	22	22	17
	Lettuce	35	35	35	35	35
	Cucumber	50	50	50	50	50
	Tomatoes	50	50	50	50	50
	Feta cheese	60	60	54	55	60
	Olive oil	25	20	19	15	15
Dinner	Chicken breast	60	60	55	50	50
	Croutons	20	30	40	50	60
	Avocado	130	120	110	100	70
	Parmesan cheese	15	15	12	15	15
	Lettuce	30	30	30	30	30
	Olives	18	18	18	18	18
	Olive oil	25	22	20	17	15
	Dates	10	15	20	25	40
Snack	Greek yoghurt	120	160	140	150	150
	Cream	10	10	5	0	0
	Raspberries	10	25	30	35	40

	Mango	0	30	50	90	115
	Pistachio nuts	25	10	17	15	12


Table S1: Example of how the menu is altered with each dose.

	Mean 24-hour	Post-prandial	CONGA	MAGE	log(LBGI)	HBGI
Constant	16.949*** (3.809)	18.217*** (4.603)	4.294* (1.952)	11.285* (5.215)	-12.029* (5.366)	19.721* (7.579)
Dose	0.017 (0.009)	0.021 (0.011)	0.007 (0.008)	0.014 (0.019)	-0.024 (0.029)	0.032 (0.022)
Age	-0.166* (0.069)	-0.173 (0.083)	-0.047 (0.035)	-0.128 (0.094)	0.179 (0.095)	-0.310 (0.136)
Male	-2.205* (0.719)	-2.717* (0.869)	-0.662 (0.364)	-1.836 (0.976)	3.007* (0.988)	-3.514* (1.426)
Kuwati	0.963 (0.715)	0.674 (0.864)	-0.072 (0.365)	-0.263 (0.977)	-1.967 (1.000)	2.447 (1.421)
Weight change (kg)	0.076 (0.098)	0.118 (0.120)		0.474* (0.195)	0.219 (0.295)	0.326 (0.233)
Mean steps per day	-0.049 (0.044)	(0.054)	0.007 (0.033)	0.024 (0.082)	-0.033 (0.109)	-0.023 (0.102)
% Variance explained†	70%	69%	42%	49%	22%	60%

Table S3: Parameter estimates with standard errors from the linear mixed-effects models (*p<0.05; **p<0.01; ***p<0.001)


†Random effect for intercept explains xx% of the variance that remains after the variance has been explained by the fixed effects.

Supplementary data: Figure S2: Post-prandial glucose excursions for each participant per meal, per day.




Supplementary data: Figure S2: Post-prandial glucose excursions for each participant per meal, per day.

Dose: 15%



Supplementary data: Figure S2: Post-prandial glucose excursions for each participant per meal, per day.



Supplementary data: Figure S2: Post-prandial glucose excursions for each participant per meal, per day.



Supplementary data: Figure S2: Post-prandial glucose excursions for each participant per meal, per day.

Dose: 30%

