

SUPPLEMENTS

Table S1. Description and categorization of variables used in analyses.

Variable	Question in Questionnaire	Answer possibilities	Calculation / Categorisation	Timepoint
Test results				
Positive SARS-CoV-2 test	Did you have a positive SARS CoV-2 swab test?	True; False	True; False	Weekly
Number of negative swab tests	Have you been tested for SARS-CoV-2 in the past week?	True; False	Cumulative number of negative tests up to this date	Weekly
	What was the result?	Positive; Negative; Pending		Weekly
Exposure to COVID-19				
Frequency of contacts to COVID-19 patients	How many COVID-19 patients were you exposed to since March 2020?	None; 1 to 5; 6 to 10; 11 to 20; more than 20; I don't know	None, 1-20, >20	Follow-up
Frequency of contacts to coworkers with COVID-19	How many SARS-CoV-2 confirmed coworkers were you exposed to since March 2020?	1; 2; 3; 4 or more	1;2;3;4	Follow-up
Any positively tested household member	Has someone of your household or your intimate partner been tested positive since March 2020?	True; False	True; False	Weekly
Patient exposure (<i>time dependent</i>)	Did you have contact to a confirmed COVID-19 patient in hospital last week?	True; False	True; False	Weekly
Coworker exposure (<i>time dependent</i>)	Did you have contact to a SARS-CoV-2 confirmed coworker without wearing a mask last week?	True; False	True; False	Weekly
Household exposure (<i>time dependent</i>)	Has someone from your household been tested positive for SARS-CoV-2 last week?	True; False	True; False	Weekly
Other exposure (<i>time dependent</i>)	Has someone other from your surroundings been tested positive for SARS-CoV-2 last week?	True; False	True; False	Weekly
Sociodemographic data				
Age	What is your date of birth?	Day/Month/Year	(2020–birthyear)/10	Baseline
Sex	What is your sex?	Male; Female; Not specified	Male; Female and other	Baseline
Living in Germany or Austria	Zip code of your hometown	Numbers	GE/AU vs. CH	Baseline
Number of additional persons in household	How many people live in your place of residence?	Numbers	Numbers from 0 to 4 (larger numbers truncated to 4)	Baseline
Child in household	Year of birth for every household contact	Year	At least one year \geq 2008	Baseline
Medical conditions				
Comorbidity	Do you suffer from any of the following diseases?	None; Arterial hypertension; Diabetes mellitus; Heart disease; COPD; Asthma; Liver disease (Fatty liver; Cirrhosis); Cancer; Rheumatologic disease (Arthritis, Lupus, Vasculitis); Hay fever; Hypothyroidism; Chronic inflammatory bowel disease (M. Crohn, Colitis ulcerosa), Intermittent claudication (Peripheral artery disease), Stroke	None = False; All others = True	Baseline
Active smoker	Do you smoke or did you use to smoke?	No; Used to smoke; Active smoker	Active smoker; other	Baseline
Pregnancy during study	Are you pregnant at the moment?	True; False	True; False	Follow-up
Behaviour outside of work				
Prophylactic home remedies	Which preventive preparation do you take against COVID-19?	None; Echinacea; Zinc; Vitamin C; Umckaloabo/Kaloba; Vitamin B12;	None = False; All others =	Baseline

Social leisure activities	Which activities do you perform regularly at the moment?	Homeopathics; Others None; Sportsclub; Choir; Orchestra/Music group (Active musician); Concert/Theatre; Cinema; Church; Yoga/Fitness; Restaurant/Bar At home; Outside; On public transport; While shopping; In hospital area; In the workspace; During contact with other people	True None = False; All others = True	Baseline
Wearing a mask outside work	In which situations did you wear a mask?	1 Very exaggerated/much too long; 2 a bit exaggerated/a bit too long; 3 just perfect; 4 a bit weak/a bit too short; 5 much too weak/much too short	None = False; All others = True	Baseline
Support for stronger public restrictions	Do you support the national measures and it's duration or what is your opinion about it?	1 Very exaggerated/much too long; 2 a bit exaggerated/a bit too long; 3 just perfect; 4 a bit weak/a bit too short; 5 much too weak/much too short	Answers 1-3 = False; Answers 4-5 = True	Baseline
HCW specifics				
Job: nurse	What is your job description?	Nurse; Medical doctors assistant; Physician; Secretary; Scientist; Physiotherapist; Ergotherapist; Logopedist; Social service; Dietician; Medical technical radiology assistant (MTRA); Housekeeping; Volunteer; Gastronomy/Hotel services; Kitchen; Technical service; Administration (Human resources, management); Informatics; Laboratory; other	Nurse = True; All others = False	Baseline
Job: physician	What is your job description?	Physician; Secretary; Scientist; Physiotherapist; Ergotherapist; Logopedist; Social service; Dietician; Medical technical radiology assistant (MTRA); Housekeeping; Volunteer; Gastronomy/Hotel services; Kitchen; Technical service; Administration (Human resources, management); Informatics; Laboratory; other	Physician = True; All others = False	Baseline
Full-time job (> 80%)	What is your work percentage?	Percentage (xxx%)	>80% = True; 0-80% = False	Baseline
Involved in AGP	In which aerosol-generating procedures have you been involved since March 2020?	None; In-/Extubation; Bronchoscopy; Reanimation; Tracheal secretion suction; Non-invasive ventilation; Gastroscopy; Transesophageal echocardiography; other	None = False; All others = True	Follow-up
Working in intensive care	In which department do you work?	List of departments	Intensive care unit = True, All others = False	Baseline
Behaviour at work				
Hygiene knowledge	Which are standard hygiene measures?	Cough etiquette; Physical distancing*; Handhygiene; Vaccines; Wearing a gown when in contact with body fluids; Wearing surgical mask when having a cold (* incorrect answer, all others are correct)	Number of correct answers ≥ 3 = True; < 3 = False	Baseline
Regular meals in staff restaurant	How often do you go to the staff restaurant or the cafeteria?	None is available; Never; Less than weekly; Weekly, but not daily; Daily; More than daily; No restaurant or cafeteria visits	None, never or less than weekly; weekly or more	Baseline
Handwashing more frequent	Do you wash your hands more frequently in the current COVID-19 situation compared to before?	True; False	True; False	Baseline
Use of personal protective equipment				
Always used goggles ¹	In what situations (during COVID-19 patient contact) did you use goggles?	1 Never; 2 Occasionally; 3 Only if a contact with body fluids or secretions is expected; 4 Always	Answer 4 = True; Answers 1-3 = False	Follow-up
Always used gloves/gown ¹	In what situations (during COVID-19 patient contact) did you use gloves? In what situations (during COVID-19 patient contact) did you use a gown?	1 Never; 2 Occasionally; 3 Only if a contact with body fluids or secretions is expected; 4 Always	Answer 4 (for both gloves and gown) = True; All others = False	Follow-up
Mostly used FFP2 ¹	What mask do you usually wear in contact with COVID-19 patients, when non-aerosol generating procedures are done? Please consider the whole time since the start of the pandemic.	1 Always surgical mask; 2 mostly surgical mask; 3 Both surgical mask and FFP2; 4 Mostly FFP2; 5 Always FFP2	Answers 1-3 = False; Answers 4-5 = True	Follow-up
Always used FFP2 during AGP	What mask do you usually wear in contact with COVID-19 patients, when aerosol generating procedures are performed? Please consider the whole time since the start of the pandemic.	1 Always surgical mask; 2 Mostly surgical mask; 3 Both surgical mask and FFP2; 4 Mostly FFP2; 5 Always FFP2	Answers 1-4 = False; Answer 5 = True	Follow-up

¹ In contact with COVID-19 patients outside of AGPs

Abbreviations: AU – Austria. GE – Germany. HCW – Health care worker. COPD – Chronic obstructive pulmonary disease. AGP – Aerosol-generating procedure. FFP2 – Filtering facepiece class 2.

Table S2. Cox regression analysis (full model and sensitivity analyses) with outcome “SARS-CoV-2- positive nasopharyngeal PCR/rapid antigen test”.

Participants; Events	Full model ¹		Sensitivity analysis 1 ¹		Sensitivity analysis 2		Sensitivity analysis 3 ¹	
	HR (95% CI)	p	HR (95% CI)	p	HR (95% CI)	p	HR (95% CI)	p
	n=3'259; n=433		n=2'783; n=250		n=3'259; n=433		n=3'067; n=203	
			Excluding HCW with positive households		Treating cantons and institutions as fixed effects		Restricting analysis to events after Dec 1st 2020	
Risk or protection factor	HR (95% CI)	p	HR (95% CI)	p	HR (95% CI)	p	HR (95% CI)	p
Exposure to COVID-19								
Patient exposure (<i>time dependent</i>)	2.24 (1.63-3.10)	<0.001	2.84 (2.05-3.93)	<0.001	2.11 (1.67-2.66)	<0.001	2.07 (1.55-2.76)	<0.001
Coworker exposure (<i>time dependent</i>)	1.58 (1.26-1.99)	<0.001	1.98 (1.58-2.48)	<0.001	1.56 (1.20-2.04)	0.001	1.48 (1.13-1.94)	0.004
Household exposure (<i>time dependent</i>)	10.06 (7.48-13.51)	<0.001	NA		9.64 (7.66-12.12)	<0.001	14.21 (9.66-20.91)	<0.001
Other exposure (<i>time dependent</i>)	1.51 (1.31-1.73)	<0.001	1.70 (1.45-2.00)	<0.001	1.48 (1.20-1.83)	<0.001	1.64 (1.22-2.22)	0.001
Number of negative swabs (<i>time dependent</i>)	0.91 (0.84-0.98)	0.008	0.94 (0.83-1.05)	0.277	0.91 (0.81-1.02)	0.120	0.95 (0.87-1.04)	0.243
Sociodemographic data								
Age (per 10 years)	1.06 (1.00-1.13)	0.046	0.97 (0.89-1.05)	0.429	1.08 (0.98-1.18)	0.133	0.96 (0.88-1.05)	0.400
Gender: male	0.78 (0.61-1.01)	0.058	0.50 (0.38-0.66)	<0.001	0.82 (0.60-1.11)	0.191	0.64 (0.36-1.15)	0.137
Living in Germany or Austria	1.10 (0.71-1.70)	0.667	0.92 (0.40-2.11)	0.837	NA		0.69 (0.35-1.38)	0.295
Child in household	1.13 (0.93-1.38)	0.228	0.69 (0.55-0.87)	0.002	1.10 (0.86-1.40)	0.458	1.70 (1.25-2.29)	0.001
Medical conditions								
Comorbidity	0.99 (0.81-1.20)	0.895	1.01 (0.76-1.33)	0.965	1.00 (0.82-1.23)	0.982	0.99 (0.72-1.34)	0.923
Active smoking	0.65 (0.49-0.86)	0.003	0.59 (0.41-0.85)	0.005	0.66 (0.50-0.89)	0.006	0.55 (0.29-1.06)	0.074
Pregnancy during study	1.37 (1.05-1.77)	0.019	1.27 (0.68-2.36)	0.458	1.37 (0.87-2.16)	0.172	2.03 (1.33-3.09)	0.001
Behaviour outside of work								
Prophylactic home remedies	1.08 (0.82-1.42)	0.598	1.00 (0.76-1.30)	0.974	1.07 (0.82-1.40)	0.599	1.30 (0.94-1.78)	0.107
Social leisure activities	0.93 (0.79-1.09)	0.357	0.79 (0.67-0.94)	0.007	0.91 (0.74-1.10)	0.324	1.04 (0.79-1.37)	0.781
Wearing a mask outside work	0.74 (0.56-0.99)	0.040	0.69 (0.43-1.11)	0.127	0.75 (0.58-0.97)	0.026	0.84 (0.64-1.11)	0.222
Support for stronger public restrictions	0.88 (0.73-1.05)	0.161	0.92 (0.67-1.27)	0.619	0.96 (0.73-1.25)	0.749	1.11 (0.76-1.62)	0.593
HCW specifics								
Job: nurse	1.20 (1.03-1.39)	0.020	1.43 (1.12-1.83)	0.004	1.15 (0.89-1.49)	0.276	1.24 (0.94-1.63)	0.127
Job: physician	0.87 (0.65-1.15)	0.328	1.10 (0.73-1.67)	0.650	0.82 (0.58-1.15)	0.256	0.90 (0.55-1.46)	0.669
Full-time job (> 80%)	1.10 (0.94-1.27)	0.230	1.10 (0.90-1.34)	0.347	1.08 (0.85-1.39)	0.521	1.51 (1.05-2.16)	0.026
Involved in AGP	1.18 (0.96-1.45)	0.114	1.21 (0.90-1.62)	0.217	1.13 (0.91-1.40)	0.266	1.31 (1.04-1.67)	0.024
Working in intensive care	0.77 (0.53-1.12)	0.170	0.63 (0.42-0.94)	0.025	0.74 (0.50-1.10)	0.142	1.02 (0.63-1.66)	0.924
Behaviour at work								
Hygiene knowledge	1.09 (0.91-1.32)	0.342	1.01 (0.71-1.44)	0.964	1.00 (0.75-1.34)	0.976	1.30 (0.96-1.75)	0.089
Regular meals in staff restaurant	1.16 (1.00-1.36)	0.058	1.13 (0.95-1.34)	0.173	1.15 (0.93-1.43)	0.199	1.16 (0.84-1.59)	0.373

Use of personal protective equipment

Always used goggles ²	0·81 (0·58-1·12)	0·201	0·81 (0·53-1·25)	0·342	0·88 (0·67-1·16)	0·375	0·66 (0·40-1·08)	0·100
Always used gloves/gown ²	1·11 (0·71-1·73)	0·644	1·11 (0·69-1·80)	0·664	1·07 (0·81-1·41)	0·636	1·16 (0·69-1·96)	0·572
Mostly used FFP2²	0·80 (0·64-1·00)	0·052	0·82 (0·54-1·25)	0·360	0·88 (0·66-1·18)	0·402	0·73 (0·55-0·97)	0·029

¹ Institutions and living place (canton) included in model as random cluster term

² Germany/Austria treated like separate cantons and included as fixed effects

³ In contact with COVID-19 patients outside of AGPs

Abbreviations: PCR – Polymerase chain reaction. HCW – Health care worker. AGP – Aerosol-generating procedure. FFP2 – Filtering facepiece class 2. NA – Not applicable

Table S3. Results of multivariable logistic regression analysis (full model and sensitivity analyses) regarding outcome “SARS-CoV-2 seroconversion”.

Risk or protection factor	Full model ¹		Sensitivity analysis 1 ¹		Sensitivity analysis 2	
	Participants; Events	n=2916; n=511	Excluding HCW with positive households	n=2491; n=335	Treating cantons and institutions as fixed effects	n=2916; n=511
	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Sociodemographic data						
Age (per 10 years)	0.96 (0.87 - 1.07)	0.475	0.92 (0.82 - 1.04)	0.177	0.96 (0.86 - 1.06)	0.422
Sex: male	0.86 (0.62 - 1.18)	0.349	0.77 (0.53 - 1.13)	0.176	0.86 (0.62 - 1.19)	0.368
Living in Germany or Austria	0.86 (0.47 - 1.58)	0.628	0.81 (0.42 - 1.55)	0.522	NA	
No of additional persons in households	0.88 (0.79 - 0.98)	0.016	0.86 (0.76 - 0.98)	0.022	0.87 (0.78 - 0.97)	0.013
Child in household	0.97 (0.71 - 1.34)	0.859	0.73 (0.48 - 1.10)	0.132	0.98 (0.71 - 1.35)	0.880
Any positively tested household member	5.01 (3.89 - 6.46)	<0.001	NA		5.11 (3.96 - 6.60)	<0.001
Medical conditions						
Comorbidity	0.93 (0.74 - 1.16)	0.506	0.96 (0.74 - 1.24)	0.732	0.93 (0.74 - 1.16)	0.507
Active smoking	0.57 (0.41 - 0.79)	<0.001	0.56 (0.39 - 0.82)	0.002	0.57 (0.41 - 0.78)	0.001
Pregnancy during study	1.01 (0.56 - 1.82)	0.979	0.91 (0.45 - 1.86)	0.802	0.99 (0.53 - 1.74)	0.969
Behaviour outside of work						
Prophylactic home remedies	1.21 (0.90 - 1.63)	0.205	1.14 (0.81 - 1.62)	0.448	1.21 (0.89 - 1.63)	0.211
Social leisure activities	1.02 (0.82 - 1.27)	0.847	0.92 (0.72 - 1.18)	0.517	1.02 (0.82 - 1.27)	0.863
Wearing a mask outside work	0.96 (0.74 - 1.25)	0.779	0.92 (0.68 - 1.24)	0.588	0.97 (0.74 - 1.27)	0.833
Support for stronger public restrictions	0.85 (0.64 - 1.14)	0.282	0.93 (0.67 - 1.29)	0.657	0.88 (0.65 - 1.17)	0.386
HCW specifics						
Job: nurse	1.55 (1.18 - 2.04)	0.001	1.83 (1.33 - 2.52)	<0.001	1.56 (1.19 - 2.06)	0.002
Job: physician	0.84 (0.59 - 1.21)	0.360	0.88 (0.57 - 1.36)	0.562	0.84 (0.58 - 1.20)	0.337
Full-time job (> 80%)	1.08 (0.83 - 1.42)	0.566	1.11 (0.81 - 1.51)	0.521	1.10 (0.84 - 1.44)	0.509
Involved in AGP	0.96 (0.75 - 1.24)	0.775	1.03 (0.78 - 1.37)	0.820	0.95 (0.74 - 1.21)	0.659
Working in ICU	0.43 (0.27 - 0.68)	<0.001	0.40 (0.22 - 0.71)	0.002	0.43 (0.27 - 0.68)	<0.001
Frequency of contacts with COVID-19 patients	1.51 (1.38 - 1.66)	<0.001	1.50 (1.35 - 1.68)	<0.001	1.51 (1.38 - 1.67)	<0.001
Frequency of contacts with coworkers with COVID-19	1.22 (1.10 - 1.36)	<0.001	1.18 (1.05 - 1.33)	0.007	1.22 (1.10 - 1.36)	<0.001
Behaviour at work						
Hygiene knowledge	1.06 (0.77 - 1.46)	0.698	1.12 (0.76 - 1.65)	0.562	1.03 (0.75 - 1.44)	0.834
Regular meals in staff restaurant	1.12 (0.88 - 1.41)	0.363	1.06 (0.81 - 1.40)	0.657	1.12 (0.88 - 1.42)	0.362
Use of personal protective equipment						
Always used goggles ²	0.91 (0.68 - 1.22)	0.521	0.99 (0.71 - 1.39)	0.953	0.94 (0.70 - 1.26)	0.677

Always used gloves/gown ²	0·81 (0·59 - 1·11)	0·182	0·79 (0·55 - 1·13)	0·197	0·80 (0·58 - 1·10)	0·170
Mostly used FFP2²	0·73 (0·53 - 1·00)	0·053	0·69 (0·48 - 0·99)	0·046	0·76 (0·55 - 1·04)	0·088

¹Institutions and living place (canton) included in model as random cluster term. For Sensitivity Analysis 2, the effects of these factors are not shown in the table.

²In contact with COVID-19 patients outside of AGPs

Abbreviations: HCW – Health care worker. AGP – Aerosol-generating procedure. FFP2 – Filtering facepiece class 2. NA – Not applicable.

Table S4. Subgroup analysis of HCW with frequent COVID-19 exposure vs. HCW with less frequent COVID-19 exposure; A) Cox regression (outcome SARS-CoV-2 positive swab); B) multivariable logistic regression (outcome SARS-CoV-2 seroconversion).

A) Cox Regression

Risk or protection factor	Participants; Events	1–20 COVID-19 patient contacts		>20 COVID-19 patient contacts	
		n = 1292; n = 149		n = 1120; n = 215	
		HR	p	HR	p
Exposure to COVID-19					
Patient exposure (<i>time dependent</i>)		2.09 (1.66-2.62)	<0.001	1.52 (0.93-2.49)	0.099
Coworker exposure (<i>time dependent</i>)		1.64 (1.09-2.46)	0.018	1.61 (0.99-2.63)	0.055
Household exposure (<i>time dependent</i>)		13.93 (9.64-20.12)	<0.001	6.49 (4.02-10.47)	<0.001
Other exposure (<i>time dependent</i>)		1.71 (1.36-2.14)	<0.001	1.20 (0.86-1.66)	0.286
Number of negative swabs (<i>time dependent</i>)		0.88 (0.76-1.01)	0.060	0.93 (0.83-1.03)	0.143
Sociodemographic data					
Age (per 10 years)		1.10 (0.99-1.22)	0.086	1.05 (0.95-1.15)	0.338
Sex:male		0.53 (0.29-0.98)	0.042	0.86 (0.63-1.17)	0.334
Living in Germany or Austria		1.20 (0.35-4.06)	0.775	1.11 (0.49-2.52)	0.794
Child in household		0.93 (0.65-1.34)	0.710	1.24 (0.84-1.84)	0.283
Medical conditions					
Comorbidity		0.95 (0.71-1.27)	0.705	0.99 (0.72-1.35)	0.938
Active smoking		0.72 (0.38-1.36)	0.307	0.61 (0.32-1.14)	0.120
Pregnancy during study		1.41 (0.87-2.29)	0.165	0.88 (0.40-1.94)	0.745
Behaviour outside of work					
Prophylactic home remedies		1.02 (0.70-1.49)	0.909	1.09 (0.76-1.58)	0.640
Social leisure activities		1.03 (0.76-1.41)	0.836	0.91 (0.73-1.12)	0.373
Wearing a mask outside work		0.70 (0.47-1.04)	0.078	0.82 (0.61-1.11)	0.193
Support for stronger public restrictions		0.91 (0.57-1.48)	0.715	0.81 (0.65-1.02)	0.076
HCW specifics					
Job: nurse		1.20 (0.74-1.94)	0.450	1.77 (1.30-2.41)	<0.001
Job: physician		0.53 (0.30-0.94)	0.029	1.48 (0.91-2.39)	0.111
Full-time job (> 80%)		1.10 (0.79-1.53)	0.561	1.09 (0.85-1.39)	0.504
Involved in AGP		1.08 (0.88-1.34)	0.453	1.12 (0.76-1.65)	0.566
Working in intensive care		0.64 (0.35-1.17)	0.149	0.67 (0.45-0.99)	0.046
Behaviour at work					
Hygiene knowledge		1.12 (0.80-1.57)	0.521	1.07 (0.78-1.46)	0.691
Regular meals in staff restaurant		1.28 (0.89-1.84)	0.186	1.02 (0.80-1.30)	0.879
Use of personal protective equipment					
Always goggles ¹		0.76 (0.45-1.29)	0.309	0.93 (0.68-1.27)	0.664

Always gloves/gown ¹	0.82 (0.42-1.61)	0.562	1.28 (0.92-1.78)	0.147
Mostly using FFP2 mask¹	1.06 (0.70-1.63)	0.774	0.66 (0.54-0.81)	<0.001

B) Multivariable Logistic Regression

Risk or protection factor	Participants; Events	1–20 COVID-19 patient contacts		> 20 COVID-19 patient contacts	
		n = 1156; n = 164		n = 1019; n = 272	
		OR (95% CI)	p	OR (95% CI)	p
Exposure to COVID-19					
Frequency of contacts to coworkers with COVID-19		1.22 (1.02 - 1.45)	0.032	1.23 (1.04 - 1.45)	0.016
Any positively tested household member		3.97 (2.60 - 6.06)	<0.001	5.28 (3.59 - 7.78)	<0.001
Sociodemographic data					
Age (per 10 years)		1.03 (0.86 - 1.22)	0.776	0.91 (0.78 - 1.06)	0.233
Sex:male		0.66 (0.37 - 1.20)	0.175	0.95 (0.62 - 1.47)	0.822
Living in Germany or Austria		1.28 (0.51 - 3.22)	0.606	0.74 (0.35 - 1.57)	0.427
Number of additional persons in household		0.94 (0.78 - 1.14)	0.536	0.86 (0.74 - 1.01)	0.066
Child in household		0.81 (0.46 - 1.41)	0.450	1.08 (0.67 - 1.72)	0.759
Medical conditions					
Comorbidity		0.76 (0.52 - 1.13)	0.173	1.02 (0.74 - 1.42)	0.890
Active smoking		0.76 (0.46 - 1.27)	0.296	0.49 (0.30 - 0.80)	0.004
Pregnancy during study		1.13 (0.47 - 2.75)	0.781	0.79 (0.28 - 2.24)	0.657
Behaviour outside of work					
Prophylactic home remedies		1.29 (0.80 - 2.10)	0.298	1.11 (0.69 - 1.80)	0.665
Social leisure activities		1.02 (0.71 - 1.46)	0.931	1.10 (0.80 - 1.52)	0.545
Wearing a mask outside work		0.82 (0.51 - 1.29)	0.388	1.18 (0.80 - 1.74)	0.404
Support for stronger public restrictions		1.01 (0.62 - 1.66)	0.970	0.83 (0.54 - 1.27)	0.395
HCW specifics					
Job: nurse		1.39 (0.89 - 2.17)	0.151	2.22 (1.40 - 3.52)	<0.001
Job: physician		0.50 (0.26 - 0.96)	0.038	1.33 (0.78 - 2.28)	0.298
Full-time job (> 80%)		1.31 (0.82 - 2.08)	0.254	0.97 (0.64 - 1.48)	0.901
Involved in AGP		1.35 (0.90 - 2.03)	0.144	0.82 (0.58 - 1.15)	0.251
Working in intensive care		0.33 (0.11 - 1.00)	0.049	0.51 (0.30 - 0.87)	0.015
Behaviour at work					
Hygiene knowledge		0.99 (0.59 - 1.65)	0.968	1.12 (0.65 - 1.93)	0.691
Regular meals in staff restaurant		1.53 (1.03 - 2.28)	0.037	1.03 (0.72 - 1.47)	0.886
Use of personal protective equipment					
Always goggles ¹		0.90 (0.54 - 1.51)	0.703	1.06 (0.72 - 1.58)	0.755
Always gloves/gown ¹		0.60 (0.35 - 1.03)	0.065	0.98 (0.64 - 1.49)	0.915
Mostly using FFP2 mask¹		0.75 (0.42 - 1.32)	0.320	0.64 (0.42 - 0.97)	0.036

¹In contact with COVID-19 patients outside of AGPs

Abbreviations: HCW – Health care worker. AGP – Aerosol-generating procedure. FFP2 – Filtering facepiece class 2. NA – not applicable

Table S5. Subgroup analysis of HCW performing AGP: A) Cox regression (outcome SARS-CoV-2 positive swab); B) multivariable logistic regression (outcome SARS-CoV-2 seroconversion).

A) Cox Regression	Participants; Events	Participants involved in AGP	
		n = 1204; n = 188	
Risk or protection factor		HR	p
Exposure to COVID-19			
Patient exposure (<i>time dependent</i>)		1.50 (0.98-2.32)	0.065
Coworker exposure (<i>time dependent</i>)		1.72 (1.24-2.37)	0.001
Household exposure (<i>time dependent</i>)		9.75 (6.28-15.14)	<0.001
Other exposure (<i>time dependent</i>)		1.42 (1.07-1.88)	0.015
Number of negative swabs (<i>time dependent</i>)		0.88 (0.75-1.03)	0.108
Sociodemographic data			
Age (per 10 years)		1.02 (0.88-1.19)	0.776
Sex:male		0.87 (0.57-1.33)	0.531
Living in Germany or Austria		1.46 (0.79-2.67)	0.224
Child in household		1.15 (0.88-1.50)	0.297
Medical conditions			
Comorbidity		1.13 (0.85-1.49)	0.402
Active smoking		0.65 (0.37-1.13)	0.129
Pregnancy during study		0.56 (0.13-2.38)	0.433
Behaviour outside of work			
Prophylactic home remedies		1.11 (0.67-1.83)	0.685
Social leisure activities		0.98 (0.77-1.26)	0.900
Wearing a mask outside work		0.75 (0.51-1.11)	0.157
Support for stronger public restrictions		0.91 (0.69-1.20)	0.512
HCW specifics			
Job: nurse		1.17 (0.87-1.58)	0.309
Job: physician		1.12 (0.64-1.96)	0.697
Full-time job (> 80%)		0.95 (0.65-1.37)	0.767
Working in intensive care		0.77 (0.48-1.24)	0.277
Behaviour at work			
Hygiene knowledge		1.72 (1.03-2.88)	0.039
Regular meals in staff restaurant		1.00 (0.67-1.47)	0.981
Use of personal protective equipment			
Always goggles ¹		0.78 (0.59-1.03)	0.075
Always gloves/gown ¹		1.11 (0.67-1.82)	0.686
Mostly using FFP2 mask¹		0.71 (0.50-1.01)	0.059

Always using FFP2 mask during AGP	1.08 (0.71-1.64)	0.726
B) Multivariable Logistic Regression		
	Participants involved in AGP	
Participants; Events	n = 1087; n = 216	
Risk or protection factor	OR (95% CI)	p
Exposure to COVID-19		
Frequency of contacts to COVID-19 patients	1.44 (1.22 - 1.70)	<0.001
Frequency of contacts to coworkers with COVID-19	1.27 (1.06 - 1.51)	0.008
Any positively tested household member	4.69 (3.14 - 7.00)	<0.001
Sociodemographic data		
Age (per 10 years)	1.01 (0.85 - 1.19)	0.951
Sex:male	0.99 (0.64 - 1.53)	0.953
Living in Germany or Austria	1.38 (0.65 - 2.91)	0.404
Number of additional persons in households	0.89 (0.75 - 1.06)	0.185
Child in household	0.90 (0.53 - 1.52)	0.693
Medical conditions		
Comorbidity	1.08 (0.76 - 1.52)	0.667
Active smoking	0.46 (0.28 - 0.78)	0.004
Pregnancy during study	1.25 (0.39 - 3.95)	0.708
Behaviour outside of work		
Prophylactic home remedies	1.41 (0.88 - 2.25)	0.149
Social leisure activities	1.05 (0.75 - 1.47)	0.786
Wearing a mask outside work	0.84 (0.56 - 1.26)	0.404
Support for stronger public restrictions	1.11 (0.73 - 1.69)	0.633
HCW specifics		
Job: nurse	1.28 (0.78 - 2.09)	0.330
Job: physician	0.97 (0.55 - 1.73)	0.930
Full-time job (> 80%)	0.95 (0.62 - 1.46)	0.822
Working in intensive care	0.44 (0.27 - 0.74)	0.002
Behaviour at work		
Hygiene knowledge	1.81 (0.94 - 3.50)	0.077
Regular meals in staff restaurant	1.09 (0.75 - 1.59)	0.645
Use of personal protective equipment		
Always used goggles ¹	0.89 (0.59 - 1.32)	0.552
Always used gloves/gown ¹	0.86 (0.56 - 1.34)	0.518
Mostly used FFP2 mask¹	0.66 (0.43 - 1.01)	0.055
Always used FFP2 mask during AGP	0.89 (0.62 - 1.28)	0.535

¹ In contact with COVID-19 patients outside of AGPs

Abbreviations: HCW – Health care worker. AGP – Aerosol-generating procedure. FFP2 – Filtering facepiece class 2. NA – not applicable