

Supplementary Table 1. Sociodemographics, clinical characteristics and severity scoring for COVID-19 patients

Sociodemographics and Clinical Characteristics	Number of patients n (%)	
Sex		
Male	27 (54)	
Female	23 (46)	
Age (years)		
<35	12 (24)	
35-60	21 (42)	
>60	17 (34)	
Hospitalization		
Home	25 (50)	
Admission	25 (50)	
Disease severity		
0	0 (0)	
1	16 (32)	
2	17 (34)	
3	14 (28)	
4	3 (6)	
Serological positive (WANTAI)		
	49 (98)	

Score	Short name	Criteria
0	Asymptomatic	SARS-CoV-2 proven infection without developing symptoms of COVID-19
1	Mild	Symptoms of COVID-19 without evidence of viral pneumonia or hypoxia
2	Moderate	Clinical signs of pneumonia but no signs of severe pneumonia, including $\text{SpO}_2 \geq 90\%$ on room air
3	Severe	Clinical signs of severe pneumonia: respiratory rate > 30 breaths/min, $\text{SpO}_2 < 90\%$ on room air
4	Critical	ICU admission (ARDS, Sepsis, Septic shock)

Supplementary Table 2. Sequence identity matrices for the ectodomains of all hCoV S proteins

S

	SARS 2 S	SARS S	MERS S	OC43 S	HKU1 S	229E S	NL63 S
SARS 2 S	100%						
SARS S	75,4%	100%					
MERS S	30,9%	31,4%	100%				
OC43 S	30,0%	30,6%	31,7%	100%			
HKU1 S	30,3%	31,1%	32,2%	65,9%	100%		
229E S	27,4%	27,2%	28,4%	27,7%	28,8%	100%	
NL63 S	26,2%	24,8%	27,1%	28,1%	28,3%	65,1%	100%

S1

	SARS 2 S1	SARS S1	MERS S1	OC43 S1	HKU1 S1	229E S1	NL63 S1
SARS 2 S1	100%						
SARS S1	66,8%	100%					
MERS S1	21,0%	22,1%	100%				
OC43 S1	25,1%	24,7%	23,4%	100%			
HKU1 S1	23,9%	25,2%	26,4%	61,5%	100%		
229E S1	17,5%	16,9%	18,1%	16,5%	19,4%	100%	
NL63 S1	16,4%	14,6%	15,1%	16,9%	17,5%	51,7%	100%

S2

	SARS 2 S2	SARS S2	MERS S2	OC43 S2	HKU1 S2	229E S2	NL63 S2
SARS 2 S2	100%						
SARS S2	87,6%	100%					
MERS S2	44,5%	43,4%	100%				
OC43 S2	41,8%	40,9%	45,2%	100%			
HKU1 S2	41,3%	40,6%	43,5%	71,7%	100%		
229E S2	38,0%	37,7%	38,0%	35,0%	35,0%	100%	
NL63 S2	38,2%	35,9%	36,9%	36,6%	35,4%	77,7%	100%


RBD

	SARS 2 RBD	SARS RBD	MERS RBD	OC43 RBD	HKU1 RBD	229E RBD	NL63 RBD
SARS 2 RBD	100%						
SARS RBD	73,4%	100%					
MERS RBD	19,2%	21,3%	100%				
OC43 RBD	23,1%	24,6%	27,1%	100%			
HKU1 RBD	19,7%	23,4%	27,5%	60,0%	100%		
229E RBD	16,7%	12,8%	23,6%	17,1%	21,1%	100%	
NL63 RBD	12,5%	17,0%	20,0%	17,1%	15,6%	57,3%	100%


Sequence identity matrices were composed of all coronavirus spike proteins in this study. All sequences comprise only the truncated ectodomain of each spike as was used to generate the recombinant proteins. S1, S2 and RBD were defined as noted in the corresponding Genbank sequences (see methods). Multiple sequence alignments were performed and sequence identities calculated using Clustal Omega 1.2.4.

Supplementary Fig. 1


a



b




Supplementary Fig. 2




Supplementary Fig. 3


a




b




Supplementary Fig. 4



Supplementary Fig. 5




Supplementary Fig. 6




Supplementary Fig. 7


a



b




c




Supplementary Fig. 8


a



b




c




Supplementary Fig. 9

a



b



Supplementary Fig. 10

