

Insecticide resistance outpaces behavioural adaptation, as a response to Long-Lasting Insecticidal Net distribution, in malaria vectors in Burkina Faso.

Antoine Sanou^{1, 2*}; Luca Nelli¹; W. Moussa Guelbéogo²; Fatoumata Cissé²; Michel M. Tapsoba²; Pierre Ouédraogo²; N'falé Sagnon²; Hilary Ranson³; Jason Matthiopoulos¹; Heather M. Ferguson¹.

Supplementary Table 1: Significance of explanatory variables included in the model for assessing variation in *An. gambiae* s.l. mortality rates. Here, df is the degree of freedom, Chi-sq (χ^2) represents the values of Likelihood Ratio Test. and p indicates the p-values associated with each term. “Season” is a smoothing function on days from 1 - 365 describing a year of collection for assessing the seasonality in the proportion. “Longer-term” a discrete variable from first to the last day of collection (798) describing the long-term trend in the proportion.

Explanatory variable	0.05% deltamethrin (Model 1)			0.25% deltamethrin (Model 2)			0.5% deltamethrin (Model 3)			0.75% deltamethrin (Model 4)		
	Chi-sq	df	p	Chi-sq	df	p	Chi-sq	df	p	Chi-sq	df	p
Season	15.1	3.95 ^a	0.003 ^b	1.54	0.7 ^a	0.12	0.0	0.0	0.31	0.0	0.0	0.13
Longer-term	20.91	1	0.00 ^{bn}	3.726	1	0.05 ⁿ	0.11	1	0.74	11.25	1	0.00 ^{bn}
Village	29.61	8	0.00 ^b	17.44	8	0.00 ^b	12.72	7	0.00 ^b	16.81	7	0.019 ^b

^a indicates that the correspondent “df” represents the effective degree of freedom (edf); ^b indicates the significant terms in the model and ⁿ indicates negative therefore a long-term reduction in the mortality rate; P = 0.00 indicate p<0.001;

Supplementary Table 2: Output of the different models on proportion of *An. coluzzii*, the outdoor biting, the median biting time and the outdoor resting. Here, df is the degree of freedom, Chi-sq (χ^2) represents the values of Likelihood Ratio Test. and p indicates the p-values associated with each term. “Season” is a smoothing function on days from 1 - 365 describing a year of collection, for assessing the seasonality in the proportion. “Longer-term” a discrete variable from first to the last day of collection (798) describing the long-term trend in the proportion.

Explanatory variable	Proportion of Anopheles coluzzii (Model 5)			Human biting rate (Model 6)			Proportion of outdoor biting (Model 7)			Median biting time (Model 10)		
	Chi-sq	df	p	Chi-sq	df	p	Chi-sq	df	p	F	df	p
Season	68.35	5.22 ^a	0.04	6.63	1	0.01*	3.06	1.55 ^a	0.19	89.66	1.92 ^a	0.00
Longer-term	32.78	1	0.00 ⁿ	1165	6.84 ^a	0.0069*	1.31	1	0.25	0.127	1	0.72
Village	n/a	n/a	n/a	230.54	11	<0.0001*	13.70	11	0.25	2.32	11	0.00
Location	n/a	n/a	n/a	21.28	1	<0.0001*	n/a	n/a	n/a	0.221	1	0.64
Location: Village	24.78	11	0.01	14.93	11	0.185	n/a	n/a	n/a	0.61	11	0.82
Humidity	0.04	1	0.84	0.004	1	0.95	2.76	1	0.09	0.21	1	0.65
Temperature	1.53	1	0.22	0.15	1	0.7	0.49	1	0.48	0.346	1	0.55
Species	n/a	n/a	n/a	n/a	n/a	n/a	6.82	1	0.009	23.47	1	0.00

^a indicates that the correspondent “df” represents the effective degree of freedom (edf); ^b indicates the significant terms in the model and ⁿ indicates negative therefore a long-term reduction in the mortality rate; P = 0.00 indicate p<0.001;

Supplementary Table 3 Output of the Model 14 and 15 on respectively the proportion of mosquito when people are indoor (P_{fi}) and human exposure to mosquito bite indoor (π_i). Here, df is the degree of freedom, Chi-sq (χ^2) represents the values of Likelihood Ratio Test. and p indicates the p-values associated with each term. “Season” is a seasonal spline from 1 - 365 describing a year of collection, and “longer-term” a discrete variable from first to the last day of collection (798) describing a linear term representing longer-term linear change across the study period.

Explanatory variable	Proportion of mosquito when people are indoor (P_{fi}) (Model 14)			Human exposure to mosquito bite indoor (π_i) (Model 15)		
	Chi-sq	df	p	F	df	p
Season	0.01	0.02 ^a	0.39	1.42	0.86 ^a	0.11
Longer-term	9.86	1	0.00 ⁿ	13.81	1	0.00 ⁿ
Village	50.16	11	0	41.51	11	0
Humidity	0.34	1	0.56	0.29	1	0.59
Temperature	1.28	1	0.126	3.58	1	0.06

^a indicates that the correspondent “df” represents the effective degree of freedom (edf); ^b indicates the significant terms in the model; ⁿ and indicates negative therefore a long-term reduction in the mortality rate; P = 0.00 indicate p<0.001;

Supplementary Table 4: Significance of explanatory variables included in the Model 13 for *Plasmodium falciparum* sporozoite rates. Here, df is the degree of freedom and Chi.sq (χ^2) represents the values of Likelihood Ratio Test; “Season” is a smoothing function on days from 1 - 365 describing a year of collection, for assessing the seasonality in the proportion. “Longer-term” a discrete variable from first to the last day of collection (798), describing the long-term trend in the proportion. The temperature and relative humidity were obtained by averaging the records over the course of the collection night.

Explanatory variable	Chi.sq	df	p-values
Season	3.175	1.3 ^a	0.036*
Humidity	0.54	1	0.46
Location	0.02.49	1	0.12
Location: Species	1.02	1	0.31
Longer-term	6.26	1	0.01*
Species	0.05	1	0.82
Temperature	2.41	1	0.12
Village	34.61	11	0.0002*
Village : Species	3.35	11	0.95

* indicates the significant terms with p < 0.05 and ^a indicate the given df represent the estimate degree of freedom (edf = degree of wigginess) from smoother term.

Supplementary Table 5: Global Positioning System (GPS) coordinates and altitudes of the 12 study villages.

Village	North	East	Altitude
Dangouindougou	10° 11.970'	005° 0.763'	294
Gouera	10° 24.233'	004° 52.418'	306
Nianiagara	10° 33 42.4'	004° 41.801'	324
Nofesso	10° 08.226'	004° 46.972'	300
Ouangolodougou	10° 04.234'	004° 48.329'	307
Sitiena	10° 36.244'	004° 40.244'	286
Tengrela	10° 39.113'	004° 49.465'	285
Tiefora	10° 37.447'	004° 33.201'	302
Timperba	10° 09.897'	004° 53.983'	318
Tondoura	10° 11.792'	004° 46.311'	299
Toumousseni	10° 37.273'	004° 54.893'	274
Yendere	10° 12.473'	004° 58.729'	315

Supplementary Table 6: Number of females *Anopheles gambiae* s.l. exposed to different concentrations of deltamethrin and untreated paper (control) in bioassays between September 2016, December 2018 and summed across replicates. Numbers inside brackets are the total number of bioassay replicates conducted. "n/a" indicates no bioassays done at this level.

Village	Concentrations of deltamethrin used					Total
	0	0.05	0.25	0.5	0.75	
Dangouindougou	80 (3)	200(8)	50(2)	51(2)	49(2)	430
Gouera	150 (6)	54(2)	148(6)	125(5)	72(3)	549
Nianiagara	n/a	71(3)	n/a	n/a	n/a	91
Sitiena	315(13)	382(15)	252(10)	128(5)	27(1)	1,104
Tengrela	757(31)	411(16)	502(20)	503(20)	421(17)	2,594
Tiefora	528(21)	492(20)	488(20)	434(17)	231 (9)	2,173
Tondoura	282(11)	300(12)	240(10)	147(6)	123(5)	1,092
Toumousseni	356(14)	356(14)	315(13)	292(12)	250(10)	1,569
Yendere	205(8)	253(10)	167(7)	186(7)	71(3)	857
Total	2,693	2,519	2,162	1,866	1,244	10,484

Supplementary Table 7: Number of females *Anopheles gambiae s.l.* exposed at different seasons to deltamethrin and untreated paper (control) in bioassays between September 2016 and December 2018, summed across replicates. Numbers inside brackets are the total number bioassay of replicates conducted. “n/a” indicates no bioassays done at this level.

Village	Seasons					Total
	wet-16	dry-17	wet-17	dry-18	wet-18	
Dangouindougou	n/a	n/a	98(4)	n/a	332(13)	430
Gouera	n/a	n/a	549(22)	n/a	n/a	549
Nianiagara	n/a	n/a	91(3)	n/a	n/a	91
Sitiena	232(9)	n/a	465(17)	168(7)	239(10)	1,104
Tengrela	307(12)	447(18)	663(27)	173(7)	1,004(40)	2,594
Tiefora	116(5)	426(17)	527(21)	469(19)	635(25)	2,173
Tondoura	n/a	n/a	419(17)	n/a	673(27)	1,092
Toumousseni	464(17)	n/a	371(15)	n/a	734(29)	1,569
Yendere	249(10)	n/a	77(3)	286(11)	270(11)	857
Total	1,368	873	3,260	1,096	3,887	10,484

Supplementary Table 8: Maximal models used for modelling seasonality including the primary response variable, explanatory variables and statistical distribution used. Hyphens (-) indicates no random effect used. Physiology means the abdominal status. “Longer_term” a discrete variable from first to the last day of collection (798) indicates the longer-term trend over the collection periods fitted as a linear term. The seasonality term was fitted using a non-linear smoothing function (spline $t_2(\text{Season}, \text{bs}=\text{cc})$) on days as a period of 365-days. nHour and the $l(\text{nHour}^2)$ respectively represent here hours as discrete variables from 1 being the first hour of collection (7pm-8pm) to the last hour of collection of the night being 11 (5am – 6am) and its quadratic term. Here, “subset of *An. gambiae* s.l.” refers to subset that were individually identified to species levels.

Model	Trait	Response variables	Fixed Effect variables	Random effects	Type of data	Distribution
1	Mortality rate	(Dead/ (Dead +Alive))	Village+ longer_term + $t_2(\text{Season}, \text{bs}=\text{cc})$,	Replicate	Subset of <i>An. gambiae</i> s.l. exposed to 0.05% of deltamethrin	binomial
2	Mortality rate	(Dead/ (Dead +Alive))	Village+ longer_term + $t_2(\text{Season}, \text{bs}=\text{cc})$,	Replicate	Subset of <i>An. gambiae</i> s.l. exposed to 0.25% of deltamethrin	binomial
3	Mortality rate	(Dead/ (Dead +Alive))	Village+ longer_term + $t_2(\text{Season}, \text{bs}=\text{cc})$,	Replicate	Subset of <i>An. gambiae</i> s.l. exposed to 0.5% of deltamethrin	binomial
4	Mortality rate	(Dead/ (Dead +Alive))	Village+ longer_term + $t_2(\text{Season}, \text{bs}=\text{cc})$,	Replicate	Subset of <i>An. gambiae</i> s.l. exposed to 0.75% of deltamethrin	binomial
5	Proportion of <i>An. coluzzii</i>	(<i>An. coluzzii</i> / <i>An. coluzzii</i> + <i>An. gambiae</i>)	Village+ Location+ Temperature+ Humidity +Village : Location + Longer_term + $t_2(\text{Season})$	Compound+ Household	Subset: <i>An. gambiae</i> s.l. lab-processed data	Binomial

Model	Trait	Response variables	Fixed Effect variables	Random effects	Type of data	Distribution
6	Human biting rate (HBR)	Number of <i>An. gambiae</i> s.l.	Village + Location + Temperature + Humidity + Year + Village: Year + $t_2(\text{Season}, \text{bs}=\text{cc})$	Compound+ Household	Host-seeking nightly <i>An. gambiae</i> s.l. data from 6 villages	Negative binomial
7	Outdoor biting proportion	(Outdoor/ Outdoor +Indoor)	Village+ Temperature+ Humidity + Longer_term+ $t_2(\text{Season}, \text{bs}=\text{cc})$,	Compound+ Household	Host-seeking nightly <i>An. gambiae</i> s.l.	Binomial
8	Outdoor biting proportion	(Outdoor/ Outdoor +Indoor)	Village+ Species+ Temperature+ Humidity + Longer_term+ $t_2(\text{Season}, \text{bs}=\text{cc})$,	Compound+ Household	Subset of <i>An. gambiae</i> s.l. identified to species level	Binomial
9	Mean number of bite/person/hours	Hourly count	nHour + I(nHour ²)	-	Host-seeking hourly <i>An. gambiae</i> s.l.	Negative binomial
10	Median biting time	Median of hours	Village+ Location+ Longer_term + Temperature + Humidity + Village : Location + $t_2(\text{Season}, \text{bs}=\text{cc})$	Compound+ Household	Host-seeking hourly <i>An. gambiae</i> s.l.	Poisson
11	Median biting time by species	Median of hours	Species+ Village+ Location+ Longer_term + Temperature + Humidity + $t_2(\text{Season}, \text{bs}=\text{cc})$	Compound+ Household	Subset of <i>An. gambiae</i> s.l. data	Poisson
12	Outdoor resting proportion (female)	(Outdoor/ Outdoor +Indoor)	Longer_term+ Species + Physiology + Species: Physiology + $t_2(\text{Season}, \text{bs}=\text{cc})$	Village + Compound+ Household	Subset: <i>An. gambiae</i> s.l. resting lab-processed data	Binomial
13	Sporozoite rate	(Positive/Positive+ Negative))	Village + Location + Species + Village: Species + Location: Species + Temperature + Humidity + Longer_term + $t_2(\text{cDate}, \text{bs} = \text{cc})$,	Compound+ Household	Subset of <i>An. gambiae</i> s.l. lab-processed from 6 villages	Binomial
14	Proportion of mosquito when people are indoor (P_{fi})	$(I_{10\text{pm} \rightarrow 5\text{am}} + O_{10\text{pm} \rightarrow 5\text{am}}) / (I_{7\text{pm} \rightarrow 6\text{am}} + O_{7\text{pm} \rightarrow 6\text{am}})$	Village+ Temperature + Humidity + Longer_term + $t_2(\text{Season}, \text{bs}=\text{cc})$,	Compound+ Household	Host-seeking nightly <i>An. gambiae</i> s.l.	Binomial
15	Human exposure to mosquito bite indoor (π_i)	$I_{10\text{pm} \rightarrow 5\text{am}} / (I_{10\text{pm} \rightarrow 5\text{am}} + O_{7\text{pm} \rightarrow 10\text{pm}, 5\text{am} \rightarrow 6\text{am}})$	Village+ Temperature+ Humidity + Longer_term + $t_2(\text{Season}, \text{bs}=\text{cc})$,	Compound+ Household	Host-seeking nightly <i>An. gambiae</i> s.l.	Binomial