

Supporting information for

Formation of oxidized gases and secondary organic aerosol from a commercial oxidant-generating electronic air cleaner


Taekyu Joo^a, Jean C. Rivera-Rios^b, Daniel Alvarado-Velez^b, Sabrina Westgate^b, Nga Lee Ng^{a,b,c}

5 *^aSchool of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, Georgia 30332, United States*


^bSchool of Chemical and Biomolecular Engineering, Georgia Institute of Technology, Atlanta, Georgia 30332, United States

^cSchool of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, Georgia 30332, United States


10




15 Figure S1. Size distribution of (a) particle number concentration ($\# \text{ cm}^{-3}$) and (b) particle volume concentration ($\text{nm}^3 \text{ cm}^{-3}$) from the office measurements. The color scale is in log scale.




20 Figure S2. SMPS and CPC results from an experiment performed in a larger space (laboratory) than the office: (a) time series of number (CPC and SMPS) and volume concentrations (SMPS) with device operation period highlighted in yellow. Size distribution of (b) particle number concentration ($\# \text{cm}^{-3}$) and (c) particle volume concentration ($\text{nm}^3 \text{cm}^{-3}$). The color scale is in log scale.




25 Figure S3. Comparison of AMS total mass concentration (sum of all non-refractory species) and SMPS mass concentration (intercepts set to zero) from the office measurements. The SMPS volume concentration was converted into mass concentration using density of the individual species: 1.75 g/cm^3 for sulfate, nitrate, and ammonium; 1.52 g/cm^3 for chloride; 1.79 g cm^{-3} for organics, estimated based on the elemental ratio of organics.¹ Particle water from AMS was estimated based on the equation proposed in Engelhart et al.²



30 Figure S4. Organic families mass spectra of (a) office background and (b) during device operation.



35 Figure S5. (a) Time series of H:C, O:C, and NO^+/NO_2^+ ratio from the office measurements, with the device operation period highlighted in yellow. (b) $f_{H_2SO_4^+}$ vs. $f_{HSO_3^+}$ during the experiment. $f_{H_2SO_4^+}$ and $f_{HSO_3^+}$ are the fractions of $H_2SO_4^+$ and HSO_3^+ in $H_xSO_y^+$ fragments (SO^+ , SO_2^+ , SO_3^+ , HSO_3^+ , and $H_2SO_4^+$), respectively. Both fractions approach zero when organic sulfate contribution to total sulfate increases.³ Data are 10-min averaged data. The blue data points correspond to office background and the red data points correspond to device operation. The two blue data points at around [0.04, 0.01] are the data points immediately after the device was turned off.



40

Figure S6. Time series of ozone, NO, and NO₂ from the office measurements. The device was in operation from 2:30 pm to 4:00 pm (highlighted in yellow).

45 Reference

1. Kuwata, M.; Zorn, S. R.; Martin, S. T., Using Elemental Ratios to Predict the Density of Organic Material Composed of Carbon, Hydrogen, and Oxygen. *Environmental Science & Technology* **2012**, *46* (2), 787-794.
2. Engelhart, G. J.; Hildebrandt, L.; Kostenidou, E.; Mihalopoulos, N.; Donahue, N. M.; Pandis, S. N., Water content of aged aerosol. *Atmos. Chem. Phys.* **2011**, *11* (3), 911-920.
3. Chen, Y.; Xu, L.; Humphry, T.; Hettiyadura, A. P. S.; Ovadnevaite, J.; Huang, S.; Poulain, L.; Schroder, J. C.; Campuzano-Jost, P.; Jimenez, J. L., et al., Response of the Aerodyne Aerosol Mass Spectrometer to Inorganic Sulfates and Organosulfur Compounds: Applications in Field and Laboratory Measurements. *Environmental Science & Technology* **2019**, *53* (9), 5176-5186.

55