

Appendix

I. Additional Methods

Reporting of swab tests taken outside the study

Throughout the study, participants reporting symptoms in the previous week were also asked to report the outcome of any NHS COVID-19 swab test for that illness. From the week commencing 28 September 2020, participants could also report the results of any swab testing in the previous week, regardless of whether they were also reporting symptoms for that week. This was to ensure that all swab test results could be reported regardless of when the test results arrived or whether the swab was taken asymptotically.

Tests taken within the study

The laboratory sub-cohort is split into three groups with different swabbing protocols. In all three groups, participants are asked to provide self-administered swabs on day two of illness if they experience one or more of the trigger symptoms. Group 1 (7,368 participants) swab for respiratory symptoms, fever or feverishness or loss or change to sense of smell or taste. Group 2 (1,042 participants) swab for respiratory, gastrointestinal or general symptoms of infection including fever or feverishness or loss or change to sense of smell or taste. Group 3 (2,216 participants) swab for cough, fever or feverishness or loss or change to sense of smell or taste. In group 3, if any household member experiences a trigger symptom then all household contacts are also asked to swab on the same day as the index case regardless of whether they are symptomatic and if any of those swabs are PCR+ then all household members are asked to swab again one and two weeks later irrespective of symptoms.

Matching of illness episodes and test results

To differentiate COVID-19 from non-COVID-19 illnesses, we matched illness episodes and self-reported swab test results that were within up to two weeks of each other. Matching was undertaken in a stepwise manner, for both symptomatic swabbing and asymptomatic swabbing, on a week-by-week basis, based on the likely sequence of symptoms and swab positivity. First, swab test results were matched to illnesses occurring in the same week as receipt of the swab result. Following that, any test results that did not link to an illness in the same week, were sequentially matched to illness episodes that had commenced in the following week (+1 week), then finished in the week prior (-1 week), and then finished two weeks prior (-2 weeks). Illnesses commencing 2 weeks after a swab result were not matched as these were considered unlikely to be related. For each round of sequential matching, only test results not yet matched to an illness were used.

Some illness episodes continued for several weeks, and the associated swab test may have been completed any time during the episode, or shortly before or after it. If more than one swab test result matched to an illness episode, any positive swab test result superseded any negative swab test result. Any illness episode matching to a positive swab result was taken to be a confirmed COVID-19 illness. Any illness episode matching to negative swab test results only, was taken to be a non-COVID-19 illness. Illness episodes not matching any swab test results, and swab test results not matching to any illness episodes, were not included in these analyses. Where an individual had multiple test positive illnesses, we only analysed their first test positive illness as it was felt that this was most likely to be the result of long-term virus shedding rather than re-infection.

Calculation of test characteristics for clinical case definitions

We calculated the diagnostic test characteristics of clinical case definitions and individual symptoms using the formulae from Figure S1.

II. Additional Tables and Figures

Figure S1. Two by two contingency table and equations used to calculate test characteristics of the clinical case definitions compared to swab test results (gold standard).

		Gold standard test (Swab test result)		Totals
		Positive	Negative	
Clinical Case Definition	Positive	a	b	a+b
	Negative	c	d	c+d
Totals		a+c	b+d	a+b+c+d

True Positives = a	False Positives = b
True Negatives = d	False Negatives = c

$$\text{True prevalence} = \frac{a + c}{a + b + c + d}$$

$$\text{Sensitivity} = \frac{a}{a + c}$$

$$\text{Specificity} = \frac{d}{b + d}$$

$$\text{Predictive value of a positive test (PPV)} = \frac{a}{a + b}$$

$$\text{Predictive value of a negative test (NPV)} = \frac{d}{c + d}$$

$$\text{Number needed to test (NNT)} = \frac{1}{PPV}$$

Figure S2. Percent of illnesses with a swab test result by month, overall and by UK Case Definition status




Figure S3. Ratio of swabbing among illnesses meeting the case definition to those not meeting the case definition over time.




Table S1. Details of symptoms and symptom groupings

	Symptom name	Symptom wording and combinations
Grouped Symptoms (used in case definitions)	cough	This was a combination of the following individual symptoms: 'dry_cough', 'white_phlegm' and 'green_phlegm' (i.e. "Dry Cough" AND/OR "Coughing up White Phlegm" AND/OR "Coughing up Green Phlegm")
	fever	This was a combination of the following individual symptoms: 'fever_cat' and 'feeling_feverish' (i.e. "Fever" with reported temperature $\geq 37.8C$ AND/OR "Feeling feverish")
	smell_taste	This was a combination of the following individual symptoms: 'loss_of_smell' and 'loss_of_taste' (i.e. "Loss or change to sense of smell" AND/OR "Loss or change to sense of taste")
Individual Symptoms	fever_cat	"Fever" with reported temperature $\geq 37.8C$
	feeling_feverish	"Feeling Feverish"
	chills	"Chills and shakes"
	night_sweats	"Night Sweats"
	muscle_ache	"Muscle Ache"
	bone_ache	"Bone or Joint ache"
	loss_of_appetite	"Loss of Appetite"
	headache	"Headache"
	concentration	"Lack of Concentration"
	dizzy	"Lightheaded or Dizzy"
	not_sleeping	"Not Sleeping"
	fatigue	"Fatigue / Feeling Unusually Tired"
	daily_activities	"Difficulties with daily activities around the house"
	extra_bed	"Needed extra time in bed"
	out_of_bed	"Could not get out of bed"
	confusion	"Confusion, disorientation, or hallucinations (altered mental state)"
	runny_nose	"Runny Nose"
	blocked_nose	"Blocked Nose"
	sinus_pain	"Sinus Pain / congestion"
	dry_cough	"Dry cough"
	white_phlegm	"Coughing up White Phlegm"
	green_phlegm	"Coughing up Green Phlegm"
	loss_of_smell	"Loss or change to sense of smell"
	loss_of_taste	"Loss or change to sense of taste"
	sneezing	"Sneezing"
	sore_throat	"Sore Throat"
	swollen_tonsils	"Swollen Tonsils"
	swollen_glands	"Swollen Glands (enlarged lymph nodes)"
	ear_pain	"Ear Pain or change in hearing"
	ear_fluid	"Fluid leaking from ear"
	sob	"Shortness of Breath / difficulty breathing"
	wheezing	"Wheezing"
	chest_pain_nc	"Chest pain (not changed by breathing or moving)"
	chest_pain_br	"Chest pain when breathing in"
	diarrhoea	"Diarrhoea (even mild)"
	vomiting	"Vomiting (being sick)"
	nausea	"Nausea (feeling sick)"
	abdo_pain	"Abdominal pain (not including menstrual pain)"
	redeye	"Eye Redness"
	sticky_eye	"Sticky Eye(s)"
eye_pain	"Eye Pain"	
eye_deter	"Deterioration of eyesight"	
rash_allover	"Rash - All Over"	
rash_local	"Rash - Local"	

Table S2. Speed of symptom onset, proportion of all community cases tested, and the test characteristics of each symptom.

Symptom	Onset Mean	Onset Median	Sensitivity	Specificity	PPV	NPV	Proportion tested*
cough_bin	2.0	1	62%	63%	18%	93%	53%
smell_taste_bin	3.0	1	43%	92%	42%	93%	77%
fever_bin	1.9	1	45%	79%	22%	92%	62%
fevercat_bin	2.1	1	23%	90%	23%	90%	78%
feeling_feverish_bin	1.9	1	36%	86%	25%	91%	58%
chills_bin	2.0	1	33%	88%	26%	91%	53%
night_sweats_bin	2.4	1	25%	91%	28%	90%	51%
muscle_ache_bin	1.8	1	59%	74%	23%	93%	44%
bone_ache_bin	2.0	1	38%	85%	24%	91%	41%
loss_of_appetite_bin	2.3	1	41%	88%	31%	92%	59%
headache_bin	1.8	1	67%	62%	19%	94%	41%
concentration_bin	2.2	1	28%	91%	28%	91%	48%
dizzy_bin	2.5	1	33%	87%	25%	91%	48%
not_sleeping_bin	2.4	1	25%	90%	24%	90%	42%
fatigue_bin	1.7	1	68%	65%	20%	94%	44%
daily_activities_bin	2.3	1	39%	89%	31%	92%	51%
extra_bed_bin	2.2	1	48%	81%	24%	92%	48%
out_of_bed_bin	2.5	1	16%	96%	35%	90%	55%
confusion_bin	3.1	1	5%	99%	41%	89%	55%
runny_nose_bin	1.7	1	36%	67%	13%	89%	33%
blocked_nose_bin	2.0	1	37%	78%	18%	90%	37%
sinus_pain_bin	2.3	1	25%	87%	21%	90%	43%
dry_cough_bin	2.2	1	53%	73%	20%	92%	59%
white_phlegm_bin	3.0	1	18%	90%	19%	89%	44%
green_phlegm_bin	3.3	1	8%	94%	15%	89%	51%
loss_of_smell_bin	3.0	1	37%	95%	49%	92%	81%
loss_of_taste_bin	3.1	1	37%	94%	45%	92%	79%
sneezing_bin	1.9	1	37%	74%	16%	90%	34%
sore_throat_bin	1.7	1	40%	65%	13%	89%	41%
swollen_tonsils_bin	2.3	1	5%	96%	14%	89%	54%
swollen_glands_bin	2.7	1	9%	94%	15%	89%	51%
ear_pain_bin	2.6	1	13%	91%	16%	89%	41%
ear_fluid_bin	4.1	1	1%	99%	18%	89%	36%
sob_bin	2.7	1	26%	91%	27%	90%	56%
wheezing_bin	2.8	1	12%	93%	17%	89%	52%
chest_pain_nc_bin	3.3	1	11%	97%	30%	89%	60%
chest_pain_br_bin	3.5	1	10%	97%	30%	89%	68%
diarrhoea_bin	2.1	1	29%	86%	21%	90%	32%
vomiting_bin	2.6	1	4%	97%	17%	89%	34%
nausea_bin	2.2	1	18%	91%	21%	90%	37%
abdo_pain_bin	2.3	1	13%	90%	14%	89%	33%
redeye_bin	2.4	1	7%	96%	17%	89%	32%
sticky_eye_bin	2.3	1	5%	96%	14%	89%	29%
eye_pain_bin	2.7	1	10%	97%	28%	89%	38%
eye_deter_bin	2.7	1	3%	98%	19%	89%	33%
rash_allover_bin	2.5	1	1%	99%	14%	89%	36%
rash_local_bin	2.6	1	5%	97%	18%	89%	33%

* Included tested illnesses where test result was not reported to the study


Table S3. Severity of symptoms among swab-confirmed COVID-19 positive and negative illnesses.

Symptoms	COVID+ illnesses (N=944)					COVID- illnesses (N=7269)				
	n	%	Max severity during illness (when symptom is present)			n	%	Max severity during illness (when symptom is present)		
			% mild	% mod	% severe			% mild	% mod	% severe
cough	582	62%	40%	45%	16%	2699	37%	42%	46%	12%
smell_taste	409	43%	21%	35%	44%	568	8%	37%	45%	18%
fevercat	219	23%	80%	17%	2%	738	10%	76%	20%	4%
feeling_feverish	343	36%	27%	51%	21%	1049	14%	37%	47%	16%
chills	308	33%	25%	47%	28%	876	12%	33%	47%	20%
night_sweats	238	25%	26%	40%	34%	626	9%	29%	49%	22%
muscle_ache	561	59%	28%	48%	24%	1855	26%	34%	50%	16%
bone_ache	355	38%	27%	46%	27%	1105	15%	30%	49%	21%
loss_of_appetite	391	41%	27%	49%	25%	855	12%	33%	46%	20%
headache	632	67%	28%	45%	27%	2759	38%	31%	50%	18%
concentration	266	28%	38%	46%	15%	678	9%	37%	46%	17%
dizzy	311	33%	43%	39%	18%	925	13%	46%	39%	15%
not_sleeping	238	25%	28%	50%	22%	758	10%	32%	47%	21%
fatigue	643	68%	18%	47%	35%	2527	35%	24%	51%	25%
daily_activities	368	39%	23%	41%	36%	814	11%	19%	46%	34%
extra_bed	450	48%	22%	43%	35%	1396	19%	21%	49%	30%
out_of_bed	153	16%	14%	30%	56%	281	4%	13%	28%	59%
confusion	51	5%	47%	33%	20%	72	1%	39%	43%	18%
runny_nose	339	36%	53%	35%	12%	2369	33%	45%	41%	14%
blocked_nose	345	37%	41%	44%	14%	1602	22%	40%	46%	14%
sinus_pain	240	25%	30%	48%	21%	912	13%	33%	48%	19%
dry_cough	497	53%	42%	43%	15%	1997	27%	43%	45%	12%
white_phlegm	171	18%	47%	40%	12%	729	10%	49%	41%	10%
green_phlegm	77	8%	38%	47%	16%	448	6%	38%	47%	15%
loss_of_smell	349	37%	17%	38%	46%	370	5%	34%	45%	21%
loss_of_taste	350	37%	22%	36%	42%	424	6%	38%	45%	17%
sneezing	352	37%	64%	26%	10%	1916	26%	57%	33%	10%
sore_throat	382	40%	45%	40%	15%	2574	35%	43%	43%	14%
swollen_tonsils	43	5%	37%	37%	26%	261	4%	30%	48%	22%
swollen_glands	81	9%	43%	46%	11%	459	6%	45%	40%	15%
ear_pain	125	13%	45%	43%	12%	643	9%	47%	42%	12%
ear_fluid	8	1%	63%	38%	0%	37	1%	51%	38%	11%
sob	250	26%	34%	48%	18%	673	9%	35%	46%	18%
wheezing	109	12%	54%	36%	10%	532	7%	44%	41%	14%
chest_pain_nc	107	11%	43%	45%	12%	251	3%	45%	41%	14%
chest_pain_br	97	10%	39%	41%	20%	226	3%	45%	41%	14%
diarrhoea	271	29%	47%	35%	17%	1048	14%	43%	37%	20%
vomiting	41	4%	32%	46%	22%	202	3%	31%	41%	28%
nausea	174	18%	40%	40%	20%	645	9%	37%	43%	20%
abdo_pain	118	13%	42%	42%	16%	699	10%	31%	45%	24%
redeye	65	7%	46%	49%	5%	317	4%	49%	40%	10%
sticky_eye	44	5%	55%	45%	0%	277	4%	56%	35%	9%
eye_pain	94	10%	36%	48%	16%	242	3%	52%	35%	12%
eye_deter	31	3%	42%	45%	13%	135	2%	50%	40%	10%
rash_allover	6	1%	33%	33%	33%	37	1%	27%	43%	30%
rash_local	44	5%	43%	50%	7%	194	3%	42%	41%	17%

Table S4. Duration of illness among swab-confirmed COVID-19 positive and negative illnesses, overall and stratified by age group and sex.

Illness Type	Strata		N	Mean	Standard Deviation	Range		Percentile			
						Lower	Upper	25th	50th	75th	90th
COVID+ve	Overall	Overall	944	10.9	9.7	1	70	5	8	14	21
	By age group	0-15	76	6.6	5.4	1	25	2	6	9	14
		16-24	54	7.4	7.9	1	56	3	6	9	13
		25-44	270	9.7	8.4	1	63	4	7	12	18
		45-64	379	12.5	10.5	1	70	6	9	15	25
		65+	163	12.5	10.9	1	63	6	10	14	23
	By sex	Male	359	10.7	8.7	1	62	5	8	14	21
Female		517	11.6	10.6	1	70	5	9	14	23	
COVID-ve	Overall	Overall	7269	5.9	7.3	1	182	2	4	7	12
	By age group	0-15	1038	5.1	4.4	1	36	2	4	7	11
		16-24	246	5.1	5.5	1	45	2	4	7	10
		25-44	1883	5.7	6.1	1	66	2	4	7	11
		45-64	2779	6.0	8.0	1	182	2	4	7	12
		65+	1262	7.1	9.2	1	166	2	5	7	14
	By sex	Male	2245	5.9	8.2	1	166	2	4	7	12
Female		4680	6.0	7.0	1	182	2	4	7	12	

Figure S4. Distribution of illness duration* by age group among COVID-19 positive illnesses



*Data limited to illnesses with duration <50 days.