

1 Supplemental Table 1: Mortality rates, and crude Incidence Rate Ratios (IRRc) for all variables of interest.

2

Variable / Categories	Total N=2,685 (100.0%)	Dead N=752 (28.0%)	Person- days ϕ	Rate [†]	IRRc	IRRc 95%CI	p-value
Wave							
First	884 (100.0%)	297 (33.6%)	133.450	2.23	Intercept		
Second	1,801 (100.0%)	455 (25.3%)	274.324	1.66	0.75	(0.64, 0.86)	< 0.001
Sex at birth							
Female	1,139 (100.0%)	274 (24.1%)	165.514	1.66	Intercept		
Male	1,546 (100.0%)	478 (30.9%)	242.260	1.97	1.19	(1.03, 1.38)	< 0.001
Age grouped (years)¹							
[18,40)	217 (100.0%)	21 (9.7%)	21.905	0.96	Intercept		
[40,60)	678 (100.0%)	91 (13.4%)	92.474	0.98	1.03	(0.65, 1.69)	
[60,80)	1,067 (100.0%)	326 (30.6%)	176.710	1.84	1.92	(1.27, 3.09)	
≥80	723 (100.0%)	314 (43.4%)	116.685	2.69	2.81	(1.85, 4.50)	< 0.001
Ethnicity							
White	928 (100.0%)	270 (29.1%)	155.554	1.74	Intercept		
Asian	491 (100.0%)	157 (32.0%)	66.030	2.38	1.37	(1.12, 1.67)	
Black	392 (100.0%)	114 (29.1%)	60.200	1.89	1.09	(0.87, 1.35)	
Other	634 (100.0%)	144 (22.7%)	89.640	1.61	0.93	(0.75, 1.13)	
Unknown	240 (100.0%)	67 (27.9%)	36.350	1.84	1.06	(0.81, 1.38)	0.009
CFS							
1: Very fit	195 (100.0%)	23 (11.8%)	21.870	1.05	Intercept		
2: Fit	290 (100.0%)	63 (21.7%)	41.995	1.50	1.43	(0.90, 2.35)	
3: Managing well	898 (100.0%)	165 (18.4%)	124.380	1.33	1.26	(0.83, 2.00)	
4: Vulnerable	323 (100.0%)	94 (29.1%)	56.970	1.65	1.57	(1.01, 2.53)	
5: Mildly frail	263 (100.0%)	90 (34.2%)	42.715	2.11	2.00	(1.29, 3.24)	
6: Moderately frail	354 (100.0%)	138 (39.0%)	64.909	2.13	2.02	(1.33, 3.22)	
7-9: Severely frail	360 (100.0%)	177 (49.2%)	54.925	3.22	3.06	(2.03, 4.86)	< 0.001
Missing	2	2					
CCI							
0	295 (100.0%)	16 (5.4%)	28.764	0.56	Intercept		
1	288 (100.0%)	35 (12.2%)	35.740	0.98	1.76	(0.99, 3.27)	
2	334 (100.0%)	71 (21.3%)	51.475	1.38	2.48	(1.48, 4.42)	
3	347 (100.0%)	79 (22.8%)	57.320	1.38	2.48	(1.49, 4.40)	
4	369 (100.0%)	103 (27.9%)	59.500	1.73	3.11	(1.89, 5.47)	
5	348 (100.0%)	112 (32.2%)	58.000	1.93	3.47	(2.12, 6.09)	
6	297 (100.0%)	139 (46.8%)	49.680	2.80	5.03	(3.10, 8.78)	
7	188 (100.0%)	88 (46.8%)	30.255	2.91	5.23	(3.16, 9.24)	
8_over	219 (100.0%)	109 (49.8%)	37.040	2.94	5.29	(3.23, 9.29)	< 0.001
BMI¹ (kg/m²)							
<18.5	128 (100.0%)	44 (34.4%)	23.980	1.83	Intercept		
[18.5,25)	784 (100.0%)	225 (28.7%)	134.840	1.67	0.91	(0.67, 1.27)	
[25,30)	784 (100.0%)	197 (25.1%)	128.134	1.54	0.84	(0.61, 1.18)	
≥30	725 (100.0%)	152 (21.0%)	107.140	1.42	0.77	(0.56, 1.09)	< 0.001

Missing	264	134						
Social history								
Independent	1,566 (100.0%)	315 (20.1%)	217.410	1.45	Intercept			
Family Assistance	503 (100.0%)	209 (41.6%)	85.160	2.45	1.69	(1.42, 2.02)		
Formal Assistance	357 (100.0%)	131 (36.7%)	69.459	1.89	1.30	(1.06, 1.59)		
Care Home	235 (100.0%)	94 (40.0%)	30.435	3.09	2.13	(1.68, 2.67)	< 0.001	
Missing	24	3						
Symptomatic²								
No	408 (100.0%)	85 (20.8%)	92.110	0.92	Intercept			
Yes	2,277 (100.0%)	667 (29.3%)	315.664	2.11	2.29	(1.84, 2.89)	< 0.001	
ICU admission								
No	2,073 (100.0%)	451 (21.8%)	280.554	1.61	Intercept			
Yes	612 (100.0%)	301 (49.2%)	127.220	2.37	1.47	(1.27, 1.70)		
Oxygen requirement								
No	683 (100.0%)	11 (1.6%)	104.434	0.11	Intercept			
Yes	1,982 (100.0%)	733 (37.0%)	302.070	2.43	23.04	(13.38, 44.53)	< 0.001	
Missing	20	8						
HFNO/CPAP								
No	2,185 (100.0%)	520 (23.8%)	308.964	1.68	Intercept			
Yes	473 (100.0%)	222 (46.9%)	92.890	2.39	1.42	(1.21, 1.66)	< 0.001	
Missing	27	10						
Invasive ventilation								
No	2,264 (100.0%)	500 (22.1%)	307.749	1.62	Intercept			
Yes	405 (100.0%)	248 (61.2%)	97.145	2.55	1.57	(1.35, 1.83)	< 0.001	
Missing	16	4						
Dexamethasone								
No	1,399 (100.0%)	376 (26.9%)	215.994	1.74	Intercept			
Yes	1,286 (100.0%)	376 (29.2%)	191.780	1.96	1.13	(0.98, 1.30)	0.173	
Tocilizumab								
No	2,438 (100.0%)	665 (27.3%)	362.224	1.84	Intercept			
Yes	247 (100.0%)	87 (35.2%)	45.550	1.91	1.04	(0.83, 1.29)	0.008	
Remdesivir								
No	2,116 (100.0%)	609 (28.8%)	317.149	1.92	Intercept			
Yes	569 (100.0%)	143 (25.1%)	90.625	1.58	0.82	(0.68, 0.98)	0.085	

3 (φ) person-days per 100 (¶) mortality rate per 100 person-days

4 (1) Brackets indicate that the side of the interval is closed; parenthesis, that the number is excluded. (2) Symptomatic at admission

5 CFS: Clinical Frailty Score, CCI: Charlson Comorbidity Index, ICU: Intensive Care Unit, HFNO/CPAP: High Flow Nasal Oxygen/Continuous

6 Positive Airway Pressure

7

8 Supplemental Table 2: Crude and adjusted IRR for the effect of wave on mortality among ICU admissions

Covariate	N	Missing	IRRc	IRRa 95%CI	IRRa	IRRa 95%CI
Dexamethasone	612	0	0.98	(0.77, 1.24)	2.19	(1.30, 3.68)
Remdesivir	612	0	0.98	(0.77, 1.24)	1.15	(0.88, 1.50)
Age grouped (years)	612	0	0.98	(0.77, 1.24)	0.82	(0.65, 1.05)
Charlson Comorbidity Index (CCI)	612	0	0.98	(0.77, 1.24)	0.84	(0.66, 1.08)
Tocilizumab	612	0	0.98	(0.77, 1.24)	1.07	(0.84, 1.38)
Clinical frailty score (CFS)	612	0	0.98	(0.77, 1.24)	0.89	(0.70, 1.14)
HFNO/CPAP	604	8	0.96	(0.76, 1.21)	1.03	(0.80, 1.34)
Invasive ventilation	612	0	0.98	(0.77, 1.24)	1.03	(0.81, 1.31)
Social history	604	8	0.97	(0.77, 1.23)	0.95	(0.75, 1.20)
Ethnicity	612	0	0.98	(0.77, 1.24)	0.97	(0.76, 1.25)
BMI grouped (kg/m ²)	563	49	1.14	(0.88, 1.50)	1.14	(0.88, 1.49)
Symptomatic at admission	612	0	0.98	(0.77, 1.24)	0.98	(0.78, 1.24)
Sex	612	0	0.98	(0.77, 1.24)	0.98	(0.78, 1.24)

9 IRRc: crude RR. IRRa: adjusted RR. Covariates above the dotted line are those change in the magnitude of the effect was $\geq 5.5\%$

10 ICU: Intensive Care Unit, HFNO/CPAP: High Flow Nasal Oxygen/Continuous Positive Airway Pressure

11

12 Supplemental Table 3: Multivariable causal models for the effect of wave on in-hospital mortality

#	MODEL	N	Failures	Df	RMSE	SE(β)	HR	HR 95%CI
m1	wave	2,685	752	1	--	0.075	0.75	(0.65, 0.87)
m2	wave + (A1, A2)	2,685	752	5	--	0.075	0.76	(0.65, 0.88)
m3	wave + (A1, A2) + (V1 - V6)	2,646	738	18	0.131	0.131	1.65	(1.28, 2.24)
m4	wave + (A1, A2) + (V1 - V4)	2,665	744	16	0.129	0.128	1.62	(1.26, 2.08)
i1	wave	612	301	1	--	0.120	0.95	(0.75, 1.20)
i2	wave + (A1, A2)	612	301	5	--	0.124	0.80	(0.62, 1.01)
i3	wave + (A1, A2) + (C1 -C6)	604	298	23	0.323	0.323	1.93	(1.02, 3.63)
i4	wave + (A1, A2) + (C1 -C4)	604	298	9	0.306	0.304	2.00	(1.10, 3.62)

13 m: model in overall population, i: model in ICU admission only (sub-analysis)

14 #1: crude model, #2: adjusted for a priori confounders (forced variables), #3: full model, #4: reduced model

15 (A1) Age fitted as flexible spline, (A2) Gender.

16 (V1): Dexamethasone, (V2): Oxygen requirement, (V3): Symptomatic at admission, (V4): CCI. (V5): ICU admission, (V6): HFNO/CPAP.

17 (C1): Dexamethasone, (C2): Remdesivir, (C3): Tocilizumab, (C4): HFNO/CPAP, (C5): CFS, (C6): CCI.

18 Df: model degrees of freedom, RMSE: Root Mean Square Error, SE(β) SE: of the coefficient (log scale), HR: Hazard Ratio.

19