

Supplemental Table 1: Predictor matrix

Type	Response	Predictors																							
		migr	age_log	age	age_sb	sex	sexFemale:age	who_sqrt	who	who_sb	who:sexFemale	who:age	viol	ses	ptsd	ptsd_sb	edu	child	smok	bp_log	bp	bp_sb	hgt_log	hgt	hgt_sb
b	<i>migr</i>																								
c	<i>age_log</i>																								
pas	age		1											1	1	1	1	1				1	1		
pas	age_sb			1																					
b	<i>sex</i>																		1		1	1		1	1
pas	sexFemale:age			1		1																			
c	<i>who_sqrt</i>			1	1	1	1						1	1	1	1									
pas	who							1																	
pas	who_sb								1																
pas	who:sexFemale					1				1															
pas	who:age			1						1															
b	<i>viol</i>			1	1	1	1		1	1	1				1	1									
o	<i>ses</i>			1	1	1	1	1	1	1	1				1	1			1						
c	<i>ptsd</i>			1	1	1	1		1	1	1		1	1	1										
pas	ptsd_sb														1										
o	<i>edu</i>			1	1	1	1						1												
o	<i>child</i>	1		1	1	1	1						1				1					1	1		
b	<i>smok</i>	1				1					1	1										1	1		1
c	<i>bp_log</i>			1	1	1	1																		
pas	bp																		1						
pas	bp_sb																					1			
c	<i>hgt_log</i>					1									1	1			1			1	1		
pas	hgt																						1		
pas	hgt_sb																							1	

The predictor matrix details how each partially observed variable was imputed. Horizontally the variables enter as the response, vertically the variables enter as the predictor when the corresponding cell is “1”. The “type” column indicate how the variables should be modelled when entering as the response variable (b, binary; c, continuous; pas, passive; o, ordered/ordinal; u, unordered/nominal). Variables in red are part of the substantive model (the propensity score model); variables in italic are actively imputed. “:” denotes interaction between the variables on each side.