

The CT Scan Lung Severity Score and Vaccination Status in COVID-19 patients during the Second Wave in India: Perspective of an Independent Radiology Practice

Revat T. Lakhia, MD, MPH¹, Jaimin R. Trivedi, MBBS, MPH²

1 – Kadamb Diagnostics, Ahmedabad, India

2- University of Louisville School of Medicine, Louisville, KY

Correspondence

Jaimin R. Trivedi, MBBS, MPH

Assistant Professor

Department of Cardiovascular and Thoracic Surgery

University of Louisville School of Medicine

201 Abraham Flexner Way, Suite 1200

Louisville, KY 40202

(708) 606 – 3115, jaimin.trivedi@louisville.edu

Abstract

Background

The COVID-19 patients often undergo a high-resolution CT scan to determine extent of their lung involvement. The aim of this study was to determine lung involvement in confirmed/suspected COVID-19 patients (encountered at an independent radiology practice) and its correlation to vaccination status amidst the second COVID-19 wave in India.

Methods

We retrospectively queried our data since April 2021 to identify adult patients (>17 years) who had confirmed (positive RT-PCR or antigen test) or suspected COVID-19 (classic symptoms but negative RT-PCR) and received a high-resolution CT scan to determine the extent of their lung involvement using the CT severity (CORAD) score. Patients were classified in 3 groups based on their vaccination status to determine its correlation with the CORAD score: fully vaccinated, partially vaccinated, and unvaccinated. Basic descriptive statistics, univariate tests and multivariate linear regression analysis were used.

Results

We identified 229 patients (median age 45 years, 60% male) of which 205 (89%) had confirmed COVID-19 (positive RT-PCR) and 24 had suspected disease (negative RT-PCT but classic symptoms). Of the 229 patients 29 (13%) had complete vaccination, 38 (17%) had partial vaccination and 162 (70%) had no vaccination. The CT score of the completely vaccinated patients was significantly lower compared to partially or unvaccinated patients (median 0 v. 3.5 v. 10, respectively $p < .01$). A multivariate linear regression model showed that partial or fully vaccinated patients had lower CT severity score compared to unvaccinated patients (adjusted R squared = 0.47).

Conclusion

We present here the real-world findings from an independent radiology practice (a unique and common practice model), in India amid the second COVID-19 wave showing significantly lower CT severity score in fully or partially vaccinated patients compared to unvaccinated patients. Complete vaccination in patients could be critical in preventing severe lung disease.

Introduction

The novel coronavirus (SARS nCOV-2) impacted over 160 million people across the globe causing severe respiratory illness in many patients with mortality of over 2%¹. Countries like USA, UK and South Africa have seen deadlier second waves due to highly infectious mutated strains; one such (B1617 or delta) is currently rampant in India crippling the already constrained healthcare infrastructure². There are some patients with symptomatic disease but have negative RT-PCR or rapid antigen test. These patients often undergo a high-resolution CT scan to determine extent of their lung involvement and aid in further management³.

Unlike the western countries, the radiology practice in India is largely done by independently operated groups (without affiliations to hospitals or care centers)⁴. A common patient is often referred to these independent practices including suspected/confirmed COVID-19 patients to determine the degree of their lung involvement. We have seen several patients in our practice during the second COVID-19 wave who had negative RT-PCR test but presented with classic symptoms such as fever, fatigue, body aches, occasional shortness of breath etc. Our case mix included fully or partially vaccinated or completely unvaccinated patients. At the time of the study there were 2 vaccines approved in India, both of which required 2 doses for complete vaccination⁵. The aim of this study was to determine lung involvement in confirmed/suspected COVID-19 patients and its correlation to vaccination status.

Methods

We retrospectively queried our data since April 2021 to identify adult patients (>17 years) who had confirmed (positive RT-PCR or antigen test) or suspected COVID-19 (classic symptoms but negative RT-PCR) and received a high-resolution CT scan to determine the extent of their lung involvement using the CT severity (CORAD) score⁶. Along with the RT-PCR results, we recorded vaccination status at the time of their CT scan. Patients with both doses completed 1 week prior to CT scan were considered fully vaccinated whereas patients receiving CT scan 2 weeks after the 1st dose or within 1 week of 2nd dose were considered partially vaccinated and the rest of the patients (not receiving any vaccine or within 2 weeks of 1st dose) were considered unvaccinated⁷. Based on the vaccination age eligibility criteria we stratified the cohort in 3 groups of 18-44 years, 45-59 years, and 60 years or above.

We used basic descriptive statistics (median (interquartile range) and % (N)), univariate tests (non-parametric (Kruskal-Wallis) test for continuous variables and chi-square test for categorical variables) and multivariate linear regression analysis to determine the association of the CORAD score to RT-PCR and vaccination status. All the analyses were conducted using SAS 9.4 software (SAS Inc. Cary, NC) at 95% confidence interval level.

Results

We identified 229 patients of which 205 (89%) had confirmed COVID-19 (positive RT-PCR) and 24 had suspected disease (negative RT-PCT but classic symptoms). The median age of the cohort was 45 (36-58) years, and 138 (60%) patients were male. The median CT score of the entire cohort was 9 (2-13). There were 49% (n=112), 29% (n=67) and 22% (n=50) patients in age groups 18-44 years, 45-59 years and > 60 years, respectively. Median CT score in patients in age group 18-44 years was 9 (0-12), 45-60 years was 11 (6-15) and >60 years was 8 (1-11).

Of the 229 patients only 29 (13%) had complete vaccination, 38 (17%) had partial vaccination and 162 (70%) had no vaccination. Of the RT-PCR positive patients, 29 (14%) had complete vaccination, 32 (15%) had partial vaccination. The median age in the vaccination groups was comparable (median 42 v. 48 v. 43, $p=0.32$, Table 1). There were more male patients in the unvaccinated group compared to fully or partially vaccinated groups (67% v. 42% v. 47%, $p<.01$). The overall CT score in RT-PCR positive cohort was significantly higher compared to the RT-PCR negative (median 10 vs. 0, $p<.01$) cohort. The CT score of the completely vaccinated patients was significantly lower compared to the partially or unvaccinated patients (median 0 v. 3.5 v. 10, $p<.01$, Figure 1). Within RT-CPR positive cohort, patients with complete vaccination had significantly lower CORAD score compared to partially or unvaccinated patients (median 0 vs. 4 vs. 11, $p=0.02$).

A multivariate linear regression model showed that higher age and presence of positive RT-PCR test were associated with higher CT severity score and partial or fully vaccinated patients had lower CT severity score compared to unvaccinated patients (adjusted R squared = 0.47).

Discussion

The second COVID-19 wave has put extreme pressure on India's healthcare system anecdotally affecting younger patients². We observed that almost half of all the patients in our cohort were 45 years or younger which is correlated with then national vaccination guidelines (vaccination for healthcare and frontline workers and aged over 45 years starting). We also observed that patients with complete or even partial vaccination had lower CT severity score which reattributes the effectiveness of the vaccines in preventing the severe disease⁷. In the 11% RT-PCR negative patients, we observed a low CT severity score (median 0). This summary describes COVID-19 patient profiles and its correlation with the CORAD score from an independent radiology practice perspective in India.

Even though radiology practice in India is predominantly independent, they likely encounter majority of COVID-19 cases needing imaging tests, including severely ill patients who may eventually get admitted to the hospitals for further treatment^{4,8}. Our study represents a unique and common practice across India and report the real-world evidence of the CT scan findings and its association with vaccination status and RT-PCR positivity in COVID-19 patients.

Vaccine effectiveness has been scrutinized with emergence of SARS nCoV-2 variants as B1617 (delta) or B117 (alpha) mutants appear to have higher transmissibility and case fatality than the unmutated version of the virus^{9,10}. Previous studies have suggested that vaccines are effective against variants in preventing severe disease^{11,12}. A recent report from India suggested that fully vaccinated patients may still get infected with the delta (B1617) variant however there was no mortality and high-grade fever was predominant symptom¹³. Similarly, our study showed that 29 fully vaccinated patients had a positive RT-PCR test. Although genomic sequencing data in our cohort was unavailable, based on the previous study, delta variant (B1617) could be predominantly responsible as our patients are from the same timeframe and geography.

Our study shows that compared to unvaccinated patients, completely or partially vaccinated patients had significantly lower CT severity score, irrespective of their RT-PCR status which corroborates the evidence that vaccines are particularly effective tool in preventing a severe COVID-19 even if effectiveness in limiting the spread could be lower in leu of emerging variants^{7,12,13}. This finding of reduced lung involvement with full or partial vaccination could be crucial in highlighting the role of

vaccines as an effective tool in fight against the COVID-19 during these times of vaccine misinformation to motivate more people to get vaccinated. We found 11% patients with RT-PCR negative but with classic COVID-19 symptoms which we think could be due to false negative tests, stage of the disease where nasopharyngeal swab is inconclusive or the newer variant of the virus^{14 15}.

Our study has several limitations being a retrospective single center study. We did not have patient details other than their basic demographics, COVID-19 RT-PCR status, vaccination status and CT severity score. Patient details such as their oxygen saturations, hospital admission status, co-morbidities and mortality were not available.

In conclusion, we present here the real-world findings from an independent radiology practice (a unique and common practice model), in India amid the second COVID-19 wave showing significantly lower CT severity score in fully or partially vaccinated patients compared to unvaccinated patients. We report a higher CT severity score in patients with positive RT-PCR. Complete vaccination in patients could be critical in preventing severe lung disease.

References

1. Roser M, Ritchie H, Ortiz-Ospina E, et al. Coronavirus Pandemic (COVID-19). Our World in Data, 2020.
2. Lancet T. India's COVID-19 emergency. *Lancet (London, England)* 2021;397(10286):1683.
3. Xie X, Zhong Z, Zhao W, et al. Chest CT for typical coronavirus disease 2019 (COVID-19) pneumonia: relationship to negative RT-PCR testing. *Radiology* 2020;296(2):E41-E45.
4. Jankharia GR. Commentary-radiology in India: the next decade. *The Indian journal of radiology & imaging* 2008;18(3):189.
5. Bagcchi S. The world's largest COVID-19 vaccination campaign. *The Lancet Infectious Diseases* 2021;21(3):323.
6. Simpson S, Kay FU, Abbara S, et al. Radiological society of north America expert consensus document on reporting chest CT findings related to COVID-19: endorsed by the society of thoracic Radiology, the American college of Radiology, and RSNA. *Radiology: Cardiothoracic Imaging* 2020;2(2):e200152.
7. Pilishvili T, Fleming-Dutra KE, Farrar JL, et al. Interim estimates of vaccine effectiveness of Pfizer-BioNTech and Moderna COVID-19 vaccines among health care personnel—33 US sites, January–March 2021. 2021
8. Ahuja G, Verma M, Patkar D. Financial impact of COVID-19 on radiology practice in India. *The Indian Journal of Radiology & Imaging* 2021;31(Suppl 1):S31.
9. Davies NG, Abbott S, Barnard RC, et al. Estimated transmissibility and impact of SARS-CoV-2 lineage B. 1.1. 7 in England. *Science* 2021;372(6538)
10. Grint DJ, Wing K, Williamson E, et al. Case fatality risk of the SARS-CoV-2 variant of concern B. 1.1. 7 in England, 16 November to 5 February. *Eurosurveillance* 2021;26(11):2100256.
11. Abu-Raddad LJ, Chemaitelly H, Butt AA. Effectiveness of the BNT162b2 Covid-19 Vaccine against the B. 1.1. 7 and B. 1.351 Variants. *N Engl J Med* 2021
12. Bernal JL, Andrews N, Gower C, et al. Effectiveness of the Pfizer-BioNTech and Oxford-AstraZeneca vaccines on covid-19 related symptoms, hospital admissions, and mortality in older adults in England: test negative case-control study. *BMJ* 2021;373

13. Singh UB, Rophina M, Chaudhry R, et al. Genomic analysis of symptomatic SARS-CoV-2 vaccine breakthrough infections from a tertiary care centre in India. 2021
14. Arevalo-Rodriguez I, Buitrago-Garcia D, Simancas-Racines D, et al. False-negative results of initial RT-PCR assays for COVID-19: a systematic review. *PLoS One* 2020;15(12):e0242958.
15. Tahan S, Parikh BA, Droit L, et al. SARS-CoV-2 E gene variant alters analytical sensitivity characteristics of viral detection using a commercial RT-PCR assay. *J Clin Microbiol* 2021;JCM. 00075-21.

Table 1. Baseline characteristics of patients stratified by vaccination status

	Fully Vaccinated N=29	Partially Vaccinated N=38	Not vaccinated N=162	p-value
Age (years)	42 (32-64)	48 (40-56)	43 (36-57)	0.32
Gender Male	42% (12)	47% (18)	67% (108)	<.01
RT-PCR positive	100% (29)	84% (32)	89% (144)	0.10
Median CT score	0 (0-3)	3.5 (0-9)	10 (7-14)	<.01

Data presented as median (interquartile range), % (n).

Figure 1. Median CT severity (CORAD) score stratified by vaccination status

