

Supplementary Table 1.

Primers and probes used in this study

Assay	Sequence (5' - 3') ^a	Length (bp)	Location in SARS-CoV-2 genome (bp)
MNV	F: CACGCCACCGATCTGTTCTG R: GCGCTGCGCCATCACTC P: 6FAM-CGCTTTGGAACAATG-MGB	108	4,972 – 5,080
PEDV	F: CAGGACACATTCTTGGTGGTCTT R: CAAGCAATGTACCACTAAGGAGTGTT P: FAM-ACGCGCTTCTCACTAC-MGB	140	26,010 - 26,149
SARS-CoV-2: E_Sarbecco	F: ACAGGTACGTTAATAGTTAATAGCGT R: ATATTGCAGCAGTACGCACACA P: 6FAM-ACACTAGCCATCCTTACTGCGCTTCG-BHQ	112	26,141 – 26,253
SARS-CoV-2: RdRp	F: GTGARATGGTCATGTGTGGCGG R: CARATGTTAAASACACTATTAGCATA P1: 6FAM-CCAGGTGGWACRTCATCMGGTGATGC-BHQ P2: 6FAM-CAGGTGGAACCTCATCAGGAGATGC-BHQ	99	15,361 – 15,460
SARS-CoV-2: N_Sarbecco	F: CACATTGGCACCCGCAATC R: GAGGAACGAGAAGAGGCTTG P: 6FAM-ACTTCCTCAAGGAACAACATTGCCA-BHQ	127	28,555 – 28,682

^a W is A/T; R is G/A; M is A/C; S is G/C. FAM: 6-carboxyfluorescein; MGB: minor groove binder; BHQ: blackhole quencher.