

COVID-Q: validation of the first COVID-19 questionnaire based on patient-rated symptom gravity

Spinato G^{1,2}, Fabbris C¹, Conte F³, Menegaldo A¹, Franz L¹, Gaudio P¹, Cinetto F⁴, Agostini C⁴, Costantini G³, Boscolo-Rizzo P⁵

1: Department of Neurosciences, Section of Otolaryngology and Regional Centre for Head and Neck Cancer, University of Padova, Treviso, Italy

2: Department of Surgery, Oncology and Gastroenterology, Section of Oncology and Immunology, University of Padova, Padova, Italy

3: Department of Psychology, University of Milano Bicocca, 20126 Milan, Italy

4: Department of Medicine, Clinical Immunology and Hematology, University of Padova, Treviso, Italy

5: Section of Otorhinolaryngology, Azienda Sanitaria Universitaria Integrata di Trieste, Trieste, Italy

Corresponding author:

Cristoforo Fabbris MD

Department of Neurosciences, University of Padova, Piazza Ospedale 1, 31100, Treviso, Italy -
cristoforo.fabbris@gmail.com

Conflicts of Interest, Source of Funding and Compliance with Ethical Standards

The authors have no conflict of interest, funding or financial relationships. An informed consent has been obtained for any procedure involving the patients described in this article. The manuscript has not been submitted to more than one journal for simultaneous consideration. The manuscript has not been published previously (partly or in full). All of the authors have participated in the planning, writing or revising the manuscript.

TABLES

Table 1. Item parameters and fit from final Item Response Theory (IRT) models.

	Discrimination	Threshold 1	Threshold 2	S_X2	p
<i>Asthenia</i>					
being awake most of the night	5.600	1.370	1.794	5.008	0.286
difficulty falling asleep	3.657	1.262	1.849	3.511	0.476
waking up several times at night	4.568	1.028	1.700	3.903	0.419
poor quality of sleeping	3.904	1.119	1.626	5.492	0.359
not feeling yourself	1.320	1.067	1.122	7.446	0.489
<i>Gastrointestinal symptoms</i>					
vomit ¹	4.203	1.805	-	4.849	0.183
nausea	2.703	1.485	1.905	9.807	0.133
diarrhoea	1.915	0.952	1.885	8.865	0.181
abdominal pain	2.340	1.724	2.502	1.373	0.712
<i>Ear and nose symptoms</i>					
headache	1.254	1.417	1.455	5.923	0.549
runny nose	2.136	1.420	2.030	2.634	0.621
blocked nose	2.136	1.101	2.026	4.287	0.509
sneezing	2.136	1.252	2.492	8.535	0.129
painful pressure in ears	2.136	1.919	2.627	6.381	0.094
watery eyes	2.136	1.765	3.082	3.696	0.158
<i>Breathing issues</i>					
problems with breathing	5.022	1.083	1.731	5.001	0.544
wheezing	6.659	1.193	1.829	7.909	0.161
shortness of breath	9.601	0.896	1.700	5.696	0.127
coughing up mucus ¹	1.333	1.910	-	2.640	0.955
dry cough	1.379	1.032	1.693	13.527	0.260
felt tired	1.683	0.547	0.980	12.926	0.114
<i>Throat symptoms</i>					
swollen glands	1.691	3.038	3.300	0.149	0.699
sore throat	3.870	1.313	1.820	3.711	0.054
hoarseness	1.154	2.038	2.678	8.197	0.085
tickles in the throat	5.046	1.041	1.893	1.195	0.274

Note. Discrimination = ability of the item to discriminate between individuals with different symptom frequencies. Threshold 1 = symptom frequency value that marks the cutoff between adjacent answer categories: 0 = “no symptom” and 1 = “I experience this symptom a little.” Threshold 2 = cutoff value between category 1 = “I experience this symptom a little” and 2 = “I experience this symptom a lot”

¹ Dichotomous item: 0 = symptom absent, 1 = symptom present. Threshold 2 was not computed for dichotomized items.

Table 2. Fit indices for the final Item Response Theory (IRT) models

	M2*	df	p	RMSEA	SRMSR	TLI	CFI
Asthenia	2,028*	5	0.845	0.000	0.016	1.003	1.000
Stomachache	3,664	1	0.056	0.076	0.084	0.955	0.993
Nose Symptoms	8,478	4	0.076	0.049	0.093	0.967	0.974
Breathing issues	4,669	4	0.323	0.019	0.075	0.998	0.999
Throat symptoms	3,334*	2	0.189	0.038	0.042	0.991	0.997

Note. p values > .05 indicate good model-data fit.

* The model did not have sufficient degrees of freedom for the computation of the M2* index. The less restrictive C2 index was used instead. The interpretation of p values remains unvaried³⁴.

Table 3 The final COVID-Q questionnaire.

<i>Asthenia</i>			
“Being awake most of the night”	0 (none)	1 (a little)	2 (a lot)
“Difficulty falling asleep”	0 (none)	1 (a little)	2 (a lot)
“Waking up several times at night”	0 (none)	1 (a little)	2 (a lot)
“Poor quality of sleeping”	0 (none)	1 (a little)	2 (a lot)
“Not feeling yourself”	0 (none)	1 (a little)	2 (a lot)
<i>Gastrointestinal symptoms</i>			
“vomit”	0 (absent)	1 (present)	
“nausea”	0 (none)	1 (a little)	2 (a lot)
“diarrhea”	0 (none)	1 (a little)	2 (a lot)
“abdominal pain”	0 (none)	1 (a little)	2 (a lot)
<i>Fever</i>			
“Feeling feverish”	0 (none)	1 (a little)	2 (a lot)
“Sweat”	0 (none)	1 (a little)	2 (a lot)
“Chills”	0 (none)	1 (a little)	2 (a lot)
<i>Ear and nose symptoms</i>			
“headache”	0 (none)	1 (a little)	2 (a lot)
“runny nose”	0 (none)	1 (a little)	2 (a lot)
“blocked nose”	0 (none)	1 (a little)	2 (a lot)
“sneezing”	0 (none)	1 (a little)	2 (a lot)
“watery eyes”	0 (none)	1 (a little)	2 (a lot)
<i>Breathing issues</i>			
“problems with breathing”	0 (none)	1 (a little)	2 (a lot)
“wheezing”	0 (none)	1 (a little)	2 (a lot)
“shortness of breath”	0 (none)	1 (a little)	2 (a lot)
“coughing up mucus”	0 (absent)	1 (present)	
“dry cough”	0 (none)	1 (a little)	2 (a lot)
“felt tired”	0	1	2

	(none)	(a little)	(a lot)			
<i>Throat symptoms</i>						
“swollen glands”	0 (none)	1 (a little)	2 (a lot)			
“sore throat”	0 (none)	1 (a little)	2 (a lot)			
“hoarseness”	0 (none)	1 (a little)	2 (a lot)			
“tickles in the throat”	0 (none)	1 (a little)	2 (a lot)			
<i>Muscle pain</i>	0 (none)	1 (a little)	2 (a lot)			
<i>Anosmia/Ageusia</i>	0 (no symptom)	1	2	3	4	5 (complete absence of smell or taste)

Table 4. Rare-event logistic regression predicting SARS-CoV-2 infection with unweighted sum of the breathing symptoms scale and of the anosmia/ageusia item. COVID-19 symptom questionnaire's partial score used as predictor.

Predictor	<i>b</i>	Exp(<i>b</i>)	<i>S.E.</i>	<i>Z</i>	<i>p</i>
(Intercept)	-5.066	0.006	0.185	-27.451	0.000
Demographics					
Age ^d	0.039	1.040	0.012	3.272	0.001
Sex ^{a,d}	1.211	3.357	0.309	3.915	0.000
COVID-Q					
Partial score	0.688	1.990	0.070	9.861	0.000
Comorbidities and smoking status					
Smoking status ^{b,d}	-0.526	0.591	0.328	-1.602	0.109
Hypertension ^{c,d}	0.521	1.683	0.414	1.258	0.208
Diabetes ^{c,d}	0.579	1.784	0.811	0.714	0.475
Cardiovascular diseases ^{c,d}	-0.034	0.966	0.590	-0.058	0.953
Cerebrovascular diseases ^{c,d}	-1.652	0.192	1.401	-1.179	0.238
Tumors ^{c,d}	-0.894	0.409	0.691	-1.294	0.196
COPD ^{c,d}	-1.573	0.207	0.756	-2.082	0.037
Renal Impairment ^{c,d}	0.736	2.087	1.210	0.608	0.543

Note. COPD = chronic obstructive pulmonary disease

^a Gender was coded as 0 for females and 1 for males. Reference was made for male gender.

^b Smoking status was coded as 0 for never, and 1 for former or current smokers.

^c Hypertension, Diabetes, Cardiovascular and Cerebrovascular diseases, Tumors, COPD and Renal impairment were coded as 0 for absent and 1 for present.

^d These predictors were mean centered for the regression analysis.