

**Supplementary Table S1.** Cox regression results, stratified by the presence of ischaemic heart disease. Hazard ratios (HR) were referenced against patients without fragmented QRS.

Subgroup	Outcome		Univariate		Multivariate <sup>1</sup>	
			HR [95% CI]	p value	HR [95% CI]	p value
Ischaemic heart	Composite primary outcome <sup>2</sup>		1.703 [1.166, 2.488]	0.006	1.746 [1.191, 2.559]	0.004
disease present	Secondary	Cardiovascular mortality	1.745 [1.142, 2.668]	0.010	1.802 [1.173, 2.768]	0.007
(N=786)	outcomes	Ventricular arrhythmia	2.842 [1.152, 7.014]	0.023	2.652 [1.065, 6.602]	0.036
		Sudden cardiac death	1.831 [0.993, 3.376]	0.053	1.881 [1.014, 3.481]	0.045
		Myocardial infarction	1.085 [0.684, 1.722]	0.728	1.052 [0.661, 1.672]	0.832
		New-onset atrial fibrillation	1.072 [0.709, 1.620]	0.742	1.116 [0.737, 1.692]	0.604
No ischaemic	Composite pri	imary outcome <sup>2</sup>	1.334 [0.926, 1.920]	0.121	1.342 [0.929, 1.938]	0.117
heart disease	Secondary	Cardiovascular mortality	1.062 [0.672, 1.678]	0.797	1.049 [0.662, 1.662]	0.839
(N=1396)	outcomes	Ventricular arrhythmia	3.487 [1.414, 8.601]	0.007	3.336 [1.343, 8.282]	0.009
		Sudden cardiac death	1.809 [1.071, 3.053]	0.027	1.928 [1.135, 3.278]	0.015
		Myocardial infarction	0.788 [0.402, 1.547]	0.489	0.895 [0.454, 1.762[	0.747
		New-onset atrial fibrillation	1.168 [0.870, 1.570]	0.302	0.835 [0.620, 1.125]	0.235

Adjusted for hypertension for patients with ischaemic heart disease, and renal diseases, atrial fibrillation, diabetes mellitus, calcium channel blocker usage, and usage of statins and fibrates for patients without ischaemic heart disease

<sup>&</sup>lt;sup>2</sup> A composite of cardiovascular mortality, ventricular arrhythmia, and sudden cardiac death CI, confidence interval.

**Supplementary Table S2.** Cox regression of the 179 patients with fragmented QRS. Hazard ratios were referenced against patients with fragmented QRS present in only two contiguous leads.

Outcome		Hazard ratio [95% confidence interval])	p value
Composite primary outcome <sup>1</sup>		1.841 [1.119, 3.029]	0.016
Secondary	Cardiovascular	1.539 [0.850, 2.786]	0.155
outcomes	mortality		
	Ventricular arrhythmia	1.897 [0.611, 5.888]	0.268
	Sudden cardiac death	2.866 [1.350, 6.081]	0.006
	Myocardial infarction	1.746 [0.835, 3.649]	0.139
	New-onset atrial	1.096 [0.676, 1.775]	0.710
	fibrillation		

fibrillation fibrillation arrhythmia, and sudden cardiac death