

SUPPLEMENTAL MATERIAL

Premature Atrial Stimulation Accentuates Conduction Abnormalities in Cardiac Surgery Patients that Develop Postoperative Atrial Fibrillation

Authors: Muhammad S. Khan, PhD,¹ Matthias Lange, PhD,¹ Ravi Ranjan, MD, PhD,^{1,2,3} Vikas Sharma, MD,⁴ Jason P. Glotzbach, MD,⁴ Craig Selzman, MD,^{1,4} Derek J. Dossdall, PhD,^{1,2,3,4,*}

Affiliations: ¹ Nora Eccles Harrison Cardiovascular Research and Training Institute,

² Division of Cardiovascular Medicine, Department of Internal Medicine,

³ Department of Bioengineering,

⁴ Division of Cardiothoracic Surgery, Department of Surgery,

The University of Utah – Health, Salt Lake City, UT 84112, USA.

*** Address For Correspondence:**

Dr. Derek J. Dossdall, PhD, FHRS, FAHA

Associate Professor of Surgery, Division of Cardiothoracic Surgery,

The University of Utah – Health, Salt Lake City, UT 84112 USA

Email: derek.dossdall@utah.edu / Ph: (+1) 801-587-2036




Fig. S1. (A) Flexible array of 240-electrodes used in the intraoperative study to collect epicardial atrial activation time data in sinus and S1S2 pacing. **(B)** A schematic representation of an array positioned on posterior LA wall in between PVs (not drawn to scale). Star (★) represents the site of pacing electrodes.

Table S1. Comparing effectiveness of a shortest premature captured atrial beat S2 with S1 (600 ms) to quantify CB, CD and CDCDB in patients that remained sinus and developed POAF during 1 – 4 days post-cardiac surgery.

| Group | Conduction disorder type | S1 (mm²) | S2 (mm²) | p value |
|--------------|---------------------------------|----------------------------|----------------------------|----------------|
| Sinus | CB | 0.56±1.18 | 1.34±2.86 | 0.126 |
| | CD | 3.33±3.36 | 6.06±4.22 | 0.004 |
| | CDCB | 3.89±4.39 | 7.41±6.39 | 0.004 |
| POAF | CB | 1.32±2.16 | 4.10±3.75 | 0.125 |
| | CD | 6.39±5.93 | 13.26±6.49 | 0.032 |
| | CDCB | 7.71±6.92 | 17.36±8.75 | 0.031 |