


Supplemental Materials

Household Transmission and Clinical Features of SARS-CoV-2 Infections by Age in 2 US Communities


Huong Q. McLean, PhD, MPH,^a Carlos G. Grijalva, MD, MPH,^b Kayla E. Hanson, MPH,^a Yuwei Zhu, MD, MS,^b Jessica E. Deyoe, MPH,^c Jennifer K. Meece, PhD,^a Natasha B. Halasa, MD, MPH,^b James D. Chappell, MD, PhD,^b Alexandra Mellis, PhD,^c Carrie Reed, DSc,^c Edward A. Belongia, MD,^a H. Keipp Talbot, MD, MPH,^b Melissa A. Rolfes, PhD^c

Affiliations: ^aMarshfield Clinic Research Institute, Marshfield, Wisconsin; ^bVanderbilt University Medical Center, Nashville, Tennessee; and ^cCenters for Disease Control and Prevention, Atlanta, Georgia


Supplemental Figure. Reported symptoms, timing, and duration of specific symptoms by age group among persons with SARS-CoV-2 infection in a prospective study of SARS-CoV-2 household transmission — Tennessee and Wisconsin, April 2020–April 2021.




c. neurologic symptoms



d. lower respiratory symptoms





Abbreviation: OR, odds ratio; CI, confidence interval; IQR, interquartile range; NE, not estimated

P value from Kruskal-Wallis Test comparing median days with symptoms across age groups.

¹Measure of association between age group and odds of symptom reported during follow-up

²Among contacts who reported the specific symptoms

³Gastrointestinal symptoms were assessed only for Wisconsin participants