

**Supplemental Table 1:** Associations between percent adequacy of total gestational weight gain and perinatal outcomes, modified by maternal pre-pregnancy BMI<sup>1</sup>.

Characteristic	n	Severely Inadequate (<70%)		Inadequate (70% to <90%)		Adequate (90% to 125%)		Excessive (>125%)	
		RR (95% CI)	P int <sup>2</sup>	RR (95% CI)	P int. <sup>2</sup>	RR (95% CI)	P int. <sup>2</sup>	RR (95% CI)	P int. <sup>2</sup>
<b>Stillbirth</b>	6,579								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		0.79 (0.33, 1.88)	0.52	0.48 (0.15, 1.59)	0.95	1.0		3.13 (0.96, 10.20)	<b>0.04</b>
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.17 (0.74, 1.87)	Ref.	1.01 (0.63, 1.63)	Ref.	1.0		1.73 (0.95, 3.15)	Ref.
Overweight/Obese (≥ 25 kg/m <sup>2</sup> )		1.49 (0.71, 3.13)	0.81	1.66 (0.69, 4.00)	0.99	1.0		1.17 (0.64, 2.12)	0.40
<b>Perinatal death</b>	6,376								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		0.91 (0.47, 1.78)	0.64	1.23 (0.63, 2.40)	0.31	1.0		2.45 (0.76, 7.93)	0.09
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		0.98 (0.68, 1.40)	Ref.	0.76 (0.52, 1.12)	Ref.	1.0		1.06 (0.63, 1.80)	Ref.
Overweight/Obese (≥ 25 kg/m <sup>2</sup> )		1.87 (1.06, 3.30)	0.55	2.02 (0.99, 4.09)	0.49	1.0		1.84 (1.09, 3.12)	0.56
<b>Preterm-born</b>	6,370								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		1.03 (0.71, 1.51)	0.53	0.75 (0.49, 1.16)	0.72	1.0		1.21 (0.46, 3.18)	0.52
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.06 (0.85, 1.32)	Ref.	1.00 (0.80, 1.24)	Ref.	1.0		1.30 (0.94, 1.81)	Ref.
Overweight/Obese (≥ 25 kg/m <sup>2</sup> )		0.90 (0.59, 1.37)	0.40	0.90 (0.54, 1.49)	0.45	1.0		0.66 (0.45, 0.94)	0.78
<b>Low birthweight</b>	6,205								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		2.95 (1.95, 4.48)	<b>0.03</b>	1.65 (0.97, 2.78)	0.46	1.0		1.49 (0.42, 5.33)	0.76
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.34 (0.97, 1.87)	Ref.	1.08 (0.77, 1.52)	Ref.	1.0		1.41 (0.87, 2.30)	Ref.
Overweight/Obese (≥ 25 kg/m <sup>2</sup> )		0.94 (0.49, 1.78)	0.45	1.13 (0.54, 2.35)	0.28	1.0		0.58 (0.33, 1.02)	0.77
<b>Small-for-gestational age</b>	6,188								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		1.44 (1.07, 1.95)	0.23	1.21 (0.87, 1.67)	0.59	1.0		1.83 (0.72, 4.64)	0.28
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.79 (1.47, 2.18)	Ref.	1.49 (1.22, 1.81)	Ref.	1.0		0.79 (0.56, 1.12)	Ref.
Overweight/Obese (≥ 25 kg/m <sup>2</sup> )		0.75 (0.51, 1.11)	0.40	0.82 (0.52, 1.28)	0.63	1.0		0.93 (0.63, 1.36)	0.97
<b>Large-for-gestational age</b>	6,188								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		0.41 (0.21, 0.77)	0.82	0.72 (0.43, 1.19)	0.27	1.0		0.99 (0.36, 2.71)	0.20
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		0.65 (0.51, 0.83)	Ref.	0.72 (0.57, 0.91)	Ref.	1.0		1.32 (0.95, 1.84)	Ref.
Overweight/Obese (≥ 25 kg/m <sup>2</sup> )		1.20 (0.76, 1.88)	0.08	0.69 (0.37, 1.27)	0.91	1.0		0.87 (0.61, 1.25)	0.44
<b>Stunting at birth</b>	4,590								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		1.12 (0.79, 1.60)	0.91	0.77 (0.52, 1.14)	0.16	1.0		0.60 (0.19, 1.89)	0.29
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.26 (1.02, 1.55)	Ref.	1.23 (1.00, 1.51)	Ref.	1.0		1.03 (0.75, 1.42)	Ref.
Overweight/Obese (≥ 25 kg/m <sup>2</sup> )		1.09 (0.73, 1.62)	0.96	1.30 (0.85, 2.00)	0.46	1.0		1.03 (0.73, 1.44)	0.85
<b>Microcephaly</b>	5,435								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		1.00 (0.60, 1.66)	0.38	1.10 (0.67, 1.82)	0.54	1.0		1.81 (0.49, 6.61)	0.74
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.48 (1.09, 2.01)	Ref.	1.37 (1.01, 1.86)	Ref.	1.0		0.94 (0.56, 1.55)	Ref.
Overweight/Obese (≥ 25 kg/m <sup>2</sup> )		0.90 (0.52, 1.58)	0.32	1.34 (0.75, 2.41)	0.93	1.0		1.15 (0.67, 1.97)	0.73
<b>Macrosomia</b>	6,205								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		0.52 (0.07, 4.20)	0.86	0.52 (0.12, 2.19)	0.80	1.0		0.90 (0.11, 7.38)	0.80
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		0.27 (0.13, 0.56)	Ref.	0.58 (0.34, 1.00)	Ref.	1.0		1.47 (0.78, 2.77)	Ref.
Overweight/Obese (≥ 25 kg/m <sup>2</sup> )		1.61 (0.44, 5.87)	0.60	2.11 (0.84, 5.25)	0.24	1.0		1.29 (0.68, 2.45)	0.71

<sup>1</sup> Multivariable models adjusted for maternal age, education, marital status, wealth quintile, smoking status, alcohol consumption, maternal pre-pregnancy BMI, and supplementation group assignment.

<sup>2</sup> P-value for interaction based on Wald test evaluating the relationship between gestational weight gain adequacy ratio and perinatal outcome among normal weight women is the reference.

**Supplemental Table 2:** Associations between percent adequacy of early gestational weight gain (2<sup>nd</sup> trimester) and perinatal outcomes, modified by maternal pre-pregnancy BMI<sup>1</sup>.

Characteristic	n	Severely Inadequate (<70%)		Inadequate (70% to <90%)		Adequate (90% to 125%)		Excessive (>125%)	
		RR (95% CI)	P int <sup>2</sup>	RR (95% CI)	P int. <sup>2</sup>	RR (95% CI)	P int. <sup>2</sup>	RR (95% CI)	P int. <sup>2</sup>
<b>Stillbirth</b>	7,054								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		0.97 (0.33, 2.83)	0.49	0.71 (0.33, 1.51)	0.72	1.0		1.90 (0.66, 5.44)	0.12
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		0.94 (0.58, 1.53)	Ref.	1.07 (0.72, 1.59)	Ref.	1.0		1.14 (0.64, 2.03)	Ref.
Overweight/Obese ( $\geq$ 25 kg/m <sup>2</sup> )		2.06 (1.00, 4.23)	0.63	0.70 (0.17, 2.91)	0.31	1.0		1.55 (0.88, 2.73)	0.93
<b>Perinatal death</b>	6,750								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		1.36 (0.64, 2.90)	0.65	0.84 (0.47, 1.50)	0.58	1.0		1.26 (0.49, 3.23)	0.79
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		0.94 (0.64, 1.37)	Ref.	0.92 (0.67, 1.26)	Ref.	1.0		1.13 (0.72, 1.78)	Ref.
Overweight/Obese ( $\geq$ 25 kg/m <sup>2</sup> )		2.26 (1.30, 3.93)	0.38	1.80 (0.83, 3.89)	0.80	1.0		1.52 (0.97, 2.37)	0.90
<b>Preterm-born</b>	6,807								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		1.11 (0.71, 1.72)	0.17	0.77 (0.53, 1.11)	0.75	1.0		1.12 (0.59, 2.12)	0.24
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.55 (1.25, 1.91)	Ref.	0.86 (0.71, 1.05)	Ref.	1.0		1.17 (0.89, 1.56)	Ref.
Overweight/Obese ( $\geq$ 25 kg/m <sup>2</sup> )		0.80 (0.54, 1.19)	0.47	<b>2.19 (1.32, 3.62)</b>	<b>0.02</b>	1.0		0.62 (0.45, 0.84)	0.07
<b>Low birthweight</b>	6,579								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		3.26 (1.94, 5.45)	0.02	1.61 (1.05, 2.45)	0.54	1.0		1.63 (0.70, 3.81)	0.64
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.43 (1.03, 2.00)	Ref.	1.16 (0.87, 1.56)	Ref.	1.0		1.26 (0.81, 1.95)	Ref.
Overweight/Obese ( $\geq$ 25 kg/m <sup>2</sup> )		1.05 (0.58, 1.91)	0.94	1.14 (0.48, 2.68)	0.94	1.0		0.68 (0.42, 1.09)	0.23
<b>Small-for-gestational age</b>	6,559								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		1.31 (0.86, 2.00)	1.0	1.34 (1.03, 1.76)	0.92	1.0		1.34 (0.72, 2.50)	0.96
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.28 (1.04, 1.58)	Ref.	1.38 (1.15, 1.64)	Ref.	1.0		0.82 (0.61, 1.10)	Ref.
Overweight/Obese ( $\geq$ 25 kg/m <sup>2</sup> )		0.83 (0.55, 1.25)	0.96	0.57 (0.30, 1.09)	0.32	1.0		0.91 (0.66, 1.26)	0.77
<b>Large-for-gestational age</b>	6,559								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		0.58 (0.32, 1.07)	0.48	0.57 (0.35, 0.93)	0.46	1.0		0.52 (0.22, 1.22)	0.70
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.23 (0.98, 1.56)	Ref.	0.74 (0.59, 0.92)	Ref.	1.0		1.28 (0.95, 1.72)	Ref.
Overweight/Obese ( $\geq$ 25 kg/m <sup>2</sup> )		0.83 (0.54, 1.27)	0.53	1.79 (0.98, 3.26)	0.13	1.0		0.85 (0.63, 1.16)	0.55
<b>Stunting at birth</b>	4,741								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		0.73 (0.42, 1.24)	0.13	1.12 (0.82, 1.52)	0.70	1.0		0.67 (0.34, 1.34)	0.14
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.14 (0.91, 1.44)	Ref.	1.09 (0.90, 1.33)	Ref.	1.0		1.17 (0.88, 1.55)	Ref.
Overweight/Obese ( $\geq$ 25 kg/m <sup>2</sup> )		0.69 (0.44, 1.08)	0.04	0.88 (0.47, 1.65)	0.33	1.0		0.97 (0.72, 1.30)	0.27
<b>Microcephaly</b>	5,615								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		0.79 (0.37, 1.68)	0.43	1.33 (0.86, 2.04)	0.74	1.0		1.35 (0.56, 3.23)	0.79
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		1.09 (0.78, 1.51)	Ref.	1.02 (0.77, 1.35)	Ref.	1.0		0.98 (0.65, 1.49)	Ref.
Overweight/Obese ( $\geq$ 25 kg/m <sup>2</sup> )		0.93 (0.50, 1.75)	0.66	0.50 (0.15, 1.61)	0.22	1.0		1.06 (0.68, 1.64)	0.86
<b>Macrosomia</b>	6,579								
Underweight (BMI <18.5 kg/m <sup>2</sup> )		0.95 (0.11, 7.99)	0.82	0.45 (0.11, 1.91)	0.62	1.0		0.29 (0.04, 2.20)	0.44
Normal weight (18.5 – 24.9 kg/m <sup>2</sup> )		0.35 (0.16, 0.79)	Ref.	0.59 (0.34, 1.00)	Ref.	1.0		2.10 (1.22, 3.62)	Ref.
Overweight/Obese ( $\geq$ 25 kg/m <sup>2</sup> )		1.66 (0.43, 6.36)	0.39	2.29 (0.65, 8.06)	0.20	1.0		0.99 (0.58, 1.67)	0.72

<sup>1</sup> Multivariable models adjusted for maternal age, education, marital status, wealth quintile, smoking status, alcohol consumption, maternal pre-pregnancy BMI, and supplementation group assignment.

<sup>2</sup> P-value for interaction based on Wald test evaluating the relationship between gestational weight gain adequacy ratio and perinatal outcome among normal weight women is the reference.

**Supplemental Table 3:** Associations between percent adequacy of total gestational weight gain and perinatal outcomes, modified by maternal age<sup>1</sup>.

Characteristic	n	Severely Inadequate (<70%)		Inadequate (70% to <90%)		Adequate (90% to 125%)		Excessive (>125%)	
		RR (95% CI)	P int <sup>2</sup>	RR (95% CI)	P int. <sup>2</sup>	RR (95% CI)	P int. <sup>2</sup>	RR (95% CI)	P int. <sup>2</sup>
<b>Stillbirth</b>	6,579								
<20 years		1.24 (0.55, 2.82)	0.76	0.78 (0.32, 1.86)	0.65	1.0		1.85 (0.84, 4.04)	0.32
20 – 29 years		1.03 (0.51, 2.06)	Ref.	1.38 (0.72, 2.65)	Ref.	1.0		1.70 (0.84, 3.48)	Ref.
>30 years		2.12 (1.09, 4.14)	0.61	0.82 (0.40, 1.70)	0.12	1.0		1.23 (0.65, 2.32)	0.46
<b>Perinatal death</b>	6,376								
<20 years		1.06 (0.59, 1.94)	0.65	0.74 (0.37, 1.47)	0.71	1.0		1.57 (0.77, 3.17)	0.22
20 – 29 years		0.90 (0.56, 1.45)	Ref.	0.99 (0.62, 1.58)	Ref.	1.0		0.92 (0.54, 1.56)	Ref.
>30 years		1.42 (0.85, 2.38)	0.44	0.77 (0.44, 1.37)	0.32	1.0		1.64 (0.97, 2.77)	0.24
<b>Preterm-born</b>	6,370								
<20 years		0.93 (0.64, 1.37)	0.049	1.19 (0.79, 1.80)	0.30	1.0		1.66 (0.99, 2.81)	0.89
20 – 29 years		1.25 (0.93, 1.69)	Ref.	1.01 (0.75, 1.38)	Ref.	1.0		1.27 (0.88, 1.84)	Ref.
>30 years		0.75 (0.54, 1.04)	0.9	0.86 (0.62, 1.21)	0.6	1.0		0.71 (0.48, 1.06)	0.74
<b>Low birthweight</b>	6,205								
<20 years		1.00 (0.62, 1.62)	1.0	1.20 (0.69, 2.09)	0.66	1.0		1.18 (0.50, 2.76)	0.77
20 – 29 years		1.84 (1.19, 2.85)	Ref.	1.31 (0.83, 2.09)	Ref.	1.0		1.22 (0.66, 2.22)	Ref.
>30 years		0.88 (0.56, 1.38)	0.36	0.78 (0.47, 1.29)	0.23	1.0		1.26 (0.69, 2.30)	0.85
<b>Small-for-gestational age</b>	6,188								
<20 years		1.15 (0.85, 1.54)	0.15	1.24 (0.89, 1.73)	0.09	1.0		0.87 (0.50, 1.53)	0.83
20 – 29 years		1.71 (1.32, 2.21)	Ref.	1.44 (1.10, 1.87)	Ref.	1.0		0.90 (0.64, 1.28)	Ref.
>30 years		1.04 (0.79, 1.37)	0.97	0.93 (0.70, 1.24)	0.54	1.0		0.87 (0.59, 1.29)	0.43
<b>Large-for-gestational age</b>	6,188								
<20 years		1.06 (0.66, 1.70)	0.30	1.14 (0.67, 1.95)	0.45	1.0		1.61 (0.92, 2.81)	0.80
20 – 29 years		0.85 (0.61, 1.19)	Ref.	0.72 (0.51, 1.02)	Ref.	1.0		1.43 (0.99, 2.08)	Ref.
>30 years		0.60 (0.41, 0.88)	0.14	1.03 (0.71, 1.49)	0.45	1.0		0.85 (0.59, 1.23)	0.95
<b>Stunting at birth</b>	4,590								
<20 years		1.08 (0.75, 1.56)	0.13	0.87 (0.58, 1.28)	0.53	1.0		0.61 (0.35, 1.08)	0.62
20 – 29 years		0.95 (0.72, 1.26)	Ref.	1.10 (0.83, 1.46)	Ref.	1.0		0.99 (0.71, 1.38)	Ref.
>30 years		1.39 (1.03, 1.89)	0.01	0.94 (0.69, 1.28)	0.57	1.0		0.83 (0.59, 1.18)	0.96
<b>Microcephaly</b>	5,435								
<20 years		1.47 (0.92, 2.35)	0.07	0.89 (0.52, 1.54)	0.63	1.0		1.20 (0.58, 2.51)	0.30
20 – 29 years		1.04 (0.68, 1.58)	Ref.	1.19 (0.79, 1.79)	Ref.	1.0		0.89 (0.53, 1.49)	Ref.
>30 years		1.41 (0.91, 2.18)	0.29	1.23 (0.81, 1.85)	0.55	1.0		0.92 (0.53, 1.59)	0.72
<b>Macrosomia</b>	6,205								
<20 years		0.80 (0.16, 3.98)	0.46	1.77 (0.52, 6.01)	0.91	1.0		1.74 (0.58, 5.20)	0.92
20 – 29 years		0.69 (0.27, 1.77)	Ref.	0.76 (0.31, 1.87)	Ref.	1.0		1.81 (0.79, 4.15)	Ref.
>30 years		0.45 (0.13, 1.54)	0.03	1.69 (0.70, 4.08)	0.64	1.0		1.67 (0.85, 3.28)	0.56

<sup>1</sup>Multivariable models adjusted for maternal age, education, marital status, wealth quintile, smoking status, alcohol consumption, maternal pre-pregnancy BMI, and supplementation group assignment.

<sup>2</sup>P-value for interaction based on Wald test evaluating the relationship between gestational weight gain adequacy ratio and perinatal outcome among normal weight women is the reference.

**Supplemental Table 4:** Associations between percent adequacy of early gestational weight gain (2<sup>nd</sup> trimester) and perinatal outcomes, modified by maternal age<sup>1</sup>.

Characteristic	n	Severely Inadequate (<70%)		Inadequate (70% to <90%)		Adequate (90% to 125%)		Excessive (>125%)	
		RR (95% CI)	P int <sup>2</sup>	RR (95% CI)	P int <sup>2</sup>	RR (95% CI)	P int <sup>2</sup>	RR (95% CI)	P int <sup>2</sup>
<b>Stillbirth</b>	7,054								
<20 years		0.69 (0.28, 1.66)	0.96	1.88 (0.96, 3.67)	0.06	1.0		1.23 (0.58, 2.64)	0.31
20 – 29 years		1.06 (0.59, 1.92)	Ref.	0.67 (0.37, 1.20)	Ref.	1.0		0.97 (0.52, 1.81)	Ref.
>30 years		0.96 (0.47, 1.95)	0.94	1.71 (0.94, 3.10)	0.18	1.0		1.38 (0.79, 2.39)	0.38
<b>Perinatal death</b>	6,750								
<20 years		1.07 (0.57, 2.01)	0.34	1.39 (0.81, 2.39)	0.09	1.0		0.94 (0.50, 1.77)	0.51
20 – 29 years		0.86 (0.54, 1.38)	Ref.	0.68 (0.44, 1.05)	Ref.	1.0		0.94 (0.59, 1.50)	Ref.
>30 years		1.25 (0.72, 2.17)	0.41	1.36 (0.84, 2.19)	0.24	1.0		1.11 (0.73, 1.71)	0.56
<b>Preterm-born</b>	6,807								
<20 years		1.02 (0.70, 1.48)	0.22	1.23 (0.87, 1.75)	0.63	1.0		1.26 (0.81, 1.97)	0.73
20 – 29 years		1.56 (1.17, 2.07)	Ref.	0.88 (0.67, 1.15)	Ref.	1.0		1.07 (0.77, 1.47)	Ref.
>30 years		0.89 (0.64, 1.22)	0.33	0.80 (0.59, 1.08)	0.61	1.0		0.64 (0.46, 0.90)	0.60
<b>Low birthweight</b>	6,579								
<20 years		1.05 (0.63, 1.75)	0.94	1.19 (0.75, 1.89)	0.78	1.0		0.94 (0.45, 1.97)	0.75
20 – 29 years		1.60 (1.06, 2.43)	Ref.	1.08 (0.73, 1.59)	Ref.	1.0		0.93 (0.56, 1.55)	Ref.
>30 years		0.95 (0.59, 1.53)	0.68	0.98 (0.64, 1.50)	0.59	1.0		1.26 (0.77, 2.08)	0.21
<b>Small-for-gestational age</b>	6,559								
<20 years		1.09 (0.77, 1.56)	0.41	1.17 (0.88, 1.56)	0.22	1.0		1.09 (0.71, 1.69)	0.47
20 – 29 years		1.24 (0.94, 1.63)	Ref.	1.22 (0.96, 1.56)	Ref.	1.0		0.86 (0.63, 1.16)	Ref.
>30 years		0.92 (0.66, 1.29)	0.9	1.10 (0.85, 1.41)	0.41	1.0		0.89 (0.64, 1.23)	0.77
<b>Large-for-gestational age</b>	6,559								
<20 years		0.94 (0.59, 1.49)	0.75	1.71 (1.13, 2.58)	0.09	1.0		1.54 (0.95, 2.49)	0.22
20 – 29 years		1.27 (0.92, 1.75)	Ref.	0.68 (0.50, 0.94)	Ref.	1.0		1.10 (0.78, 1.55)	Ref.
>30 years		0.75 (0.52, 1.09)	0.91	0.84 (0.59, 1.20)	0.7	1.0		0.81 (0.58, 1.12)	0.84
<b>Stunting at birth</b>	4,741								
<20 years		1.10 (0.72, 1.68)	0.27	0.85 (0.60, 1.20)	0.82	1.0		0.63 (0.39, 1.03)	0.46
20 – 29 years		0.87 (0.64, 1.19)	Ref.	0.92 (0.71, 1.20)	Ref.	1.0		1.05 (0.77, 1.44)	Ref.
>30 years		0.97 (0.67, 1.40)	0.52	1.13 (0.86, 1.50)	0.12	1.0		0.94 (0.70, 1.26)	0.58
<b>Microcephaly</b>	5,615								
<20 years		1.15 (0.66, 2.03)	0.17	1.24 (0.77, 1.99)	0.09	1.0		1.22 (0.65, 2.32)	0.16
20 – 29 years		0.99 (0.64, 1.53)	Ref.	0.87 (0.59, 1.29)	Ref.	1.0		0.85 (0.53, 1.34)	Ref.
>30 years		0.90 (0.52, 1.56)	0.53	1.24 (0.82, 1.86)	0.7	1.0		1.33 (0.84, 2.10)	0.54
<b>Macrosomia</b>	6,579								
<20 years		1.06 (0.20, 5.52)	0.73	1.55 (0.52, 4.69)	0.37	1.0		2.19 (0.87, 5.51)	0.17
20 – 29 years		0.63 (0.22, 1.80)	Ref.	0.77 (0.33, 1.78)	Ref.	1.0		1.58 (0.70, 3.58)	Ref.
>30 years		0.73 (0.19, 2.82)	0.22	1.07 (0.44, 2.58)	0.33	1.0		2.18 (1.20, 3.99)	0.71

<sup>1</sup>Multivariable models adjusted for maternal age, education, marital status, wealth quintile, smoking status, alcohol consumption, maternal pre-pregnancy BMI, and supplementation group assignment.

<sup>2</sup>P-value for interaction based on Wald test evaluating the relationship between gestational weight gain adequacy ratio and perinatal outcome among normal weight women is the reference.

**Supplementary Table 5:** Associations between gestational weight gain (GWG) z-scores, using Intergrowth-21st standards, and perinatal outcomes, among normal weight women<sup>1</sup>.

Characteristic	Total n	No. of cases (%)	GWGz <-2 SD RR (95% CI)	GWGz -2 SD to <-1 SD RR (95% CI)	GWGz -1 SD to <1 SD Reference	GWGz ≥1 SD RR (95% CI)
<b>Total GWG</b>						
Stillbirth	4314	123 (2.85)	1.27 (0.65, 2.48)	0.98 (0.63, 1.54)	1.0	2.51 (1.26, 4.99)
Perinatal death	4165	195 (4.68)	1.06 (0.60, 1.87)	0.94 (0.66, 1.35)	1.0	1.45 (0.73, 2.85)
Preterm-born	4191	630 (15.0)	0.82 (0.57, 1.18)	0.80 (0.65, 0.98)	1.0	1.48 (0.97, 2.25)
Low birthweight	4076	243 (5.96)	1.62 (1.02, 2.55)	0.93 (0.68, 1.29)	1.0	1.43 (0.73, 2.77)
Small-for-gestational age	4066	866 (21)	1.72 (1.31, 2.27)	1.54 (1.30, 1.83)	1.0	0.73 (0.45, 1.18)
Large-for-gestational age	4066	512 (12.6)	0.51 (0.33, 0.81)	0.62 (0.49, 0.79)	1.0	1.52 (0.97, 2.37)
Stunting at birth	2996	797 (27)	1.16 (0.84, 1.59)	1.16 (0.96, 1.41)	1.0	0.64 (0.39, 1.08)
Microcephaly	3564	307 (8.62)	1.46 (0.96, 2.22)	1.39 (1.06, 1.82)	1.0	0.61 (0.26, 1.41)
Macrosomia	4076	83 (2.04)	0.16 (0.02, 1.13)	0.37 (0.18, 0.74)	1.0	1.62 (0.66, 4.01)
<b>Early GWG (2<sup>nd</sup> trimester)</b>						
Stillbirth	4630	145 (3.13)	1.89 (0.95, 3.76)	0.87 (0.45, 1.67)	1.0	1.08 (0.60, 1.95)
Perinatal death	4411	231 (5.24)	1.26 (0.65, 2.42)	0.94 (0.56, 1.57)	1.0	1.22 (0.77, 1.92)
Preterm-born	4485	753 (16.8)	1.51 (1.01, 2.24)	1.58 (1.22, 2.06)	1.0	1.19 (0.90, 1.58)
Low birthweight	4325	286 (6.61)	1.36 (0.75, 2.46)	1.30 (0.85, 1.97)	1.0	1.39 (0.92, 2.10)
Small-for-gestational age	4313	910 (21)	1.05 (0.70, 1.56)	1.10 (0.85, 1.43)	1.0	0.63 (0.47, 0.86)
Large-for-gestational age	4313	586 (13.6)	1.30 (0.83, 2.05)	1.28 (0.94, 1.73)	1.0	1.51 (1.13, 2.01)
Stunting at birth	3094	814 (26)	1.09 (0.72, 1.67)	1.23 (0.92, 1.66)	1.0	1.23 (0.93, 1.62)
Microcephaly	3679	312 (8.48)	1.68 (1.01, 2.81)	0.95 (0.61, 1.50)	1.0	0.89 (0.57, 1.38)
Macrosomia	4325	86 (1.99)	0.82 (0.20, 3.32)	0.46 (0.14, 1.47)	1.0	2.82 (1.65, 4.80)

<sup>1</sup> Multivariable models adjusted for maternal age, education, marital status, wealth quintile, smoking status, alcohol consumption, maternal pre-pregnancy body-mass-index, parity, and supplementation group assignment.