

1 Supplementary material

2 GISAID acknowledgements table

Table S1: Molecular clock model configurations and parameterisation. The term 'clades' correspond to monophyletic groups of either of the four variants of concern (VOC) in the data set (Alpha, Beta, Gamma, Delta) and the stems are the branches leading up to them. Note that 'clock rate' refers to the global clock rates of the strict clock and 'b. rate' is for the background rate.

Model family	Model abbreviation	Parameters	Number of parameters
Strict clock	SC	clock rate	1
Relaxed	UCLN	Num. branches, mean, s.d.	358+2
Relaxed	UGM	Num. branches, mean, shape	358+2
Random local clock	RLC*	b. rate, num. rate changes	variable
Fixed local clock	FLC clades	b. rate, num. clades	1+4
Fixed local clock	FLC stems+clades	b. rate, num. clades & stems	1+4
Fixed local clock	FLC stems only	b. rate, num. stems	1+4
Fixed local clock	FLC shared stem	b. rate, stem rate	1+1
Fixed local clock	FLC shared clade	b. rate, clade rate	1+1
Fixed local clock	FLC shared clades+stems	b. rate, clade & stem rate	1+1




Figure S1: Illustration of molecular clock models considered in hypothetical trees with the four variants of concern (VOCs) and background genetic diversity. SC stands for strict clock and FLC for fixed local clock. Branches are coloured according to the rate assigned in each case, with grey corresponding to the 'background' branches and those in colours being the 'foreground' branches. Model names match those of Supplementary material Table 1.




Figure S2: Violin plots of the number of weeks and expected number of substitutions along VOC stem branches for the FLC shared stems model (panels (A) and (B)) and FLC stems (panels (C) and (D)).