

Supplementary material

Summary of quality assessment criteria for reasonable quality studies

Study design	Assessment tool	Questions from the overall scale number	Criteria to be met
Cohort study	JBI	1	Two groups were similar and recruited from the same population
		2	Exposures were measured similarly to assign exposure status
		6	Were strategies to deal with confounding factors stated
		7	Suicidal behaviour was measured in a valid and reliable way
		9	Follow-up was complete or reasons for loss to follow up were described and explored
Before and After	NIH tool	3	All eligible participants that met the prespecified entry criteria were enrolled
		6	Outcome measures were prespecified, clearly defined, valid, reliable, and assessed consistently across all study participants
		7	The same approach/data source was used for the pre-covid measures of suicidal behaviour as those collected during the pandemic period
		8	There were no differences in level of missing data pre and during the pandemic period
Time series	EPOC RoB Tool	1	Intervention independent of other changes
		2	Intervention unlikely to affect data collection (low or unclear risk)
		5	Missing outcome measures were unlikely to bias the results
Cross sectional	JBI	1	Criteria for inclusion was clearly defined
		2	Study participants and setting were described in detail
		6	Were strategies to deal with confounding factors stated
		7	Suicidal behaviour was measured in a valid and reliable way
Case series	JBI	1	Criteria for inclusion was clearly defined
		3	Valid methods were used for identification of the condition for all participants included in the case series
		4	Consecutive cases were included
		6	Clear reporting of demographics of the participants in the study

JBI – Joanna Briggs Institute; NIH – National Institutes of Health; EPOC – (Cochrane) Effective Practice and Organisation of Care; RoB – risk of bias

Number of cases of suicide and self-harm used to calculate rate ratios

Author (year)	Number of cases* by pandemic period	
	Pre-	During
Thongchuam (2021)	1	7
Fidanci (2021)	187	31
Eray (2021)	23	11
Acharya (2020)	9	4
Sengupta (2020)	33	50
Behara (2021)	105	61
Jhanwar (2020)	3	11
Shrestha (2020)	38	55

* Cases were either suicide deaths or self-harm attempts